

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

ANBEH-D-15-00493-R1

Voice parameters predict sex-specific body morphology in men and women

Katarzyna Pisanski^{a,*}, Benedict C. Jones^b, Bernhard Fink^{c,d}, Jillian J.M. O'Connor^a, Lisa M. DeBruine^b, Susanne Röder^{c,d}, David R. Feinberg^a

^{a)} Department of Psychology, Neuroscience & Behaviour, McMaster University, Canada

^{b)} Institute of Neuroscience and Psychology, University of Glasgow, Scotland

^{c)} Department of Psychology, University of Göttingen, Germany

^{d)} Courant Research Centre Evolution of Social Behavior, University of Göttingen, Germany

Running Title: Voice parameters predict sex-specific body size and shape

Word Count (main text and tables): 7001

* Address correspondence to: Katarzyna Pisanski, pisanska@mcmaster.ca

Department of Psychology, Neuroscience & Behaviour, McMaster University, 1280 Main Street West,
Hamilton, Ontario, Canada, L8S 4K1. Phone: +1 (905) 525-9140, x. 28664. Fax: +1 (905) 529-6225.

1 **Abstract**

2 Studies of several mammalian species confirm that formant frequencies (vocal tract resonances)
3 predict height and weight better than does fundamental frequency (F_0 , perceived as pitch) in same-sex
4 adults due to differential anatomical constraints. However, our recent meta-analysis [Pisanski et al.
5 (2014) *Animal Behaviour*, 95, 85-99] indicated that formants and F_0 could explain no more than 10%
6 and 2% of the variance in human height, respectively, controlling for sex and age. Here, we examined
7 whether other voice parameters, many which are affected by sex hormones, can indicate additional
8 variance in human body size or shape, and whether these relationships differ between the sexes. Using
9 a cross-cultural sample of 700 men and women, we examined relationships among 19 voice parameters
10 (min-max F_0 , mean F_0 , F_0 variability, formant-based vocal tract length estimates, shimmer, jitter,
11 harmonics-to-noise ratio) and 8 indices of body size or shape (height, weight, body-mass-index BMI,
12 hip-, waist- and chest-circumferences, waist-to-hip ratio WHR, chest-to-hip ratio CHR). Our results
13 confirm that formant measures explain the most variance in men's and women's heights and weights,
14 whereas shimmer, jitter, and HNR do not indicate height, weight, or BMI in either sex. In contrast,
15 these perturbation and noise parameters, in addition to F_0 range and variability, explained more
16 variance in body shape than did formants or mean F_0 , particularly among men. Shimmer or jitter
17 explained the most variance in men's hip circumferences (12%) and CHRs (6%) whereas HNR and
18 formants explained the most variance in women's WHRs (11%), and significantly more than in men's
19 WHRs. Our study represents the most comprehensive analysis of vocal indicators of human body size
20 to date and offers a foundation for future research examining the hormonal mechanisms of voice
21 production in humans and perceptual playback experiments.

22 *Keywords:* voice; acoustic communication; sexual selection; formant; fundamental frequency;
23 jitter; shimmer; body size; waist-to-hip ratio; chest-to-hip ratio

24 Many animals use vocalizations to communicate in social contexts. Vocalizations may
25 communicate an animal's motivation state (Morton, 1977) but can also function as indexical cues to
26 identity, sex, and various physical traits (Ghazanfar & Rendall, 2008; Owren, 2011). Bioacoustic
27 analyses suggest that the vocalizations of mammals contain reliable and perpetually salient information
28 about a vocalizer's body size and mass (Ey, Pfefferle, & Fischer, 2007; Pisanski, Fraccaro, Tigue,
29 O'Connor, Röder, et al., 2014a; Taylor & Reby, 2010), and playback experiments suggest that both
30 human and non-human listeners may use vocalizations to gauge the body size of conspecifics (e.g.,
31 humans, *Homo sapiens*: Charlton, Taylor, & Reby, 2013; Pisanski, Fraccaro, Tigue, O'Connor, &
32 Feinberg, 2014b; Rendall, Vokey, & Nemeth, 2007; Smith & Patterson, 2005; red deer, *Cervus*
33 *elaphus*: Charlton, Reby, & McComb, 2007; koalas, *Phascolarctos cinereus*: Charlton, Whisson, &
34 Reby, 2013; rhesus macques, *Macaca mulatta*: Fitch & Fritz, 2006; dogs, *Canis lupus familiaris*:
35 Taylor, Reby, & McComb, 2010).

36 **Vocal Indicators of Body Size**

37 Following the source-filter theory of speech production (Fant, 1960), researchers attempting to
38 uncover which voice parameters may reliably indicate body size in humans and other mammals have
39 focused on two largely independent features of the voice: mean fundamental frequency (F_0 ,
40 produced by vocal fold vibration and perceived as voice pitch) and formant frequencies (produced by
41 filtering of the supralaryngeal vocal tract; Titze, 1994). Among humans, our recent meta-analysis
42 showed that formants predict height and weight more reliably than does F_0 when sex and age are
43 controlled (Pisanski et al., 2014a). This finding supports the prediction that mammalian formants are
44 more anatomically constrained than is F_0 (Fitch, 1994, 2000) and corroborates findings from several
45 other mammalian species (reviewed in Kreiman & Sidtis, 2011). However, the meta-analysis also
46 highlighted that formants could explain no more than 10% of the variance in men's heights whereas
47 mean F_0 explained less than 2%. Formants accounted for even less of the variance in women's heights

48 (6%) whereas mean $F0$ was not significantly correlated with height among women (Pisanski et al.,
49 2014a). Due to the limited number of studies investigating other kinds of voice-body relationships, the
50 meta-analysis did not test whether vocal features other than mean $F0$ or formants could explain
51 additional variance in human body size, and did not examine relationships between the voice and body
52 shape, such as circumference parameters.

53 **Fundamental Frequency Range and Variability**

54 A growing literature suggests that several voice parameters, in addition to formants and mean
55 $F0$, may indicate body size and shape in one sex or the other. These voice parameters include non-
56 mean-based measures of fundamental frequency such as minimum $F0$, maximum $F0$, and $F0$
57 variability (the standard deviation of $F0$, $F0\ sd$) that are sexually dimorphic (Puts, Apicella, &
58 Cardenas, 2012). These source measures indicate the upper and lower range of an individual's voice
59 pitch and the degree to which voice pitch deviates from baseline across an utterance. The standard
60 deviation of men's $F0$ appears to be a particularly reliable indicator of status, correlating negatively
61 with self-reported dominance, reproductive success, and testosterone levels (Hodges-Simeon, Gaulin,
62 & Puts, 2010, 2011). In a cross-cultural study, Puts and colleagues (2012) found that $F0\ sd$ predicted
63 self-reported physical aggression in American men, and was marginally negatively related to arm
64 strength among American but not Hadza men. In that study, however, formants reliably predicted
65 height in both samples of men, whereas $F0\ sd$ did not.

66 **Vocal Perturbation and Noise**

67 Vocal frequency perturbation (jitter), amplitude perturbation (shimmer), and noise (harmonics-
68 to-noise ratio) parameters may also correlate with body size or shape as they relate to the mass and
69 oscillating properties of the vocal folds. Jitter and shimmer measure the mean deviation in voice pitch
70 or amplitude between adjacent cycles, whereas harmonics-to-noise ratio (HNR) measures the relative

71 degree of periodicity to aperiodicity in the voice. A relatively high degree of jitter or shimmer or a low
72 HNR can indicate irregular vocal fold vibration, often caused by laryngeal asymmetry in mass or
73 tension, which can result in vocal breathiness and hoarseness (Buder, 2000). Traditionally these
74 measures have been used by clinicians to assess voice quality in pathological voices (Maryn, Roy, De
75 Bodt, Van Cauwenberge & Corthals, 2009), however several researchers have criticized the validity of
76 jitter and shimmer as reliable indices of voice quality (Hillenbrand, 1987; Maryn et al., 2009; Kreiman
77 & Gerratt, 2005).

78 Linders, Massa, Boersma and Dejonckere (1995) suggested that jitter and body size may be
79 negatively related to the extent that larger, more massive vocal folds may result in a mechanical
80 dampening of vocal fold oscillation, producing a steadier voice pitch (see also Lieberman 1963; Titze,
81 1988). However, vocal fold mass is more closely related to sex hormone levels than to height, where
82 for example pubertal increases in testosterone masculinise and enlarge the vocal folds causing F_0 to
83 drop (Hollien, Green, & Massey, 1994; Prelevic, 2013). Indeed, researchers have long proposed that
84 sex hormones may influence voice perturbation and noise parameters, either by affecting the mass of
85 the vocal folds, or the motor and sensory processes involved in laryngeal control (e.g., Higgins &
86 Saxman, 1989; Silverman & Zimmer, 1978; for more recent work see Gugatschka, Kiesler,
87 Obermayer-Pietsch, Schoekler, Schmid, Groselj-Strele, & Friedrich, 2010; Prelevic, 2013). It follows
88 that jitter, shimmer and HNR may relate to body size and in particular body shape via the shared
89 influence of sex hormones on these vocal properties and on the development and distribution of fat and
90 muscle on the body.

91 Relationships between perturbation or noise parameters and the human body have been
92 examined in only a small number of studies with mixed results. González (2007) found that jitter
93 correlated positively with women's body mass, such that heavier women showed more irregularities in
94 their voice pitch, whereas shimmer and HNR were relatively poor indicators of women's, and even

95 less so men's, heights and weights. In contrast, Linders et al. (1995) reported a negative correlation
96 between jitter and height in prepubescent girls and boys independent of gender, suggesting that before
97 puberty, shorter children show more irregularities in their voice pitch than do taller children. Finally,
98 Hamdan et al. (2012) failed to find relationships between jitter or HNR and body size, but reported
99 weak positive relationships between shimmer and trunk fat or muscle mass in men. The largest same-
100 sex sample among these studies included only 81 individuals (González, 2007), which may be too few
101 to detect various voice-body relationships.

102 **Vocal Indicators of Body Shape**

103 There is some evidence that information about body *shape*, not only height and weight, may be
104 present in the human voice. The principle mechanism linking voice to body shape may be hormonal
105 (Hughes & Gallup, 2008). In addition to affecting voice *F0* and formants, and possibly also
106 perturbation parameters (Abitbol, Abitbol, & Abitbol, 1999; Dabbs & Mallinger, 1999; Lieberman,
107 McCarthy, Hiiemae, & Palmer, 2001), estrogens and androgens affect the circumferences of the waist,
108 hips, and chest and the ratios among them (waist-to-hip ratio, WHR and chest-to-hip ratio, CHR), as
109 well as an individual's body-mass-index or BMI (Blouin, Boivin, & Tchernof, 2008; Derby, Zilber,
110 Brambilla, Morales, & McKinlay, 2006; Evans, Hoffmann, Kalkhoff, & Kissebah, 1983)¹.

111 Similar to physical height, indices of body shape such as WHR and CHR can provide socially
112 relevant information about an individual (Hughes & Gallup, 2008). For instance, body shape predicts a
113 wide range of health-related factors in both sexes, controlling for body mass (Blouin et al., 2008;
114 Larsson et al., 1984; Seidell, 2009). Among women, WHR and BMI are robust predictors of fecundity
115 and correlate with ratings of women's physical attractiveness from photographs (Kaye, Folsom,

¹ These indices of body shape are sexually dimorphic and can vary independently of one another within the same individual. It is also important to note that the distribution of fat and muscle mass on the body that determines body shape is largely independent of the amount of fat and muscle on the body that determines body mass (Singh & Singh, 2011).

116 Prineas, Potter, & Gapstur, 1990; Singh, 1993; Zaadstra et al., 1993). Women with lower WHRs are
117 also rated as having more attractive voices (Hughes, Dispenza, & Gallup, 2009), and listeners are able
118 to gauge women's WHRs from their voices alone (Hughes, Harrison, & Gallup, 2009). Among men,
119 CHR and height positively predict physical attractiveness and reproductive success (Pawlowski,
120 Dunbar, & Lipowicz, 2000; Swami et al., 2007). Like body size, body shape influences mate
121 preferences across a range of human cultures (Pisanski & Feinberg, 2013) and is likely to be important
122 for both mate selection and intersexual competition.

123 Few studies have examined vocal indicators of body shape compared to body size, and again
124 the results of this work are mixed. Early studies examined relationships between principal components
125 of voice and body shape (i.e., factor scores) in small samples of men or women ($n = 26-34$), making
126 interpretation of results difficult. In these studies, Collins (2000) and Bruckert et al. (2006) failed to
127 find relationships between voice and body shape components among men, whereas Collins and Missing
128 (2003) reported that women with higher harmonics (integer multiples of $F0$) had lower scores on a
129 body component comprised of BMI, weight, waist- and hip-circumference. Evans, Neave and Wakelin
130 (2006) reported negative relationships between men's mean $F0$ and their shoulder- and chest-
131 circumferences or shoulder-to-hip ratios, but no relationship between men's $F0$ and shoulder-to-waist
132 or waist-to-hip ratios. More recently, in a sample of 109 women, Vukovic, Feinberg, DeBruine, Smith
133 and Jones (2010) reported negative relationships between women's mean $F0$'s and their BMIs and hip-
134 circumferences, but not waist circumferences or WHRs.

135 **Key Research Questions**

136 The present study addresses the key open research questions: (1) Do voice parameters other
137 than formants and mean $F0$ explain additional variance in men's and women's heights and weights
138 (i.e., body size)? (2) Does any voice parameter explain variance in the circumferences and

139 circumference ratios of the waist, hips, and chest (i.e., body shape)? (3) Do voice parameters explain
140 more variance in the body size or shape of one sex than the other? To answer these questions we
141 examined relationships among 19 voice parameters and 8 indices of body size or shape in a large cross-
142 cultural sample of adult men and women. To our knowledge, our study is the first to examine
143 relationships between body shape and any of the following vocal parameters: minimum $F0$, maximum
144 $F0$, $F0$ variability, jitter, shimmer and HNR. Although the voice-body relationships investigated in this
145 study were chosen on the basis of the theoretical and empirical work reviewed above, the study is
146 exploratory in nature. The principle aim of the study is to offer a comprehensive account of vocal
147 correlates of body size and shape in humans that may help researchers to generate novel testable
148 hypotheses concerning the mechanisms and functions of these relationships, and ultimately allow for a
149 meta-analysis of less commonly studied voice-body relationships.

150 **Methods**

151 **Sample Characteristics**

152 Voice recordings and body measures derived from a total of 700 (N) adults from Canada (n =
153 118 women; 185 men), Scotland (n =235 women, 111 men) and Germany (n =85 women). Age data
154 were available for the Canadian (men: 18.7 ± 1.5 , women: 19 ± 2.3 , range 17-30 years) and German
155 samples (23.1 ± 2.2 , range 19-30 years). Voice recordings and body measures were initially collected
156 for other research; as a result, age data were unavailable for the Scottish sample and only female
157 participants were included in the German sample. All participants were students at local universities
158 who provided written informed consent to participate in the study and all procedures were approved by
159 the research ethics review board.

160 **Voice Recording**

161 Participants were recorded in a sound attenuated chamber using a professional condenser
162 microphone with a cardioid pick-up pattern and at an approximate distance of 5–10 cm. All participants
163 were recorded speaking five vowel sounds. For the Canadian and German samples the five vowels
164 were /a/, /i/, /ɛ/, /o/, and /u/ (International Phonetic Alphabetic notation). For the UK sample the vowels
165 were /eI/, /i/, /aI/, /o/, and /ju/.

166 **Voice Measurement and Analysis**

167 Voice measurements and analyses were performed in Praat (Boersma & Weenink, 2013). For
168 each vocalizer we analyzed 19 voice parameters including minimum and maximum F_0 , mean F_0 , the
169 standard deviation of F_0 ($F_0\ sd$), three perturbation or noise parameters (shimmer, jitter, and
170 harmonics-to-noise ratio, HNR), the first to fourth formants (F_1 - F_4), and several amalgamated
171 formant-based parameters, henceforth termed vocal tract length (VTL) estimates, that included:
172 Average Formant Frequency, F_n (Pisanski & Rendall, 2011); Formant Dispersion, D_f (Fitch, 1997),
173 Formant Position, P_f (Puts et al., 2012); Formant Spacing and Apparent Vocal Tract Length derived
174 from formant spacing, ΔF and $VTL(\Delta F)$ (Reby & McComb, 2003); Apparent Vocal Tract Length
175 derived from mean formants, $VTL(F_i)$ (adapted from Fitch, 1997; see also Titze, 1994); Geometric
176 Mean Formant Frequency, MFF (Smith & Patterson, 2005); and factor scores from a confirmatory
177 factor analysis, CFA (Turner, Walters, Monaghan, & Patterson, 2009). The algorithms used to compute
178 VTL estimates are provided in Pisanski et al. (2014a). All mean voice measurements were taken from
179 the steady-state portion of each of five isolated vowels per vocalizer, averaged within vocalizers, and
180 then within sex to obtain mean values.

181 We measured all F_0 parameters using Praat's autocorrelation algorithm with a search range set
182 to 65-300 Hz for men and 100-600 Hz for women and measured formants F_1 - F_4 using Praat's Burg
183 Linear Predictive Coding algorithm with the initial settings of maximum formant set to 5000 Hz for
184 men and 5500 Hz for women. Formants were first overlaid on a spectrogram and formant number was

185 manually adjusted until the best visual fit of predicted onto observed formants was obtained (Boersma
186 & Weenink, 2013; see Praat user manual, www.praat.org). The fundamental frequency and formant
187 measures we obtained (see Table 1) agree well with weighted population-level averages (Pisanski et al.,
188 2014a). From the mean $F1-F4$ values we computed eight different VTL estimates (F_n , D_f , P_f , ΔF ,
189 $VTL(\Delta F)$, $VTL(F_i)$, MFF, and CFA; see Table 1 for descriptive statistics, and Pisanski et al. (2014a)
190 for additional details and algorithms used to compute VTL estimates).

191 We measured one noise parameter (HNR), five frequency perturbation or jitter parameters
192 (local, local absolute, rap, ppq5, and ddp), and six amplitude perturbation or shimmer parameters
193 (local, local dB, apq3, apq5, apq11, dda) using Praat's cross-correlation algorithm (Table 1; see also
194 Baken & Orlikoff, 2000). The five jitter measures correlated significantly with one another (all $r > 0.43$,
195 all $P < 0.001$), and the five shimmer measures correlated significantly with one another (all $r > 0.88$, all
196 $P < 0.001$). Hence, using principal component analyses, we reduced each set of measures to a single
197 dimension (henceforth termed Jitter and Shimmer) for which 78% and 94% of the variance was
198 explained, respectively.

199 **Body Size and Shape Measurement**

200 We assessed a total of eight body size and shape measures including height, weight, body-
201 mass-index (BMI), hip-circumference, waist-circumference, chest-circumference, waist-to-hip ratio
202 (WHR), and chest-to-hip ratio (CHR) (see Table 1). Height was measured using a stadiometer or metric
203 tape affixed to the wall and weight was measured using an electronic scale. Participant's BMI was
204 computed as $\text{weight (kg)} / \text{height (m)}^2$ (where 18.5 to 24.9 indicates normal weight as defined by the
205 World Health Organization). Circumference measures were taken using metric tape following previous
206 work, i.e., waist-circumference was taken at the narrowest point between the rib cage and iliac crest,
207 hip-circumference was taken at the widest point between the waist and thigh, and chest circumference
208 was taken at the widest point with the tape measure placed under the arm pits and, for women, above

209 the breasts (Evans et al., 2006; Hughes, Dispenza, & Gallup, 2004; Singh, 1993; Vukovic et al., 2010).
210 Participant's WHRs were computed as the ratio of the waist circumference to the hip circumference
211 and CHRs were computed as the ratio of the chest circumference to the hip circumference.

212 Table 1. Means and standard deviations ($M \pm sd$) of individual vocal parameters and individual indices
 213 of body size or shape.

	Men	Women
Voice Parameters ($M \pm sd$)		
<i>F0</i> mean (Hz)	114.16 \pm 17.01	210.52 \pm 21.58
<i>F0</i> min (Hz)	90.03 \pm 17.78	162.76 \pm 40.20
<i>F0</i> max (Hz)	179.19 \pm 65.14	370.67 \pm 146.33
<i>F0</i> <i>sd</i> (Hz)	14.85 \pm 11.16	33.53 \pm 24.19
<i>F1</i> (Hz)	466.61 \pm 45.92	516 \pm 79
<i>F2</i> (Hz)	1520.45 \pm 140.33	1848 \pm 200
<i>F3</i> (Hz)	2592.49 \pm 132.29	3020 \pm 199
<i>F4</i> (Hz)	3493.87 \pm 187.54	4100 \pm 217
<i>F_n</i> (Hz)	2015.99 \pm 93.32	2293.55 \pm 247.80
<i>Df</i> (Hz)	1008.84 \pm 62.40	1194.79 \pm 69.34
<i>Pf</i> (Z(Hz))	-0.58 \pm 0.63	0.30 \pm 0.47
ΔF (Hz)	1010.66 \pm 46.09	1187.16 \pm 66.21
VTL(<i>F_i</i>) (cm)	18.75 \pm 2.10	15.61 \pm 1.63
VTL(ΔF) (cm)	17.35 \pm 0.79	14.79 \pm 0.84
MFF (Hz)	1589.07 \pm 71.37	1847.59 \pm 133.57
Jitter local (%)	0.0958 \pm 0.037	0.0911 \pm 0.05
Jitter local absolute (s)	0.00013 \pm 0.00007	0.00006 \pm 0.00003
Jitter rap (%)	0.0066 \pm 0.004	0.0072 \pm 0.004
Jitter ppq5 (%)	0.0067 \pm 0.004	0.0070 \pm 0.004
Jitter ddp (%)	0.0199 \pm 0.011	0.2146 \pm 0.013
Shimmer local (%)	0.0958 \pm 0.037	0.0911 \pm 0.054
Shimmer local (dB)	0.9438 \pm 0.315	0.8690 \pm 0.454
Shimmer apq3 (%)	0.0416 \pm 0.018	0.0425 \pm 0.026
Shimmer apq5 (%)	0.0587 \pm 0.028	0.0615 \pm 0.044
Shimmer apq11(%)	0.0868 \pm 0.039	0.0920 \pm 0.076
Shimmer dda (%)	0.1249 \pm 0.055	0.1285 \pm 0.076
HNR (dB)	14.15 \pm 3.21	14.73 \pm 4.62
Body Measure ($M \pm sd$)		
Height (cm)	179.34 \pm 7.16	166.37 \pm 6.93
Weight (kg)	74.96 \pm 12.44	63.28 \pm 10.77
BMI	23.27 \pm 3.26	22.87 \pm 3.64
hip circ. (cm)	99.55 \pm 7.87	99.20 \pm 7.69
waist circ. (cm)	83.81 \pm 8.12	74.54 \pm 7.97
chest circ. (cm)	95.87 \pm 8.57	88.67 \pm 7.34
WHR	0.84 \pm 0.06	0.75 \pm 0.05
CHR	0.96 \pm 0.07	0.90 \pm 0.05

214 Abbreviations: F_0 = fundamental frequency; F_1 - F_4 = first to fourth formant; F_n = average formant frequency; MF =
215 geometric mean formant frequency; D_f = formant dispersion; P_f = formant position; ΔF = formant spacing; $VTL(F_i)$ =
216 apparent vocal-tract length derived from mean formants; $VTL(\Delta F)$ = apparent vocal-tract length derived from formant
217 spacing; HNR = harmonics-to-noise-ratio; BMI = body-mass-index; WHR = waist-to-hip ratio; CHR = chest-to-hip
218 ratio; circ. = circumference.
219 a. See Baken & Orlikoff (2000) for detailed description and comparison of different jitter and shimmer measures.

220 **Statistical Analysis**

221 Shapiro-Wilk tests of normality indicated that many individual voice or body parameters were
222 non-normally distributed. Hence, we examined relationships between the 19 individual voice
223 parameters and 8 individual body size or shape parameters using non-parametric Spearman rank
224 correlations (r_s). We then tested whether the strength of various voice-body relationships differed
225 significantly for samples of men and women by transforming correlation coefficients using Fisher's r -
226 to- z transformations and running a series of independent-samples inference tests (Myers & Sirois,
227 2006). As our goal was to examine predictive utility differences in the strength of relationships between
228 sexes, we used the more conservative approach of comparing absolute r_s values (i.e., ignoring the sign
229 of the correlation, which in some cases would have inflated the apparent sex difference). Effect sizes
230 for sex differences in $|r_s|$ are given as Cohen's q (Cohen, 1988). All analyses were conducted for each
231 sex separately and all statistical tests were two-tailed with an alpha of .05.

232 **Results**

233 **Relationships Between Individual Voice and Body Parameters**

234 Correlations between individual voice parameters and indices of body size and shape are
235 reported in Table 2.

236

237 Table 2. Relationships between individual voice parameters and individual indices of body size or shape.

Parameter	Men								Women							
	height <i>n</i> =262	weight <i>n</i> =259	BMI <i>n</i> =259	hip circ. <i>n</i> =100	waist circ. <i>n</i> =100	chest circ. <i>n</i> =100	WHR <i>n</i> =100	CHR <i>n</i> =100	height <i>n</i> =438	weight <i>n</i> =436	BMI <i>n</i> =436	hip circ. <i>n</i> =297	waist circ. <i>n</i> =297	chest circ. <i>n</i> =297	WHR <i>n</i> =297	CHR <i>n</i> =297
<i>F0</i> mean	-.17**	-.09	-.01	-.10	-.08	-.09	.09	.03	-.10*	-.20**	-.16**	-.14*	-.16**	-.20**	-.09	-.11
<i>F0</i> min	-.11†	-.13*	-.08	.06	-.06	-.11	-.11	-.20*	.04	-.03	-.05	.03	.01	-.03	-.04	-.09
<i>F0</i> max	-.15*	-.19*	-.13*	-.33**	-.18†	-.21*	.10	.09	-.07	-.16**	-.14**	-.09†	-.19**	-.16**	-.14*	-.10*
<i>F0</i> <i>sd</i>	-.06	-.11†	-.09	-.31**	-.13	-.13	.14	.15	-.07	-.15**	-.12*	-.13*	-.22**	-.19**	-.15*	-.09
Shimmer	.03	.01	-.01	-.33**	-.12	-.10	.14	.24*	-.05	-.08†	-.06	-.01	-.22**	-.08	-.27**	-.10
Jitter	-.01	-.05	-.06	-.34**	-.15	-.15	.13	.15	-.04	-.08	-.06	-.02	-.22**	-.11*	-.27**	-.12*
HNR ^a	-.08	-.10	-.07	-.01	-.01	-.17	-.07	-.21*	-.11*	-.08	-.05	-.21**	.08	-.11	.33**	.14†
<i>F1</i>	-.15*	-.26**	-.21*	-.24**	-.12	-.18*	.13	.06	-.15**	-.18**	-.11*	-.18**	.06	-.14*	.26**	.06
<i>F2</i>	-.09	-.02	.04	-.14	-.05	-.14	.16	.06	-.15**	-.07	-.01	-.17**	-.09	-.14*	.08	.06
<i>F3</i>	-.21**	-.18**	-.08	-.22*	-.10	-.23*	.17†	.01	-.18**	-.15**	-.07	-.18**	-.16**	-.17**	-.02	.01
<i>F4</i>	-.25**	-.19**	-.08	-.03	-.02	-.07	.10	-.06	-.26**	-.25**	-.14**	-.27**	-.13*	-.22**	.10†	.09
<i>F_n</i>	-.26**	-.18*	-.06	-.14	.04	-.15	.15	-.01	-.22**	-.19**	-.10*	-.25**	-.01	-.18**	.30**	.12*
<i>D_f</i>	-.21**	-.13*	-.02	.04	-.03	-.01	.07	-.08	-.23**	-.22**	-.12*	-.24**	-.16**	-.20**	.03	.08
<i>P_f</i>	-.13*	-.20**	-.15*	-.33**	-.14	-.25*	.24*	.11	-.24**	-.24**	-.13**	-.24**	-.07	-.20**	.17**	.08
ΔF	-.25**	-.18**	-.06	-.09	.03	-.13	.15	-.05	-.24**	-.22**	-.11*	-.26**	-.15**	-.22**	.08	.07
VTL(<i>F_i</i>)	.18**	.12*	.04	-.01	.02	.06	-.03	.07	.16**	.18**	.11*	.17**	-.06	.09†	-.27**	-.11†
VTL(ΔF)	.25**	.18**	.05	.09	.03	.13	-.14	.05	.25**	.22**	.11*	.26**	.15**	.22**	-.08	-.07
MFF	-.23**	-.23**	-.13*	-.23*	-.11	-.22*	.17†	.04	-.23**	-.21**	-.11*	-.23**	-.06	-.18**	.18**	.07
CFA ^a	.27**	.21**	.09	.18†	.07	.17†	-.20*	-.05	-.24**	-.22**	-.11*	-.26**	-.04	-.22*	.20**	.05

239 Abbreviations: F_0 = fundamental frequency; F_1 - F_4 = first to fourth formant; F_n = average formant frequency; MFF = geometric mean formant frequency; D_f = formant
240 dispersion; P_f = formant position; ΔF = formant spacing; $VTL(F_i)$ = apparent vocal-tract length derived from mean formants; $VTL(\Delta F)$ = apparent vocal-tract length
241 derived from formant spacing; CFA = confirmatory factor analysis (factor scores); HNR = harmonics-to-noise-ratio; BMI = body-mass-index; WHR = waist-to-hip
242 ratio; CHR = chest-to-hip ratio; circ. = circumference.
243 Statistical significance of bivariate Spearman's ρ correlations (r_s) is based on a two-tailed t test, where ** $P < 0.01$, * $P < 0.05$, † $P < 0.10$. Significant correlations
244 ($P < 0.05$) are bolded.
245a. Sample sizes for women's HNRs were 185, and for women's CFA scores were 326 (height), 324 (weight, BMI), and 100 (circumference measures, WHR, CHR).

246 **Vocal tract length estimates.**

247 Compared to all other voice parameters, VTL estimates most strongly predicted height and
248 weight within each sex, explaining upwards of 7.3% and 6.7% of the variance in men's heights and
249 weights, respectively, and 6.7% and 6.3% of the variance in women's heights and weights,
250 respectively. The VTL estimates correlated with men's and women's heights and weights more
251 strongly than did mean F_0 (barring F_1 and F_2) replicating the findings of Pisanski et al. (2014a).
252 Although several VTL estimates correlated significantly with men's and women's BMIs, these
253 relationships were weaker than for height or weight.

254 Among women, most VTL estimates also correlated significantly with women's hip-, chest-
255 and waist-circumferences. The VTL estimates were also good predictor's of women's WHRs, wherein
256 F_n explained 9% of the variance in WHRs among women. Despite our large sample size ($n=297$),
257 power analysis ($\alpha=.05$, $\text{power}=.80$) indicated that a sample size of only 85 is required for F_n to
258 reliably predict women's WHRs. Among the VTL estimates, only F_n correlated significantly with
259 women's CHRs, but explained a mere 1.4% of the variance. Compared to women, VTL estimates were
260 relatively poor predictors of men's body shapes. Only F_1 , F_3 , P_f , or MFF correlated significantly with
261 men's hip- and chest-circumferences, only P_f and CFA correlated with men's WHRs, and no VTL
262 estimate correlated with men's waist circumferences or CHRs.

263 **Fundamental frequency parameters.**

264 Mean F_0 , while explaining some variance in height within each sex (2.6% among men and
265 1.9% among women) and in women's weights, BMIs, and circumference measures (up to 4%), did not
266 significantly indicate WHR or CHR in either sex. Rather, F_0 range and variability were better
267 predictor's of body shape than was mean F_0 . Among these F_0 parameters, minimum and maximum F_0
268 significantly predicted men's and women's CHRs, respectively, and maximum F_0 and F_0 sd predicted

269 women's WHRs. Maximum F_0 and F_0 sd also correlated significantly with circumference measures in
270 each sex. Compared to mean F_0 , which did not indicate body shape among men, maximum F_0 and F_0
271 sd explained a noteworthy 11% of the variance in men's hip circumferences.

272 **Vocal perturbation and noise parameters.**

273 Shimmer, jitter, and HNR did not correlate with height, weight, or BMI in either sex, with the
274 exception of HNR that explained a significant but small (1.2%) amount of the variance in women's
275 heights. However, each of these parameters significantly predicted one or more indices of body shape
276 including circumference measures, WHRs and CHRs. Compared to all other voice parameters,
277 shimmer explained the most variance in men's CHRs (5.7%), followed by HNR (4.4%), whereas HNR
278 explained the most variance in women's WHRs (10.8%), followed by F_n (9%), $VTL(F_i)$, jitter and
279 shimmer (each 7.3%).

280 **Sex differences.**

281 Following Fisher's r -to- z transformations controlling for sample size, we tested whether the 19
282 individual voice parameters reported in Table 1 indicated body size or shape significantly better in one
283 sex than the other. Our results indicated that HNR and $VTL(F_i)$ predicted women's WHRs better than
284 men's, and the VTL estimates F_4 and D_f predicted women's hip circumferences better than men's. In
285 contrast, maximum F_0 , shimmer and jitter each predicted men's hip circumferences better than
286 women's (see Table 3). Several trends were observed that did not reach statistical significance, namely:
287 VTL estimates generally explained more variance in men's than women's heights; noise and
288 perturbation parameters explained more variance in men's than women's CHRs; and mean F_0
289 explained more variance in women's than men's weights, BMIs, and circumference measures.

290 Table 3. Significant sex differences in relationships between the voice and body shape

Body shape parameter	Voice measure	<i>z</i>	<i>p</i>	<i>q</i>
Stronger relationships in men:				
Hip circumference	Shimmer	2.84	<.01	.33
	Jitter	2.85	<.01	.33
	<i>F0</i> max	2.16	.03	.25
Stronger relationships in women:				
WHR	HNR	2.33	.02	.27
	VTL(<i>F_i</i>)	2.11	.03	.25
Hip circumference	<i>F4</i>	2.11	.03	.25
	<i>Df</i>	2.43	.01	.29

291

292 Abbreviations: *F0* = fundamental frequency; WHR = waist-to-hip ratio; HNR = harmonics-to-noise-ratio; VTL(*F_i*) =
 293 apparent vocal-tract length derived from mean formants; *F4* = fourth formant; *Df* = formant dispersion.

294 Statistical significance is based on a two-tailed test comparing correlation coefficients following Fisher's *r*-to-*z*
 295 transformations. Only significant relationships (*p*<.05) are shown in this table. Effect sizes are given as Cohen's *q*, and all
 296 fall around the lower threshold of a medium effect size (.30).

297

298

299

300

301

302

General Discussion

303 Our findings demonstrate that the human voice can indicate both body size and shape among
304 adult men and women. We extend the findings of a recent meta-analysis (Pisanski et al., 2014a), and
305 show that in addition to formant frequencies, several other voice parameters including minimum and
306 maximum F_0 , F_0 variability, jitter, shimmer, and HNR may communicate meaningful information
307 about body size or shape. We emphasize, however, that these voice parameters could explain only a
308 small amount of the variation in various indices of human body size and shape controlling for sex and
309 age, and in some cases, the strength of voice-body relationships varied between men and women. Our
310 key findings are discussed in detail below.

311 Vocal Tract Length Estimates

312 Formants are constrained by the length and dimensions of the mammalian vocal tract that in
313 turn is positively related to skull size and height between and within sexes (Fitch, 2000; Fitch & Giedd,
314 1999). Thus, formants predict vocal tract length in many mammals including humans (Fitch & Hauser,
315 2003; Taylor & Reby, 2010). In men, formants also appear to predict circulating levels of testosterone
316 (Bruckert et al., 2006; Evans, Neave, Wakelin, & Hamilton, 2008) that can affect muscularity and the
317 distribution of fat on men's bodies (Blouin et al., 2008), but it remains unknown whether sex hormones
318 affect the formant frequencies of women's voices. Our results indicate that formant-based VTL
319 estimates correlate with men's and women's heights and weights more reliably than other voice
320 parameter investigated in this study. In addition, our study shows that VTL estimates correlate with one
321 or more indices of body shape in each sex, including hip-, waist-, and chest-circumferences, WHRs and
322 CHRs.

323 Formants were particularly robust indicators of body shape among women. One previous study
324 tested but failed to find significant relationships between women's voice parameters (principal

325 components representing F_0 and formants) and body shape. This may be because the authors used
326 factor scores in their regression analyses and a sample of only 30 women (Collins & Missing, 2003).
327 Thus, our study presents the first evidence that formants can explain variation in women's body shapes
328 given an adequate sample size (here, $n=297$, but power analysis suggested that a sample of $n=85$ would
329 suffice). Moreover, many individual VTL estimates explained several times more variance in women's
330 than men's circumference measures and WHRs. Indeed, only four formant measures (F_1 , F_3 , P_f and
331 MFF) significantly predicted variation in men's hip-circumferences and chest-circumferences, and one
332 (CFA) predicted variation in men's WHRs, whereas no voice parameter predicted men's waist-
333 circumferences. Theoretically, the finding that women's voices may carry information about their
334 WHRs is in line with a growing body of literature implicating the 'hour-glass' shape of a woman's
335 body as a key indicator of her age, fertility status, and health (Pisanski & Feinberg, 2013; Singh &
336 Singh, 2011). As an important determinant of her physical attractiveness and desirability as a potential
337 mate, a woman's body shape may be advertised through various modalities, including her voice.
338 However, on a proximate level, the mechanism linking formants to women's WHRs remains unclear
339 and will require a more comprehensive understanding of the relative roles of androgens and estrogens
340 on formant production (particularly among women) and on regional fat distribution.

341 **Fundamental Frequency Parameters**

342 Previous studies have shown that when sex and age are controlled, mean fundamental
343 frequency (F_0 or voice pitch) is a weak predictor of height in humans (Pisanski et al., 2014a) and of
344 body size in many other mammals (for reviews, see Ey et al., 2007; Fitch & Hauser, 2003). The mass
345 and tension of the vocal folds determine mean F_0 (Titze, 1994). However, human vocal folds can
346 develop and grow independently of the rest of the body, as their mass appears more closely related to
347 testosterone levels at puberty and into adulthood than to body size (Dabbs & Mallinger, 1999; Harries,
348 Walker, Williams, Hawkins, & Hughes, 1997).

349 In our study, mean $F0$ predicted weight, BMI and body circumferences only among women,
350 and did not predict WHR or CHR in either sex. Our results suggest that while mean $F0$ may indicate
351 women's body masses, it is a relatively poor predictor of body shape in either sex, generally supporting
352 the results of past work (Bruckert et al., 2006; Collins, 2000; but see controlling for age: Evans et al.,
353 2006). Also in line with our results are those of a meta-analysis that showed a negative relationship
354 between mean $F0$ and weight among women but not men (Pisanski et al. 2014a). One other study
355 reported a significant negative correlation between women's mean $F0$ and factor scores derived from a
356 principal component that included women's weights, BMIs, percentage body fat, waist- and hip-
357 circumferences, and WHRs (Vukovic et al., 2010), however it is difficult to ascertain whether this
358 relationship was driven by body mass, body shape, or both. One possible explanation for the apparent
359 negative relationship between women's mean $F0$ and body mass is that relatively higher levels of
360 androgens and/or lower levels of estrogens may cause some women to develop both more masculine
361 voices (larger vocal folds and lower $F0$, Abitol et al., 1999; Titze, 1994) as well as more masculine
362 bodies (heavier and more muscular, Björntorp, 1991; Blouin et al., 2008). The lack of a relationship
363 between mean $F0$ and body mass in men suggests that the ratio of estrogens to androgens may play a
364 key role in driving this relationship.

365 Although research in other animals indicates that a variety of voice features produced by the
366 vocal source (i.e., the larynx and vocal folds for terrestrial mammals) play a role in acoustic
367 communication (see, e.g., Reby & McComb, 2003; Tyack & Miller, 2002), most human studies
368 examining the indexical functions of voice have focused on mean $F0$ (for reviews, see Feinberg, 2008;
369 Puts, Jones, & DeBruine, 2012b). Our results indicate that $F0$ range and variability are generally better
370 predictors of body shape than is mean $F0$. In particular, minimum $F0$ explained several times more
371 variation in men's CHRs, and maximum $F0$ in men's circumference measures, than did mean $F0$.
372 Indeed, physical height and the girth of the chest relative to the lower body are key predictor's of
373 men's, but not women's, physical attractiveness (Pawlowski et al., 2000; Swami et al., 2007).

374 Similar to Puts et al. (2012), whose study samples included men from the northeastern US and
375 Tanzania, we found that *F0* variability was unrelated to height among men (and women) from three
376 additional cultures. However, our results indicated that *F0 sd* explained 10% of the variance in men's
377 hip-circumferences and correlated negatively with women's weight's, BMIs, circumferences measures
378 and WHRs. Thus, in our study, low *F0* variability indicated larger body circumferences in both sexes
379 and more masculine (lower) WHRs among women. Other recent work investigating *F0* variability in
380 humans suggests that this sexually dimorphic voice parameter may be an important signal of quality.
381 Low *F0* variability produces a perceptually monotone voice that is more common among men than
382 women (Puts et al., 2012), is associated with self-reported physical dominance and reproductive
383 success in men (Hodges-Simeon et al., 2010, 2011), and may also predict circulating levels of
384 testosterone and physical strength (Puts et al., 2012).

385 **Vocal Perturbation and Noise Parameters**

386 To our knowledge, no previous study has investigated relationships between perturbation or
387 noise parameters and body shape. We found that jitter, shimmer and HNR each indicated one or more
388 indices of body shape in both men and women. Shimmer and jitter correlated negatively with hip
389 circumferences among men and with waist- and chest-circumferences among women. Both parameters
390 explained around 12% of the variance in men's hip-circumferences, significantly more than in
391 women's hip-circumferences.

392 Among women, shimmer and jitter correlated negatively with WHR (explaining 7% of the
393 variance) and jitter with CHR, whereas HNR correlated positively with WHR and explained more
394 variance in women's WHRs (11%) than did any other voice parameter. Thus, women with relatively
395 low jitter and shimmer (less perturbation) and high HNRs (less noise) had relatively more masculine
396 body shapes (higher WHRs or CHRs) than did other women. Some researchers have speculated that
397 relationships between perturbation or noise parameters and body shape may be related to sex hormone

398 levels, particularly among women (Linders et al., 1995; Silverman & Zimmer, 1978; Prelevic, 2013).
399 As the vocal folds have androgen receptors that are sensitive to an influx in circulating testosterone,
400 which increases vocal fold mass (Titze, 1994), women with relatively high androgen and/or low
401 estrogen levels may experience a greater increase in vocal fold mass compared to other women. On the
402 basis that larger vocal folds may oscillate with fewer irregularities than smaller vocal folds (Linders et
403 al., 1995), jitter and shimmer may be lower, and HNR higher, among women with more masculine
404 hormonal profiles and more masculine body shapes. Our results support this prediction for women, but
405 not men, among whom shimmer correlated positively, and HNR negatively, with CHR. The possible
406 mechanism linking voice perturbation and noise parameters to men's body shapes is unclear.

407 Our results further indicate that although perturbation and noise parameters predicted body
408 shape, these parameters could not reliably predict height, weight, or BMI in either sex. Previous work
409 has also generally failed to find robust relationships between these parameters and adult height
410 (González, 2007; Hamdan et al., 2012; but see in children: Linders et al., 1995), however two studies
411 reported significant positive relationships between jitter or shimmer and certain indices of body mass
412 including body surface area, trunk fat, or muscle mass (González, 2007; Hamdan et al., 2012). As the
413 results of studies to date are mixed, additional studies are needed to determine whether relatively taller
414 or heavier adults show more irregularities in the pitch and amplitude of their voices than do others.

415 **Limitations and Future Directions**

416 Our voices and bodies change throughout the lifespan, but the most drastic changes occur at
417 puberty and after the age of about 50 (Abitol et al., 1999; Hollien, Green & Massey, 1994). Hence, in
418 the present study, we focused our analyses on adults aged 17-30 years to reduce possible age effects on
419 voice-body relationships. Age data were unavailable for one sample (Scotland), however the sample
420 was comprised of University students whose ages likely fell within this range. To test whether the
421 strength of voice-body relationships changes across the lifespan, future work should include samples

422 with a wider age range, including pre-pubescent children. Moreover, although our study included three
423 large and independent samples of adults from Canada, Scotland and Germany, future replications of
424 this work would also benefit from including multiple other cross-cultural samples, particularly from
425 less industrialized regions of the world.

426 The voice-body relationships reported in this exploratory study warrant replication, particularly
427 those between perturbation or noise parameters and body shape. Reliable measurement of jitter and
428 shimmer is inherently difficult, particularly with voices in which these parameters are high, as their
429 measurement requires accurate identification of cycle boundaries and may also vary as a function of
430 recording hardware, acoustic analysis software, and verbal content (Buder, 2000; Maryn et al., 2009).
431 These parameters can also be difficult to detect and discriminate acoustically (Hillenbrand, 1987;
432 Kreiman & Gerratt, 2005). The average values of jitter, shimmer, and HNR in our samples fell within a
433 non-pathological normal range. However, the mechanisms potentially linking these parameters to body
434 size or shape remain unclear. In our study we have speculated about possible hormonal mechanisms
435 linking variation in the human voice to variation in body size and shape, but we did not measure
436 hormone levels in this study. Future studies may focus on further identifying the hormonal mechanisms
437 of voice production in humans and elucidating the proximate causes of the relationships and sex
438 differences in vocal cues to body size and shape reported here.

439 Unfortunately we cannot infer from our data whether sex differences in vocal cues to body size
440 and shape are the product of sexual selection. It is equally likely, for instance, that men have been
441 selected to exaggerate their body size by lowering the frequencies of their voices (Fitch & Giedd, 1999;
442 Fitch & Reby, 2001; Morton, 1977), and may modulate their formants more than women in ways that
443 reduce the degree to which formants honestly indicate body mass and shape, which are more malleable
444 and variable than is height. At a proximate level, many factors may contribute to sex differences in
445 vocal cues to body size and shape. The vocal tract and resultant formants are sexually dimorphic (Fitch

446 & Giedd, 1999; Titze, 1989). There are also marked sex differences in steroid hormone concentrations
447 and in their effect on vocal anatomy (Abitbol et al., 1999; Lieberman et al., 2001) and on fat
448 distribution (Blouin et al., 2008; Singh, 1993). All or any of these factors may affect the relative degree
449 to which vocal parameters predict variation in body size and shape within and between sexes.

450 Intimately tied to the question of whether reliable indicators of body size and shape are present
451 in the voice is whether listeners are able to accurately gauge size and shape from the voice and, if so,
452 which vocal parameters listeners use to do so. Perceptual studies of voice have generally focused on
453 listener's assessments of height and weight (Bruckert et al., 2006; Collins, 2000; Gonzalez, 2006;
454 Krauss, Freyberg, & Morsella, 2002; Pisanski et al., 2014b; Rendall et al., 2007; Smith & Patterson,
455 2005). Although accuracy in these tasks is generally low, these studies indicate that listeners can gauge
456 height and weight from the voice above chance, and that accuracy is highest (about 60%) in two-
457 alternative forced choice paradigms. Compared to height and weight, relatively few studies have
458 examined assessment of body shape from the voice. Hughes, Harrison and Gallup (2009) found that
459 listeners were able to gauge women's WHRs (but not shoulder-to-hip ratios, SHRs), and men's SHRs
460 (but not WHRs) from the voice alone. Future studies may test whether the sex of the listener affects the
461 accuracy of body shape assessments, as there is some evidence that men are better than women in
462 voice-based assessments of height (Charlton et al., 2013; but see also Rendall et al., 2007). Moreover,
463 while it is clear that listeners utilize both *F*₀ and formant information to gauge height (Charlton et al.,
464 2013; Rendall et al., 2007; Pisanski et al, 2014b), it remains unknown whether listeners use jitter,
465 shimmer or HNR to gauge either body size or shape from the voice. Evidence that listeners are
466 generally insensitive to perturbation parameters in the normal range of variation (Kreiman & Gerratt,
467 2005) suggests that this is unlikely.

468 **Conclusions**

469 We examined relationships among a wide array of voice parameters and indices of both body
470 size and shape in a large cross-cultural sample of men and women. In response to our research
471 questions outlined in the Introduction, our results revealed that: (1) Formants predict height and weight
472 in both men and women better than does any other voice parameter, including mean *F0* and non-mean-
473 based *F0* parameters (*F0* range and variability); (2) Various *F0*, formant, noise and perturbation
474 parameters predicted body shape among men or women. Notably, *F0* range and variability were better
475 predictors of body shape than was mean *F0* in both sexes, formants could explain a similar amount of
476 variance in women's body shapes as in women's body sizes, and jitter, shimmer and HNR were
477 particularly good predictors of body shape including women's WHRs and men's CHRs; (3) Various
478 VTL estimates predicted women's WHRs and hip circumferences significantly better than men's, but
479 shimmer, jitter and maximum *F0* predicted men's hip circumferences better than women's. By
480 informing and guiding future research investigating the mechanisms and functions of vocal
481 communication in humans, these findings may provide further insight into how the human voice and
482 body have been shaped by sexual selection (Puts et al., 2012b), and may offer practical applications for
483 estimating the body size of vocalizers in criminal profiling and remote medical monitoring of obese or
484 malnourished patients.

References

- 485
486
- 487 Abitbol, J., Abitbol, P., & Abitbol, B. (1999). Sex hormones and the female voice. *Journal of Voice*,
488 13(3), 424–446. [http://doi.org/10.1016/S0892-1997\(99\)80048-4](http://doi.org/10.1016/S0892-1997(99)80048-4)
- 489 Baken, R. J., & Orlikoff, R. F. (2000). *Clinical measurement of speech and voice*. San Diego, CA:
490 Singular Publishing Group.
- 491 Björntorp, P. (1991). Adipose tissue distribution and function. *International Journal of Obesity*, 15,
492 67-81.
- 493 Blouin, K., Boivin, A., & Tchernof, A. (2008). Androgens and body fat distribution. *The Journal of*
494 *Steroid Biochemistry and Molecular Biology*, 108(3-5), 272–280.
495 <http://doi.org/10.1016/j.jsbmb.2007.09.001>
- 496 Boersma, P. & Weenink, D. (2013). *Praat: Doing phonetics by computer*. [Computer program].
- 497 Bruckert, L., Liénard, J. S., Lacroix, A., Kreutzer, M., & Leboucher, G. (2006). Women use voice
498 parameters to assess men's characteristics. *Proceedings of the Royal Society Biological*
499 *Sciences Series B*, 273, 83-89. <http://doi.org/10.1098/rspb.2005.3265>
- 500 Buder, E. H. (2000). Acoustic analysis of voice quality: A tabulation of algorithms 1902–1990. In
501 Kent, R. D. & Ball, M. J. (Eds.), *Voice quality measurement* (pp. 119–244). San Diego, CA:
502 Singular Publishing Group.
- 503 Charlton, B. D., Reby, D., & McComb, K. (2007). Female red deer prefer the roars of larger males.
504 *Biology Letters*, 3(4), 382–385. <http://doi.org/10.1098/rsbl.2007.0244>
- 505 Charlton, B. D., Taylor, A. M., & Reby, D. (2013). Are men better than women at acoustic size
506 judgements?. *Biology Letters*, 9, 20130270. <http://doi.org/10.1098/rsbl.2013.0270>

507 Charlton, B. D., Whisson, D. A., & Reby, D. (2013). Free-ranging male koalas use size-related
508 variation in formant frequencies to assess rival males. *PLoS ONE*, *8*(7), e70279.
509 <http://doi.org/10.1371/journal.pone.0070279>

510 Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Mahwah, NJ:
511 Lawrence Erlbaum Associates.

512 Collins, S. A. (2000). Men's voices and women's choices. *Animal Behaviour*, *60*(6), 773–780.
513 <http://doi.org/10.1006/anbe.2000.1523>

514 Collins, S. A., & Missing, C. (2003). Vocal and visual attractiveness are related in women. *Animal*
515 *Behaviour*, *65*(5), 997–1004. <http://doi.org/10.1006/anbe.2003.2123>

516 Dabbs, J. M., & Mallinger, A. (1999). High testosterone levels predict low voice pitch among men.
517 *Personality and Individual Differences*, *27*(4), 801–804. <http://doi.org/10.1016/S0191->
518 [8869\(98\)00272-4](http://doi.org/10.1016/S0191-8869(98)00272-4)

519 Derby, C. A., Zilber, S., Brambilla, D., Morales, K. H., & McKinlay, J. B. (2006). Body mass index,
520 waist circumference and waist to hip ratio and change in sex steroid hormones: the
521 Massachusetts male ageing study. *Clinical Endocrinology*, *65*(1), 125–131.
522 <http://doi.org/10.1111/j.1365-2265.2006.02560.x>

523 Evans, D. J., Hoffmann, R. G., Kalkhoff, R. K., & Kissebah, A. H. (1983). Relationship of androgenic
524 activity to body fat topography, fat cell morphology, and metabolic aberrations in
525 premenopausal women. *The Journal of Clinical Endocrinology & Metabolism*, *57*(2), 304–
526 310. <http://doi.org/10.1210/jcem-57-2-304>

527 Evans, S., Neave, N., & Wakelin, D. (2006). Relationships between vocal characteristics and body
528 size and shape in human males: An evolutionary explanation for a deep male voice.
529 *Biological Psychology*, *72*(2), 160–163. <http://doi.org/10.1016/j.biopsycho.2005.09.003>

- 530 Evans, S., Neave, N., Wakelin, D., & Hamilton, C. (2008). The relationship between testosterone
531 and vocal frequencies in human males. *Physiology & Behavior*, *93*(4-5), 783–788.
532 <http://doi.org/10.1016/j.physbeh.2007.11.033>
- 533 Ey, E., Pfefferle, D., & Fischer, J. (2007). Do age- and sex-related variations reliably reflect body
534 size in non-human primate vocalizations? A review. *Primates*, *48*(4), 253–267.
535 <http://doi.org/10.1007/s10329-006-0033-y>
- 536 Fant, G. (1960). *Acoustic theory of speech production*. The Hague: Mouton.
- 537 Feinberg, D. R. (2008). Are human faces and voices ornaments signaling common underlying cues
538 to mate value? *Evolutionary Anthropology*, *17*(2), 112–118.
539 <http://doi.org/10.1002/evan.20166>
- 540 Fitch, W. T. (1994). *Vocal tract length perception and the evolution of language* (Doctoral
541 dissertation). Brown University, Providence, RI.
- 542 Fitch, W. T. (1997). Vocal tract length and formant frequency dispersion correlate with body size
543 in rhesus macaques. *Journal of the Acoustical Society of America*, *102*(2), 1213–1222.
544 <http://doi.org/10.1121/1.421048>
- 545 Fitch, W. T. (2000). The evolution of speech: a comparative review. *Trends in Cognitive Sciences*,
546 *4*(7), 258–267. [http://doi.org/10.1016/S1364-6613\(00\)01494-7](http://doi.org/10.1016/S1364-6613(00)01494-7)
- 547 Fitch, W. T., & Fritz, J. B. (2006). Rhesus macaques spontaneously perceive formants in
548 conspecific vocalizations. *Journal of the Acoustical Society of America*, *120*(4), 2132–2141.
549 <http://doi.org/10.1121/1.2258499>
- 550 Fitch, W. T., & Giedd, J. (1999). Morphology and development of the human vocal tract: A study
551 using magnetic resonance imaging. *Journal of the Acoustical Society of America*, *106*(3),
552 1511–1522. <http://doi.org/10.1121/1.427148>

553 Fitch, W. T., & Hauser, M. (2003). Unpacking "honesty": vertebrate vocal production and the
554 evolution of acoustic signals. In A. Simmons, A. N. Popper, & R. R. Fay (Eds.), *Acoustic*
555 *Communication* (65-137). New York, NY: Springer.

556 Ghazanfar, A. A., & Rendall, D. (2008). Evolution of human vocal production. *Current Biology*,
557 *18*(11), R457–R460. <http://doi.org/10.1016/j.cub.2008.03.030>

558 González, J. (2007). Correlations between speakers' body size and acoustic parameters of voice.
559 *Perceptual and Motor Skills*, *105*(1), 215–220. <http://doi.org/10.2466/pms.105.1.215-220>

560 Gugatschka, M., Kiesler, K., Obermayer-Pietsch, B., Schoekler, B., Schmid, C., Groselj-Strele, A., &
561 Friedrich, G. (2010). Sex hormones and the elderly male voice. *Journal of Voice*, *24*(3), 369-
562 373.

563 Hamdan, A.-L., Al-Barazi, R., Tabri, D., Saade, R., Kutkut, I., Sinno, S., & Nassar, J. (2012).
564 Relationship between acoustic parameters and body mass analysis in young males.
565 *Journal of Voice*, *26*(2), 144–147. <http://doi.org/10.1016/j.jvoice.2011.01.011>

566 Harries, M. L. L., Walker, J. M., Williams, D. M., Hawkins, S., & Hughes, I. A. (1997). Changes in the
567 male voice at puberty. *Archives of Disease in Childhood*, *77*(5), 445–447.
568 <http://doi.org/10.1136/adc.77.5.445>

569 Hodges-Simeon, C. R., Gaulin, S. J. C., & Puts, D. A. (2010). Different vocal parameters predict
570 perceptions of dominance and attractiveness. *Human Nature*, *21*(4), 406–427.
571 <http://doi.org/10.1007/s12110-010-9101-5>

572 Hodges-Simeon, C. R., Gaulin, S. J. C., & Puts, D. A. (2011). Voice correlates of mating success in
573 men: examining “contests” versus “mate choice” modes of sexual selection. *Archives of*
574 *Sexual Behavior*, *40*(3), 551–557. <http://doi.org/10.1007/s10508-010-9625-0>

575 Hollien, H., Green, R., & Massey, K. (1994). Longitudinal research on adolescent voice
576 change in males. *Journal of the Acoustical Society of America*, *96*, 2646–2654.

577 Hughes, S. M., Dispenza, F., & Gallup Jr, G. G. (2004). Ratings of voice attractiveness predict sexual
578 behavior and body configuration. *Evolution and Human Behavior*, 25(5), 295–304.
579 <http://doi.org/10.1016/j.evolhumbehav.2004.06.001>

580 Hughes, S. M., & Gallup Jr, G. G. (2008). Why are we attracted to certain voices? Voice as an evolved
581 medium for the transmission of psychological and biological information. In K. Ivdebski (Ed.),
582 *Emotions in the Human Voice, Volume 2: Clinical Evidence*. California: Plural Publishing, Inc.

583 Hughes, S. M., Harrison, M. A., & Gallup Jr, G. G. (2009). Sex-specific body configurations can be
584 estimated from voice samples. *Journal of Social, Evolutionary, and Cultural Psychology*, 3(4),
585 343–355.

586 Kaye, S. A., Folsom, A. R., Prineas, R. J., Potter, J. D., & Gapstur, S. M. (1990). The association of
587 body fat distribution with lifestyle and reproductive factors in a population study of
588 postmenopausal women. *International Journal of Obesity*, 14(7), 583–591.

589 Krauss, R. M., Freyberg, R., & Morsella, E. (2002). Inferring speakers' physical attributes from their
590 voices. *Journal of Experimental Social Psychology*, 38(6), 618-625.

591 Kreiman, J., & Gerratt, B. R. (2005). Perception of aperiodicity in pathological voice. *Journal of the*
592 *Acoustical Society of America*, 117(4), 2201-2211.

593 Kreiman, J., & Sidtis, D. (2011). *Foundations of voice studies: An interdisciplinary approach to voice*
594 *production and perception*. West Sussex, UK: Wiley-Blackwell.
595 [doi:10.1002/9781444395068](https://doi.org/10.1002/9781444395068)

596 Larsson, B., Svärdsudd, K., Welin, L., Wilhelmsen, L., Björntorp, P., & Tibblin, G. (1984). Abdominal
597 adipose tissue distribution, obesity, and risk of cardiovascular disease and death: 13 year
598 follow up of participants in the study of men born in 1913. *British Medical Journal*,
599 288(6428), 1401–1404.

600 Lieberman, P. (1963). Some acoustic measures of the fundamental periodicity of normal and
601 pathologic larynges. *Journal of the Acoustical Society of America*, 35(3), 344-353.

602 Lieberman, D. E., McCarthy, R. C., Hiiemae, K. M., & Palmer, J. B. (2001). Ontogeny of postnatal
603 hyoid and larynx descent in humans. *Archives of Oral Biology*, 46(2), 117-128.
604 [http://doi.org/10.1016/S0003-9969\(00\)00108-4](http://doi.org/10.1016/S0003-9969(00)00108-4)

605 Linders, B., Massa, G. G., Boersma, B., & Dejonckere, P. H. (1995). Fundamental voice frequency
606 and jitter in girls and boys measured with electroglottography: influence of age and
607 height. *International Journal of Pediatric Otorhinolaryngology*, 33(1), 61-65.
608 [http://doi.org/10.1016/0165-5876\(95\)01197-J](http://doi.org/10.1016/0165-5876(95)01197-J)

609 Maryn, Y., Roy, N., De Bodt, M., Van Cauwenberge, P., & Corthals, P. (2009). Acoustic measurement
610 of overall voice quality: A meta-analysis. *Journal of the Acoustical Society of America*, 126(5),
611 2619-2634.

612 Morton, E. S. (1977). On the occurrence and significance of motivation-structural rules in some bird
613 and mammal sounds. *The American Naturalist*, 111(981), 855-869.

614 Myers, L. and Sirois, M. J. (2006). Spearman correlation coefficients, differences between.
615 *Encyclopedia of Statistical Sciences*, 12. doi: 10.1002/0471667196.ess5050.pub2

616 Nadhuroua, B., Gambiaa, M., Andriaholinirinab, N.V., & Ouledic, A., & Giacomaa, C. (2015). The
617 vocal communication of the mongoose lemur (*Eulemur mongoz*): Phonation mechanisms,
618 acoustic features and quantitative analysis. *Ethology Ecology & Evolution*.
619 <http://dx.doi.org/10.1080/03949370.2015.1039069>

620 Owren, M. J. (2011). Human voice in evolutionary perspective. *Acoustics Today*, 7(4), 24-33.

621 Pawlowski, B., Dunbar, R. I. M., & Lipowicz, A. (2000). Evolutionary fitness: Tall men have more
622 reproductive success. *Nature*, 403(6766), 156-156. <http://doi.org/10.1038/35003107>

- 623 Pisanski, K., & Feinberg, D. (2013). Cross-cultural variation in mate preferences for averageness,
624 symmetry, body size, and masculinity. *Cross-Cultural Research*, 47(2), 162–197.
625 <http://doi.org/10.1177/1069397112471806>
- 626 Pisanski, K., Fraccaro, P. J., Tigue, C. C., O'Connor, J. J. M., & Feinberg, D. R. (2014b). Return to Oz:
627 Voice pitch facilitates assessments of men's body size. *Journal of Experimental Psychology:*
628 *Human Perception and Performance*, 40(4), 1316–1331.
629 <http://doi.org/10.1037/a0036956>
- 630 Pisanski, K., Fraccaro, P. J., Tigue, C. C., O'Connor, J. J. M., Röder, S., Andrews, P. W., ... Feinberg, D.
631 R. (2014a). Vocal indicators of body size in men and women: a meta-analysis. *Animal*
632 *Behaviour*, 95, 89–99. <http://doi.org/10.1016/j.anbehav.2014.06.011>
- 633 Pisanski, K., & Rendall, D. (2011). The prioritization of voice fundamental frequency or formants
634 in listeners' assessments of speaker size, masculinity, and attractiveness. *Journal of the*
635 *Acoustical Society of America*, 129(4), 2201. <http://doi.org/10.1121/1.3552866>
- 636 Prelevic, G. M. (2013). The effects of sex hormones on the female voice. *The Journal of Musical*
637 *Performance*, 2(4), 93-103.
- 638 Sitverman, E. & Zimmer, C. (1978). Effect of the menstrual cycle on voice quality. *Archives of*
639 *Otolaryngology*, 104, 7-10.
- 640 Puts, D. A., Apicella, C. L., & Cardenas, R. A. (2012a). Masculine voices signal men's threat potential
641 in forager and industrial societies. *Proceedings of the Royal Society B: Biological Sciences*,
642 279(1728), 601–609. <http://doi.org/10.1098/rspb.2011.0829>
- 643 Puts, D. A., Jones, B. C., & DeBruine, L. M. (2012b). Sexual selection on human faces and voices. *The*
644 *Journal of Sex Research*, 49(2-3), 227–243.
645 <http://doi.org/10.1080/00224499.2012.658924>

- 646 Reby, D., & McComb, K. (2003). Anatomical constraints generate honesty: acoustic cues to age
647 and weight in the roars of red deer stags. *Animal Behaviour*, *65*(3), 519–530.
648 <http://doi.org/10.1006/anbe.2003.2078>
- 649 Rendall, D., Vokey, J. R., & Nemeth, C. (2007). Lifting the curtain on the Wizard of Oz: Biased
650 voice-based impressions of speaker size. *Journal of Experimental Psychology: Human*
651 *Perception and Performance*, *33*(5), 1208–1219. <http://doi.org/10.1037/0096->
652 [1523.33.5.1208](http://doi.org/10.1037/0096-1523.33.5.1208)
- 653 Seidell, J. C. (2009). Waist circumference and waist/hip ratio in relation to all-cause mortality,
654 cancer and sleep apnea. *European Journal of Clinical Nutrition*, *64*(1), 35–41.
655 <http://doi.org/10.1038/ejcn.2009.71>
- 656 Silverman, E. M., & Zimmer, C. H. (1978). Effect of the menstrual cycle on voice quality. *Archives of*
657 *Otolaryngology*, *104*(1), 7–10.
- 658 Singh, D. (1993). Adaptive significance of female physical attractiveness: Role of waist-to-hip
659 ratio. *Journal of Personality and Social Psychology*, *65*(2), 293–307.
660 <http://doi.org/10.1037/0022-3514.65.2.293>
- 661 Singh, D., & Singh, D. (2011). Shape and significance of feminine beauty: an evolutionary
662 perspective. *Sex Roles*, *64*(9–10), 723–731. <http://doi.org/10.1007/s11199-011-9938-z>
- 663 Smith, D. R. R., & Patterson, R. D. (2005). The interaction of glottal-pulse rate and vocal-tract
664 length in judgments of speaker size, sex, and age. *Journal of the Acoustical Society of*
665 *America*, *118*(5), 3177–3186. <http://doi.org/10.1121/1.2047107>
- 666 Swami, V., Smith, J., Tsiokris, A., Georgiades, C., Sangareau, Y., Tovée, M. J., & Furnham, A. (2007).
667 Male physical attractiveness in Britain and Greece: A cross-cultural study. *The Journal of*
668 *Social Psychology*, *147*(1), 15–26. <http://doi.org/10.3200/SOCP.147.1.15-26>

669 Taylor, A. M., & Reby, D. (2010). The contribution of source-filter theory to mammal vocal
670 communication research: Advances in vocal communication research. *Journal of Zoology*,
671 280(3), 221–236. <http://doi.org/10.1111/j.1469-7998.2009.00661.x>

672 Taylor, A. M., Reby, D., & McComb, K. (2010). Size communication in domestic dog, *Canis*
673 *familiaris*, growls. *Animal Behaviour*, 79(1), 205–210.
674 <http://doi.org/10.1016/j.anbehav.2009.10.030>

675 Titze, I. R. (1988). The physics of small-amplitude oscillation of the vocal folds. *Journal of the*
676 *Acoustical Society of America*, 83(4), 1536-1552.

677 Titze, I. R. (1989). Physiologic and acoustic differences between male and female voices. *Journal*
678 *of the Acoustical Society of America*, 85(4), 1699–1707. <http://doi.org/10.1121/1.397959>

679 Titze, I. R. (1994). *Principles of vocal production*. Englewood Cliffs: Prentice-Hall.

680 Turner, R. E., Walters, T. C., Monaghan, J. J. M., & Patterson, R. D. (2009). A statistical, formant-
681 pattern model for segregating vowel type and vocal-tract length in developmental formant
682 data. *Journal of the Acoustical Society of America*, 125(4), 2374–2386.
683 <http://doi.org/10.1121/1.3079772>

684 Tyack, P. L., & Miller, E. H. (2002). Vocal anatomy, acoustic communication and echolocation. In
685 *Marine Mammal Biology: An Evolutionary Approach* (pp. 142–184). Oxford, England:
686 Blackwell Science.

687 Vukovic, J., Feinberg, D., DeBruine, L., Smith, F., & Jones, B. (2010). Women’s voice pitch is
688 negatively correlated with health risk factors. *Journal of Evolutionary Psychology*, 8(3),
689 217–225. <http://doi.org/10.1556/JEP.8.2010.3.2>

690 Wadewitz, P., Hammerschmidt, K., Battaglia, D., Witt, A., Wolf, F., & Fischer, J. (2015).
691 Characterizing vocal repertoires-hard vs. soft classification approaches. *PloS One*, 10(4),
692 e0125785. <http://doi.org/10.1371/journal.pone.0125785>

693 Zaadstra, B. M., Seidell, J. C., Van Noord, P. A., te Velde, E. R., Habbema, J. D., Vrieswijk, B., &
694 Karbaat, J. (1993). Fat and female fecundity: Prospective study of effect of body fat
695 distribution on conception rates. *British Medical Journal*, 306(6876), 484–487.
696 <http://dx.doi.org/10.1136/bmj.306.6876.484>

Acknowledgements

1

2

3

4

5

6

7

8

9

We thank Diana Borak, Michael Burt, R. Elisabeth Cornwell, Lisa De Couto, Paul J. Fraccaro, Anthony Little, Fiona Moore, David Perrett, Miriam Law Smith, Cara C. Tigue, and Michael Stirrat for assistance with data collection, Stephanie Wu for assistance with voice measurements, and Paul W. Andrews for statistical aid. Portions of this work were supported by grants from the Social Sciences and Humanities Research Council of Canada, the Canadian Foundation for Innovation, the European Research Council (Starting Grant 282655, OCMATE), Ontario's Ministry of Research and Innovation (Early Researcher Award Program ER11-08-084) and the German Research Foundation.