

An Algebraic Theory of Markov Processes

Giorgio Bacci
Aalborg University, Denmark

Prakash Panangaden
McGill University, Montreal, Canada

Radu Mardare
Aalborg University, Denmark

Gordon Plotkin
University of Edinburgh, UK

Abstract

Markov processes are a fundamental model of probabilistic transition systems and are the underlying semantics of probabilistic programs. We give an algebraic axiomatization of Markov processes using the framework of quantitative equational logic introduced in [14]. We present the theory in a structured way using work of Hyland et al. [10] on combining monads. We take the interpolative barycentric algebras of [14] which captures the Kantorovich metric and combine it with a theory of contractive operators to give the required axiomatization of Markov processes both for discrete and continuous state spaces. This work apart from its intrinsic interest shows how one can extend the general notion of combining effects to the quantitative setting.

Keywords Markov processes, equational logic, quantitative reasoning, combining monads

1 Introduction

The theory of effects began with the pioneering work of Moggi [16, 17] on an algebraic treatment of programming languages via the theory of monads. This allowed a compositional treatment of various semantic phenomena such as state, IO, exceptions etc. This work was followed up by the program of Plotkin and Power [19, 20] on understanding the monads as arising from operations and equations; see also the survey of Hyland and Power [11]. A fundamental contribution, due to Hyland et al. [10], was a way of combining effects by taking the “sum” of theories.

In the present paper we use the framework of [14] which introduced the quantitative analogue of equational logic and the techniques of [10] to develop an algebraic theory of Markov processes. In [14] it was shown how a certain set of equations gave as free algebras the space of probability distributions with the Kantorovich metric. A challenge at the time was to extend this to the theory of Markov processes, which are dynamically evolving probability distributions. Instead of developing an equational theory in an *ad-hoc* way, we use the ideas of [10] to obtain a very general theory of probability distributions equipped with additional operators. Markov processes (or labelled Markov processes [18]) are just a very special instance of these where there is a set of unary operators for the transitions.

It is very pleasing that one can obtain the axiomatisation of Markov processes in this systematic way. Some effort is involved in showing that the techniques apply to the quantitative setting; in that sense our results go beyond the example of Markov processes as they can be seen as an example of a general paradigm of forming sums of quantitative theories. Overall, we see our work as a first step towards a full-blown theory of quantitative effects.

The main conceptual advance we feel we have attained is unifying an algebraic presentation of Markov processes with their well-known coalgebraic presentation. On the one hand, one sees them as algebras arising from a very natural quantitative theory; on the other hand, they arise from the theory of quantitative bisimulation via final coalgebras of behaviour functors. This all comes about as we have a coincidence of initial and final coalgebras. Such a coincidence is known in domain theory [21, 22] but seems not to have been developed in the metric case.

As far as we are aware, the algebraic and coalgebraic viewpoints in semantics have largely developed independently. We feel our work contributes to building a bridge between the two and we hope in future work to use this unified perspective in applications.

Technical summary

In [14] it is shown that any quantitative equational theory \mathcal{U} induces a monad $T_{\mathcal{U}}$ on \mathbf{Met} (the category of metric spaces), namely the monad assigning to an arbitrary space M the quantitative algebra freely generated over M and satisfying the quantitative inferences (conditional equations) of \mathcal{U} . One can readily show that if one considers a signature Σ and the empty theory, the induced monad is the free monad Σ^* over the signature endofunctor (also called Σ) in \mathbf{Met} .

Similarly, suppose that with each operator $f \in \Sigma$ of arity n we associate a contractive factor $0 < c < 1$ (written $f: \langle n, c \rangle \in \Sigma$) and add, for each $\delta \geq c\epsilon$, the axiom

$$\{x_1 =_{\epsilon} y_1, \dots, x_n =_{\epsilon} y_n\} \vdash f(x_1, \dots, x_n) \equiv_{\delta} f(y_1, \dots, y_n)$$

obtaining the *quantitative theory of contractive operators* $\mathcal{O}(\Sigma)$. Then the induced monad is the free monad $\tilde{\Sigma}^*$ on the endofunctor $\tilde{\Sigma} = \prod_{f: \langle n, c \rangle \in \Sigma} c \cdot Id^n$ (where $c \cdot X$ is the space X with metric rescaled by a factor of c).

In [14], the monad induced by the quantitative equational theory \mathcal{B} of interpolative barycentric algebras was shown to be the monad Π of finitely supported Borel probability measures with Kantorovich metric. By taking the (disjoint) union of the axioms of interpolative barycentric algebras and of the algebra of contractive operators for Σ , one obtains $\mathcal{B} + \mathcal{O}(\Sigma)$, the *quantitative theory of interpolative barycentric algebras with contractive operators* in Σ .

We show that the free monad induced by $\mathcal{B} + \mathcal{O}(\Sigma)$ is isomorphic to the sum of monads $\tilde{\Sigma}^* + \Pi$. Because of this characterisation, by using results in [10], we can show that $T_{\mathcal{B} + \mathcal{O}(\Sigma)}$ assigns to an arbitrary metric space M the *initial* solution of the functorial equation

$$X_M \cong \Pi(\tilde{\Sigma}X_M + M).$$

We obtain analogous results for complete separable metric spaces by taking the completion of the monad. In this case the monad assigns to any complete separable metric space M the *unique solution* of the functorial equation

$$Y_M \cong \Delta(\tilde{\Sigma}Y_M + M).$$

where Δ is the Giry monad of Borel probability measures with Kantorovich metric.

By observing that the maps from left to right of the above isomorphisms are coalgebra structures, we can algebraically recover Markov processes by using as the signature Σ which has a constant symbol $\mathbf{0}$, representing termination, and a unary operator $\diamond(t)$, representing the capability of performing a transition to t .

The above findings fit into a more general pattern: we show that under certain assumptions on the quantitative theories \mathcal{U} , \mathcal{U}' , the free monad $T_{\mathcal{U}+\mathcal{U}'}$ that arises from the disjoint union $\mathcal{U} + \mathcal{U}'$ of the two theories is the categorical sum $T_{\mathcal{U}} + T_{\mathcal{U}'}$ of the free monads on \mathcal{U} and \mathcal{U}' , respectively. The only requirement on the theories is that they can be axiomatised by a set of quantitative inferences involving only quantitative equations between variables as hypotheses. In [15] this type of theory is called *simple*.

For simple quantitative theories we have another main result: the quantitative algebras satisfying a simple theory \mathcal{U} are in one-to-one correspondence with the Eilenberg-Moore algebras for the free monad $T_{\mathcal{U}}$. This result generalises the classical isomorphism between the algebras of a functor F and the Eilenberg-Moore algebras of the free monad F^* on F [3].

2 Preliminaries

The basic structures with which we work are metric spaces of various kinds. A metric induces a topology on M , and different metrics can induce the same topology. In particular, for any metric d , the 1-bounded function $d'(x, y) =_{\text{def}} d(x, y)/(1 + d(x, y))$ is also a metric and yields the same topology as d . We henceforth restrict to 1-bounded metrics. If a metric space has a countable dense subset we say it is *separable*; this is equivalent to having a countable base for the topology. A sequence (x_i) in a metric space is said to be *Cauchy* if $\forall \epsilon > 0, \exists N, \forall i, j \geq N, d(x_i, x_j) \leq \epsilon$. If every Cauchy sequence converges we say, the space is said to be *complete*. If a space is not complete it can be isometrically embedded in a complete space by adding the limits of Cauchy sequences, this is a standard construction [8]. If (X, d) is a metric space we write (\bar{X}, d) or just \bar{X} for its completion.

Completeness is a metric concept: the same topological space can be described by two different metrics, one complete and the other one not. If, for a given topology, there is *some* metric that is complete we say that the topology is completely metrizable. Topological spaces underlying complete separable metric space are called *Polish*.

The categories of metric spaces that we consider are **Met**: metric spaces, **CMet**: complete metric spaces and **CSMet**: complete separable metric spaces (recall that all the spaces we consider are 1-bounded). The morphisms are the non-expansive maps, i.e. the $f : (X, d_X) \rightarrow (Y, d_Y)$ such that $d_Y(f(x), f(y)) \leq d_X(x, y)$. These categories have all countable products and coproducts. One can define products by taking the set theoretic product and defining the metric to be the sup of the pointwise metrics, i.e. given $\{(M_i, d_i) \mid i \in I\}$ the metric on $\prod_i M_i$ is

$$d((x_1, \dots, x_n, \dots), (y_1, \dots, y_n, \dots)) =_{\text{def}} \sup_i d_i(x_i, y_i).$$

We assume that the reader is familiar with the basic notions of σ -algebras, measurable functions and measures. Given a topology the σ -algebra generated is called its *Borel algebra* and its elements are called *Borel sets*. A probability measure defined on the Borel

sets is a *Borel probability measure*. Given a topological space with its Borel σ -algebra, we define the *support* of a measure to be the complement of the union of all open sets with zero measure. A measure is said to be *finitely supported* if its support is a finite set. A finitely supported probability measure is just a convex sum of point measures; i.e. measures whose support is a single point.

We assume the reader is familiar with monads and with algebras of a monad (the Eilenberg-Moore algebras of a monad).

2.1 Kantorovich Metric

We review some well-known facts about metrics between spaces of probability distributions.

Let M be a metric space. The *Kantorovich metric*¹ between Borel probability measures μ, ν over M is defined as:

$$\mathcal{K}(d_M)(\mu, \nu) = \sup_{f \in \Phi_M} \left| \int f d\mu - \int f d\nu \right|.$$

with supremum ranging over the set Φ_M of positive 1-bounded non-expansive real-valued functions $f : M \rightarrow [0, 1]$.

Under suitable restrictions on the type of measures, the above distance has a well-known dual characterization, based on the notion of *coupling*. A coupling for a pair of Borel probability measures (μ, ν) over M , is a Borel probability measure ω on the product space $M \times M$, such that, for all Borel sets $E \subseteq M$

$$\omega(E \times M) = \mu(E) \quad \text{and} \quad \omega(M \times E) = \nu(E).$$

A Borel probability measure μ over M is *Radon* if for any Borel set $E \subseteq M$, $\mu(E)$ is the supremum of $\mu(K)$ over all compact subsets K of E . We write $C(\mu, \nu)$ for the set of Radon couplings for a pair of Borel probability measures (μ, ν) .

Theorem 2.1 (Kantorovich-Rubinstein Duality [27, Thm. 5.10]). *Let M be a metric space. Then, for arbitrary Radon probability measures μ, ν over M*

$$\mathcal{K}(d_M)(\mu, \nu) = \min \left\{ \int d d\omega \mid \omega \in C(\mu, \nu) \right\}.$$

Examples of Radon probability measures are finitely supported Borel probability measures on any metric space and generic Borel probability measures over complete separable metric spaces.

We write $\Delta(M)$ for the space of Borel probability measures over M with the Kantorovich metric and $\Pi(M)$ for the subspace of $\Delta(M)$ of the finitely supported Borel probability measures over M .

Lemma 2.2. *Let M be a separable metric space. Then, the Cauchy completion of $\Pi(M)$ is isomorphic to the set of Borel probability measures over the Cauchy completion of M , i.e., $\overline{\Pi(M)} \cong \Delta(\bar{M})$.*

3 Quantitative Equational Theories

Quantitative equations were introduced in [14]. In this framework equalities $t \equiv_\epsilon s$ are indexed by a positive rational number, to capture the idea that t is “within ϵ ” of s . This informal notion is formalized in a manner analogous to traditional equational logic and it is shown that one can axiomatize quantitative analogues of algebras. Analogues of Birkhoff’s completeness theorem [14] and variety theorem [15] were established. The collection of equationally defined quantitative algebras form the algebras for monads on suitable categories of metric spaces. In this section we review this formalism.

¹Sometimes called the Wasserstein-1 metric.

Let Σ be an algebraic signature of function symbols $f: n \in \Sigma$ of arity $n \in \mathbb{N}$. Let X be a countable set of variables, ranged over by x, y, z, \dots . We write $\mathbb{T}(\Sigma, X)$ for the set of Σ -terms freely generated over X , ranged over by t, s, u, \dots

A *substitution of type* Σ is a function $\sigma: X \rightarrow \mathbb{T}(\Sigma, X)$ that is homomorphically extended to terms as

$$\sigma(f(t_1, \dots, t_n)) = f(\sigma(t_1), \dots, \sigma(t_n));$$

we write $\mathcal{S}(\Sigma)$ for the set of substitutions of type Σ .

A *quantitative equation of type* Σ over X is an expression of the form $t \equiv_\varepsilon s$, for $t, s \in \mathbb{T}(\Sigma, X)$ and a positive rational number $\varepsilon \in \mathbb{Q}_{>0}$. We write $\mathcal{E}(\Sigma, X)$ for the set of quantitative equations of type Σ over X , and its subsets will be ranged over by $\Gamma, \Theta, \Pi, \dots$

Fix X a countable set of *metavariables*. A *quantitative deduction system of type* Σ is a relation $\vdash \subseteq 2^{\mathcal{E}(\Sigma, X)} \times \mathcal{E}(\Sigma, X)$ from the powerset of $\mathcal{E}(\Sigma, X)$ to $\mathcal{E}(\Sigma, X)$ satisfying the following meta-axioms, for each $f: n \in \Sigma$

$$\text{(RefI)} \quad \emptyset \vdash x \equiv_0 x,$$

$$\text{(Symm)} \quad \{x \equiv_\varepsilon y\} \vdash y \equiv_\varepsilon x,$$

$$\text{(Triang)} \quad \{x \equiv_\varepsilon z, z \equiv_{\varepsilon'} y\} \vdash x \equiv_{\varepsilon+\varepsilon'} y,$$

$$\text{(1-Bdd)} \quad \emptyset \vdash x \equiv_1 y,$$

$$\text{(Max)} \quad \{x \equiv_\varepsilon y\} \vdash x \equiv_{\varepsilon+\varepsilon'} y, \text{ for all } \varepsilon' > 0,$$

$$\text{(Arch)} \quad \{x \equiv_{\varepsilon'} y \mid \varepsilon' > \varepsilon\} \vdash x \equiv_\varepsilon y,$$

$$\text{(f-NE)} \quad \{x_1 =_\varepsilon y_1, \dots, x_n =_\varepsilon y_n\} \vdash f(x_1, \dots, x_n) \equiv_\varepsilon f(y_1, \dots, y_n),$$

$$\text{(Subst)} \quad \text{If } \Gamma \vdash t \equiv_\varepsilon s, \text{ then } \sigma(\Gamma) \vdash \sigma(t) \equiv_\varepsilon \sigma(s), \text{ for all } \sigma \in \mathcal{S}(\Sigma),$$

$$\text{(Cut)} \quad \text{If } \Gamma \vdash \Theta \text{ and } \Theta \vdash t \equiv_\varepsilon s, \text{ then } \Gamma \vdash t \equiv_\varepsilon s,$$

$$\text{(Assum)} \quad \text{If } t \equiv_\varepsilon s \in \Gamma, \text{ then } \Gamma \vdash t \equiv_\varepsilon s,$$

where we write $\Gamma \vdash \Theta$ to mean that $\Gamma \vdash t \equiv_\varepsilon s$ holds for all $t \equiv_\varepsilon s \in \Theta$ and $\sigma(\Gamma) = \{\sigma(t) \equiv_\varepsilon \sigma(s) \mid t \equiv_\varepsilon s \in \Gamma\}$.

The rules (Subst), (Cut), (Assum) are the usual rules of equational logic. The axioms (RefI), (Symm), (Triang) correspond, respectively, to reflexivity, symmetry, and the triangle inequality; (Max) represents inclusion of neighborhoods of increasing diameter; (Arch) is the Archimedean property of the reals w.r.t. a decreasing chain of neighborhoods with converging diameters; and (f-NE) expresses nonexpansiveness of the $f \in \Sigma$. We have added the axiom (1-Bdd) to ensure that the algebras we get also have 1-bounded metrics. This is a minor variation of the theory presented in [14]. The results that we use from that paper all hold with this change.

A *quantitative equational theory of type* Σ over X is a set \mathcal{U} of universally quantified *quantitative inferences* over $\mathcal{E}(\Sigma, X)$, (i.e., expressions of the form

$$\{t_1 \equiv_{\varepsilon_1} s_1, \dots, t_n \equiv_{\varepsilon_n} s_n\} \vdash t \equiv_\varepsilon s,$$

with a finite set of hypotheses) closed under \vdash -deducibility.

A set A of quantitative inferences *axiomatises* a quantitative equational theory \mathcal{U} , if \mathcal{U} is the smallest quantitative equational theory containing A .

The models of quantitative equational theories are universal Σ -algebras equipped with a metric, called *quantitative algebras*.

Definition 3.1. A *quantitative Σ -algebra* is a tuple $\mathcal{A} = (A, \Sigma^{\mathcal{A}})$, where A is a metric space and $\Sigma^{\mathcal{A}} = \{f^{\mathcal{A}}: A^n \rightarrow A \mid f: n \in \Sigma\}$ is a set of non-expansive *interpretations* for the algebraic operators in Σ , i.e., satisfying the following, for all $0 \leq i \leq n$ and $a_i, b_i \in A$,

$$\max_i d_A(a_i, b_i) \geq d_A(f^{\mathcal{A}}(a_1, \dots, a_n), f^{\mathcal{A}}(b_1, \dots, b_n)).$$

The morphisms between quantitative Σ -algebras are non-expansive Σ -homomorphisms. Quantitative Σ -algebras and their morphism form a category $\mathbf{QA}(\Sigma)$.

A quantitative algebra $\mathcal{A} = (A, \Sigma^{\mathcal{A}})$ *satisfies* the quantitative inference $\Gamma \vdash t \equiv_\varepsilon s$ over $\mathcal{E}(\Sigma, X)$, written $\Gamma \models_{\mathcal{A}} t \equiv_\varepsilon s$, if for any assignment $\iota: X \rightarrow A$,

$$(\text{for all } t' \equiv_{\varepsilon'} s' \in \Gamma, d_A(\iota(t'), \iota(s')) \leq \varepsilon') \text{ implies } d_A(\iota(t), \iota(s)) \leq \varepsilon,$$

where $\iota(t)$ is the homomorphic interpretation of $t \in \mathbb{T}(\Sigma, X)$ in \mathcal{A} .

A quantitative algebra \mathcal{A} is said to *satisfy* (or be a *model* for) the quantitative theory \mathcal{U} , if $\Gamma \models_{\mathcal{A}} t \equiv_\varepsilon s$ whenever $\Gamma \vdash t \equiv_\varepsilon s \in \mathcal{U}$. We write $\mathbb{K}(\Sigma, \mathcal{U})$ for the collection of models of a theory \mathcal{U} of type Σ .

Sometimes it is convenient to consider quantitative Σ -algebras whose carrier is a complete metric space. This class of algebras forms a full subcategory of $\mathbf{QA}(\Sigma)$, written $\mathbf{CQA}(\Sigma)$. Similarly, we write $\mathbf{CK}(\Sigma, \mathcal{U})$ for the full subcategory of quantitative Σ -algebras in $\mathbf{CQA}(\Sigma)$ which are models of \mathcal{U} .

The following definition lifts the Cauchy completion of metric spaces to quantitative algebras.

Definition 3.2. The *Cauchy completion of a quantitative Σ -algebra* $\mathcal{A} = (A, \Sigma^{\mathcal{A}})$, is the quantitative Σ -algebra $\overline{\mathcal{A}} = (\overline{A}, \overline{\Sigma^{\mathcal{A}}})$, where \overline{A} is the Cauchy completion of A and $\overline{\Sigma^{\mathcal{A}}} = \{f^{\overline{\mathcal{A}}}: \overline{A}^n \rightarrow \overline{A} \mid f: n \in \Sigma\}$ is such that for Cauchy sequences $(b_j^i)_j$ converging to $b^i \in \overline{A}$, for $1 \leq i \leq n$, we have:

$$f^{\overline{\mathcal{A}}}(b^1, \dots, b^n) = \lim_j f^{\mathcal{A}}(b_j^1, \dots, b_j^n).$$

The above extends to a functor $\mathbb{C}: \mathbf{QA}(\Sigma) \rightarrow \mathbf{CQA}(\Sigma)$ which is the left adjoint to the functor embedding $\mathbf{CQA}(\Sigma)$ into $\mathbf{QA}(\Sigma)$.

The completion of quantitative Σ -algebras extends also to a functor from $\mathbb{K}(\Sigma, \mathcal{U})$ to $\mathbf{CK}(\Sigma, \mathcal{U})$, whenever \mathcal{U} can be axiomatised by a collection of *continuous schemata* of quantitative inferences, i.e., by axiom schemata of the form

$$\{x_1 \equiv_{\varepsilon_1} x_1, \dots, x_n \equiv_{\varepsilon_n} y_n\} \vdash t \equiv_\varepsilon s, \text{ for all } \varepsilon \geq f(\varepsilon_1, \dots, \varepsilon_n),$$

where $f: \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ is a continuous real-valued function, and $x_i, y_i \in X$. We call such a theory *continuous*.

Free Monads on Quantitative Theories. Recall that to every signature Σ , one can associate a *signature endofunctor* (also called Σ) on \mathbf{Met} by:

$$\Sigma = \coprod_{f: n \in \Sigma} Id^n.$$

It is easy to see that, by couniversality of the coproduct, quantitative Σ -algebras correspond to Σ -algebras for the functor Σ in \mathbf{Met} , and the morphisms between them to non-expansive homomorphisms of Σ -algebras. Below we pass between the two points of view as convenient.

In [14] it is shown that any quantitative theory \mathcal{U} of type Σ induces a monad $T_{\mathcal{U}}$ on \mathbf{Met} , called the *free monad on \mathcal{U}* . The relevant results leading to its definition are summarized in the following theorem.

Theorem 3.3 (Free Algebra). *Let \mathcal{U} be a quantitative theory of type Σ . Then, for any $X \in \mathbf{Met}$ there exists a metric space $T_X \in \mathbf{Met}$, a non-expansive map $\eta_X^{\mathcal{U}}: X \rightarrow T_X$, and a quantitative Σ -algebra $(T_X, \psi_X^{\mathcal{U}})$ satisfying \mathcal{U} , such that, for any quantitative Σ -algebra (A, α) in $\mathbb{K}(\Sigma, \mathcal{U})$ and non-expansive map $\beta: X \rightarrow A$, there exists*

a unique homomorphism $h: T_X \rightarrow A$ of quantitative Σ -algebras making the following diagram commute

$$\begin{array}{ccccc} X & \xrightarrow{\eta_X^{\mathcal{U}}} & T_X & \xleftarrow{\psi_X^{\mathcal{U}}} & \Sigma T_X \\ & \searrow \beta & \downarrow h & & \downarrow \Sigma h \\ & & A & \xleftarrow{\alpha} & \Sigma A \end{array}$$

The map h is also called the homomorphic extension of α along β .

The universal property above says that $(T_X, \psi_X^{\mathcal{U}})$ is the free quantitative Σ -algebra for X in $\mathbb{K}(\Sigma, \mathcal{U})$. From Theorem 3.3, one can define the monad $(T_{\mathcal{U}}, \eta^{\mathcal{U}}, \mu^{\mathcal{U}})$ as follows: the functor $T_{\mathcal{U}}: \mathbf{Met} \rightarrow \mathbf{Met}$ associates to $X \in \mathbf{Met}$ the carrier T_X of the free quantitative Σ -algebra for X in $\mathbb{K}(\Sigma, \mathcal{U})$; the maps $\eta_X^{\mathcal{U}}$ form the components of the unit $\eta^{\mathcal{U}}: Id \Rightarrow T_{\mathcal{U}}$; and the multiplication $\mu^{\mathcal{U}}: T_{\mathcal{U}}T_{\mathcal{U}} \Rightarrow T_{\mathcal{U}}$ is defined at X as the unique map that, by Theorem 3.3, satisfies $\mu_X^{\mathcal{U}} \circ \eta_X^{\mathcal{U}} = id$ and $\mu_X^{\mathcal{U}} \circ \psi_{T_{\mathcal{U}}X}^{\mathcal{U}} = \psi_X^{\mathcal{U}} \circ \Sigma \mu_X^{\mathcal{U}}$.

A similar free construction also holds for quantitative algebras in $\mathbf{CQA}(\Sigma)$ for continuous quantitative equational theories:

Theorem 3.4 (Free Complete Algebra). *Let \mathcal{U} be a continuous quantitative theory of type Σ . Then, for any $X \in \mathbf{CMet}$, quantitative Σ -algebra (A, α) in $\mathbf{C}\mathbb{K}(\Sigma, \mathcal{U})$ and non-expansive map $\beta: X \rightarrow A$, there exists a unique homomorphism $h: \mathbf{C}T_{\mathcal{U}}X \rightarrow A$ of quantitative Σ -algebras making the following diagram commute*

$$\begin{array}{ccccc} X & \xrightarrow{\mathbf{C}\eta_X^{\mathcal{U}}} & \mathbf{C}T_{\mathcal{U}}X & \xleftarrow{\mathbf{C}\psi_X^{\mathcal{U}}} & \mathbf{C}\Sigma T_{\mathcal{U}}X \\ & \searrow \beta & \downarrow h & & \downarrow \Sigma h \\ & & A & \xleftarrow{\alpha} & \Sigma A \end{array}$$

The above is equivalent to saying that the forgetful functor from $\mathbf{C}\mathbb{K}(\Sigma, \mathcal{U})$ to \mathbf{CMet} has a left adjoint. In particular, $\mathbf{C}T_{\mathcal{U}}$ is the free monad on \mathcal{U} in \mathbf{CMet} , provided that the quantitative theory is continuous.

4 Disjoint Union of Quantitative Theories

One of the advantages of the approach followed in [10] is that it allows one to combine different computational phenomena in a smooth way. In our setting we need to combine quantitative theories. The major example discussed in this paper is the combination of interpolative barycentric algebras (which we had shown [14] to axiomatize probability distributions with the Kantorovich metric) and the algebras that give a transition structure. This combination gives us the usual theory of Markov processes, but now enriched with metric reasoning principles for the underlying probability distributions.

In this section we develop the theory of the *disjoint union* of quantitative equational theories. Let Σ, Σ' be two disjoint signatures. The disjoint union of two quantitative equational theories $\mathcal{U}, \mathcal{U}'$ of respective types Σ and Σ' , written $\mathcal{U} + \mathcal{U}'$, is the smallest quantitative theory containing \mathcal{U} and \mathcal{U}' . Following Kelly [13], we show that any model for $\mathcal{U} + \mathcal{U}'$ is a $\langle \mathcal{U}, \mathcal{U}' \rangle$ -bialgebra: a metric space A with both a Σ -algebra structure $\alpha: \Sigma A \rightarrow A$ satisfying \mathcal{U} and a Σ' -algebra structure $\beta: \Sigma' A \rightarrow A$ satisfying \mathcal{U}' . Formally, let $\mathbb{K}((\Sigma, \mathcal{U}) \oplus (\Sigma', \mathcal{U}'))$ be the category of $\langle \mathcal{U}, \mathcal{U}' \rangle$ -bialgebras with non-expansive maps that preserve their two algebraic structures. Then, the following isomorphism of categories holds.

Proposition 4.1. $\mathbb{K}(\Sigma + \Sigma', \mathcal{U} + \mathcal{U}') \cong \mathbb{K}((\Sigma, \mathcal{U}) \oplus (\Sigma', \mathcal{U}'))$.

We already saw that any quantitative equational theory induces a free monad on \mathbf{Met} . Next, we would like show that under certain assumptions on the theories, the free monad $T_{\mathcal{U} + \mathcal{U}'}$ on the disjoint union $\mathcal{U} + \mathcal{U}'$ corresponds to the categorical sum $T_{\mathcal{U}} + T_{\mathcal{U}'}$ of the free monads on \mathcal{U} and \mathcal{U}' , respectively.

The only requirement we ask is that the theories can be axiomatised by a set of *simple* quantitative inferences, i.e., inferences having as hypothesis only equations of the form $x \equiv_{\varepsilon} y$, for $x, y \in X$. As in [15], we call these theories *simple*. Note that any continuous quantitative theory is simple.

Let $T\text{-Alg}$ be the category of Eilenberg-Moore T -algebras of a monad T . Then, we have:

Theorem 4.2. *For any simple quantitative equational theory \mathcal{U} of type Σ , $T_{\mathcal{U}}\text{-Alg} \cong \mathbb{K}(\Sigma, \mathcal{U})$.*

Proof. The isomorphism is given by the following pair of functors

$$T_{\mathcal{U}}\text{-Alg} \begin{array}{c} \xrightarrow{H} \\ \xleftarrow{K} \end{array} \mathbb{K}(\Sigma, \mathcal{U})$$

both mapping morphisms essentially to themselves and on objects acting as follows: for $(A, \alpha) \in T_{\mathcal{U}}\text{-Alg}$ and $(B, \beta) \in \mathbb{K}(\Sigma, \mathcal{U})$,

$$H(A, \alpha) = (A, \alpha \circ \psi_A^{\mathcal{U}} \circ \Sigma \eta_A^{\mathcal{U}}), \quad K(B, \beta) = (B, \beta^b),$$

where $\beta^b: T_{\mathcal{U}}B \rightarrow B$ is the unique map that, by Theorem 3.3, satisfies the equations $\beta^b \circ \eta_B^{\mathcal{U}} = id_B$ and $\beta^b \circ \psi_B^{\mathcal{U}} = \beta \circ \Sigma \beta^b$.

To show that K is well defined, we need to prove that the unit and associativity laws for the $T_{\mathcal{U}}$ -algebra hold. The unit law follows directly by definition of β^b . The associativity law follows by Theorem 3.3, since both $\beta^b \circ \mu_B^{\mathcal{U}}$ and $\beta^b \circ T_{\mathcal{U}}\beta^b$ fit as the unique homomorphic extension of β along $\eta_B^{\mathcal{U}}$. Given any morphism $h: (B, \beta) \rightarrow (B', \beta')$ of quantitative Σ -algebras in $\mathbb{K}(\Sigma, \mathcal{U})$, $K(h) = h$ is proved to be a $T_{\mathcal{U}}$ -homomorphism by showing that both $(\beta')^b \circ T_{\mathcal{U}}h$ and $h \circ \beta^b$ fit as the unique homomorphic extension of $\psi_B^{\mathcal{U}}$ along h . Functoriality of K follows similarly, using the universal property in Theorem 3.3.

To show that H is well defined, we need to show that for any $(A, \alpha) \in T_{\mathcal{U}}\text{-Alg}$, $H(A, \alpha)$ satisfies \mathcal{U} . Since, by hypothesis, \mathcal{U} is a simple quantitative theory, it is axiomatised by a set $\mathcal{A} \subseteq \mathcal{U}$ of simple quantitative inferences. Thus, $H(A, \alpha)$ is a model for \mathcal{U} iff it satisfies all the quantitative inference in \mathcal{A} . To this end, note that for any assignment $\iota: X \rightarrow A$ of the metavariables, the following diagram commutes

$$\begin{array}{ccccc} X & \xrightarrow{\eta_X^{\Sigma}} & T(\Sigma, X) & \xleftarrow{\psi_X^{\Sigma}} & \Sigma T(\Sigma, X) \\ \downarrow \iota & \searrow \eta_A^{\mathcal{U}} & \downarrow T_{\Sigma} \iota & & \downarrow \Sigma T_{\Sigma} \iota \\ A & \xrightarrow{\eta_A^{\mathcal{U}}} & T_{\mathcal{U}}A & \xleftarrow{\psi_A^{\mathcal{U}}} & \Sigma T_{\mathcal{U}}A \\ & \searrow id & \downarrow \alpha & & \downarrow \Sigma \alpha \\ & & A & \xleftarrow{\alpha} & T_{\mathcal{U}}A & \xleftarrow{\psi_A^{\mathcal{U}}} & \Sigma T_{\mathcal{U}}A & \xleftarrow{\Sigma \eta_A^{\mathcal{U}}} & \Sigma A \end{array} \quad (1)$$

where the commutativity of the bottom-right square is proven as follows, by naturality of the maps, by the monad laws, and the unit

and associativity laws of $(A, \alpha) \in T_{\mathcal{U}}\text{-Alg}$

$$\begin{array}{ccccc}
 & T_{\mathcal{U}}A & \xleftarrow{\psi_A^{\mathcal{U}}} & \Sigma T_{\mathcal{U}}A & \xleftarrow{id} & \Sigma T_{\mathcal{U}}A \\
 & \swarrow id & \uparrow \mu_A^{\mathcal{U}} & \Sigma \mu_A^{\mathcal{U}} \uparrow & \swarrow \Sigma T_{\mathcal{U}} \eta_A^{\mathcal{U}} & \downarrow id \\
 T_{\mathcal{U}}A & \xleftarrow{T_{\mathcal{U}}\alpha} & T_{\mathcal{U}}T_{\mathcal{U}}A & \xleftarrow{(\psi^{\mathcal{U}}T^{\mathcal{U}})_A} & \Sigma T_{\mathcal{U}}T_{\mathcal{U}}A & \xrightarrow{\Sigma T_{\mathcal{U}}\alpha} & \Sigma T_{\mathcal{U}}A \\
 \alpha \downarrow & & \downarrow \mu_A^{\mathcal{U}} & & \downarrow \Sigma \mu_A^{\mathcal{U}} & & \downarrow \Sigma \alpha \\
 A & \xleftarrow{\alpha} & T_{\mathcal{U}}A & \xleftarrow{\psi_A^{\mathcal{U}}} & \Sigma T_{\mathcal{U}}A & \xrightarrow{\Sigma \alpha} & \Sigma A \\
 & & & \Sigma \eta_A^{\mathcal{U}} \uparrow & \swarrow id & & \\
 & & & \Sigma A & & &
 \end{array}$$

By the commutativity of (1) and uniqueness of homomorphic extension, we have that the homomorphic extension $i^{\sharp}: \mathbb{T}(\Sigma, X) \rightarrow A$ of i on (A, α) can be factorised as $i^{\sharp} = \alpha \circ T_{\mathcal{U}}i$. Moreover $T_{\Sigma}i$ is the homomorphic extension of $\eta_A^{\mathcal{U}} \circ i: X \rightarrow T_{\mathcal{U}}A$ on $(T_{\mathcal{U}}A, \psi_A^{\mathcal{U}})$.

Let $\Gamma \vdash t \equiv_{\varepsilon} s \in \mathcal{A}$ and $i: X \rightarrow A$ be an assignment of the metavariables. Assume that, for all $x \equiv_{\delta} y \in \Gamma$, $d_A(i^{\sharp}(x), i^{\sharp}(y)) \leq \delta$. Since $x, y \in X$, then we have $d_A(i(x), i(y)) \leq \delta$.

By Theorem 3.3, $(T_{\mathcal{U}}A, \psi_A^{\mathcal{U}})$ satisfies \mathcal{U} , so that, because $T_{\Sigma}i$ is the homomorphic extension of $\eta_A^{\mathcal{U}} \circ i: X \rightarrow T_{\mathcal{U}}A$ on $(T_{\mathcal{U}}A, \psi_A^{\mathcal{U}})$, we have that

$$\begin{aligned}
 & \text{(for all } x \equiv_{\delta} y \in \Gamma, d_{T_{\mathcal{U}}A}(\eta_A^{\mathcal{U}} \circ i(x), \eta_A^{\mathcal{U}} \circ i(y)) \leq \delta) \\
 & \quad \text{implies} \\
 & \quad d_{T_{\mathcal{U}}A}(T_{\Sigma}i(t), T_{\Sigma}i(s)) \leq \varepsilon.
 \end{aligned} \tag{2}$$

By definition of $d_{T_{\mathcal{U}}A}$, we have that $d_{T_{\mathcal{U}}A}(\eta_A^{\mathcal{U}} \circ i(x), \eta_A^{\mathcal{U}} \circ i(y)) = d_A(i(x), i(y))$, hence, by (2), we get $d_{T_{\mathcal{U}}A}(T_{\Sigma}i(t), T_{\Sigma}i(s)) \leq \varepsilon$. Thus,

$$\begin{aligned}
 \varepsilon & \geq d_{T_{\mathcal{U}}A}(T_{\Sigma}i(t), T_{\Sigma}i(s)) \\
 & \geq d_A(\alpha \circ T_{\Sigma}i(t), \alpha \circ T_{\Sigma}i(s)) \quad (\alpha \text{ non-expansive}) \\
 & \geq d_A(i^{\sharp}(t), i^{\sharp}(s)). \quad (\alpha \circ T_{\Sigma}i = i^{\sharp})
 \end{aligned}$$

Therefore $H(A, \alpha)$ satisfies \mathcal{U} . Note the crucial role played by the requirement of simple equations. This allows us to have only variables on the left-hand side and we know these are isometrically embedded by η .

It remains to show that K and H are inverses of each other. On morphisms this is clear. As for objects, for $(A, \alpha) \in T_{\mathcal{U}}\text{-Alg}$ and $(B, \beta) \in \mathbb{K}(\Sigma, \mathcal{U})$, we have

$$\begin{aligned}
 KH(A, \alpha) &= (A, (\alpha \circ \psi_A^{\mathcal{U}} \circ \Sigma \eta_A^{\mathcal{U}})^b), \\
 HK(B, \beta) &= (B, \beta^b \circ \psi_B^{\mathcal{U}} \circ \Sigma \eta_B^{\mathcal{U}}),
 \end{aligned}$$

thus, we need to show $\beta^b \circ \psi_B^{\mathcal{U}} \circ \Sigma \eta_B^{\mathcal{U}} = \beta$ and $(\alpha \circ \psi_A^{\mathcal{U}} \circ \Sigma \eta_A^{\mathcal{U}})^b = \alpha$. These are proved by the commutativity of the following diagrams

$$\begin{array}{ccccc}
 & T_{\mathcal{U}}A & \xleftarrow{\psi_A^{\mathcal{U}}} & \Sigma T_{\mathcal{U}}A & \\
 & \swarrow \eta_A^{\mathcal{U}} & & \swarrow \mu_A^{\mathcal{U}} & \swarrow (\theta T^{\mathcal{U}})_A \\
 T_{\mathcal{U}}B & \xleftarrow{\psi_B^{\mathcal{U}}} & \Sigma T_{\mathcal{U}}B & \xleftarrow{\Sigma \eta_B^{\mathcal{U}}} & \Sigma B \\
 \beta^b \downarrow & & \downarrow \Sigma \beta^b & \swarrow id & \\
 B & \xleftarrow{\beta} & \Sigma B & & \\
 & & & \downarrow id & \\
 & & & A & \xleftarrow{\alpha} & T_{\mathcal{U}}A & \xleftarrow{\theta_A} & \Sigma A \\
 & & & \downarrow id & & \downarrow T_{\mathcal{U}}\alpha & & \downarrow \Sigma \alpha
 \end{array}$$

where $\theta = \psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}}$. In particular, the second one proves that α satisfies the equalities $\alpha \circ \eta_A^{\mathcal{U}} = id_A$ and $\alpha \circ \psi_A^{\mathcal{U}} = \alpha \circ \psi_A^{\mathcal{U}} \circ \Sigma \eta_A^{\mathcal{U}} \circ \Sigma \alpha$ which, by Theorem 3.3, are uniquely satisfied by $(\alpha \circ \psi_A^{\mathcal{U}} \circ \Sigma \eta_A^{\mathcal{U}})^b$. Hence, $(\alpha \circ \psi_A^{\mathcal{U}} \circ \Sigma \eta_A^{\mathcal{U}})^b = \alpha$. \square

When the quantitative equational theories $\mathcal{U}, \mathcal{U}'$ are simple, by Theorem 4.2, we get a refinement of Proposition 4.1 as follows.

Let T, G be two monads on a category \mathbf{C} . A $\langle T, G \rangle$ -bialgebra is an object $A \in \mathbf{C}$ with Eilenberg-Moore algebra structures $\alpha: TA \rightarrow A$ and $\beta: GA \rightarrow A$. We write $\langle T, G \rangle$ -**biAlg** for the category of $\langle T, G \rangle$ -bialgebras with morphisms those in \mathbf{C} preserving the two algebraic structures.

Corollary 4.3. *Let $\mathcal{U}, \mathcal{U}'$ be simple quantitative equational theories. Then $\mathbb{K}(\Sigma + \Sigma', \mathcal{U} + \mathcal{U}') \cong \langle T_{\mathcal{U}}, T_{\mathcal{U}'} \rangle$ -**biAlg**.*

Proof. Immediate from Theorem 4.2 and Proposition 4.1. \square

Now we are ready to state the main theorem of the section.

Theorem 4.4. *Let $\mathcal{U}, \mathcal{U}'$ be simple quantitative theories. Then, the monad $T_{\mathcal{U} + \mathcal{U}'}$ in \mathbf{Met} is the sum of monads $T_{\mathcal{U}} + T_{\mathcal{U}'}$.*

Proof. By Corollary 4.3 and Theorem 3.3 the obvious forgetful functor from $\langle T_{\mathcal{U}}, T_{\mathcal{U}'} \rangle$ -**biAlg** to \mathbf{Met} has a left adjoint. The monad generated by this adjunction is isomorphic to $T_{\mathcal{U} + \mathcal{U}'}$. Thus, by [13] (cf. also [1, Proposition 2.8]), the monad $T_{\mathcal{U} + \mathcal{U}'}$ is also isomorphic to $T_{\mathcal{U}} + T_{\mathcal{U}'}$. \square

The above constructions do not use any specific property of the category \mathbf{Met} , apart from requiring its morphisms to be non-expansive. Thus, under mild restrictions on the type of theories and conditions on the free monad induced by them, we can reformulate alternative versions of Theorem 4.4 which are valid on specific full subcategories of \mathbf{Met} .

The first one applies to \mathbf{CMet} , provided that the quantitative equational theories are continuous. Recall that, the disjoint union $\mathcal{U} + \mathcal{U}'$ of two continuous quantitative theories $\mathcal{U}, \mathcal{U}'$ is also continuous, so that, by Theorem 3.4, the free monad on it in \mathbf{CMet} is $\mathbf{CT}_{\mathcal{U} + \mathcal{U}'}$. Moreover, continuous theories are simple.

Theorem 4.5. *Let $\mathcal{U}, \mathcal{U}'$ be continuous quantitative theories. Then, the monad $\mathbf{CT}_{\mathcal{U} + \mathcal{U}'}$ in \mathbf{CMet} is the sum of $\mathbf{CT}_{\mathcal{U}}$ and $\mathbf{CT}_{\mathcal{U}'}$.*

Note that, for a continuous quantitative theory \mathcal{U} , if the functor $T_{\mathcal{U}}$ preserves separability of the metric spaces, then the free monad on \mathcal{U} in \mathbf{CSMet} is given by $\mathbf{CT}_{\mathcal{U}}$. This is the case, for example, for countable signatures (see Lemma A.3 in the appendix). Thus, under an additional condition on the free monads, the theorem above can also be stated for the case of complete separable metric spaces.

Corollary 4.6. *Let $\mathcal{U}, \mathcal{U}'$ be continuous quantitative theories and assume $T_{\mathcal{U} + \mathcal{U}'}, T_{\mathcal{U}}$, and $T_{\mathcal{U}'}$ preserve separability of metric spaces. Then, the monad $\mathbf{CT}_{\mathcal{U} + \mathcal{U}'}$ in \mathbf{CSMet} is the sum of $\mathbf{CT}_{\mathcal{U}}$ and $\mathbf{CT}_{\mathcal{U}'}$.*

5 Interpolative Barycentric Algebras

Interpolative barycentric algebras [14] are the quantitative algebras for the signature

$$\Sigma_{\mathcal{B}} = \{+_e : 2 \mid e \in [0, 1]\}$$

having a binary operator $+_e$, for each $e \in [0, 1]$ (a.k.a. *barycentric signature*), and satisfying the following axioms

$$(B1) \vdash x +_1 y \equiv_0 x,$$

$$(B2) \vdash x +_e x \equiv_0 x,$$

$$(SC) \vdash x +_e y \equiv_0 y +_{1-e} x,$$

$$(SA) \vdash (x +_e y) +_{e'} z \equiv_0 x +_{ee'} (y +_{\frac{e'-ee'}{1-ee'}} z), \text{ for } e, e' \in [0, 1),$$

$$(IB) \{x \equiv_{\varepsilon} y, x' \equiv_{\varepsilon'} y'\} \vdash x +_e x' \equiv_{\delta} y +_e y', \text{ for } \delta \geq e\varepsilon + (1-e)\varepsilon'.$$

The quantitative theory induced by the axioms above, written \mathcal{B} , is called *interpolative barycentric quantitative equational theory*. The axioms (B1), (B2), (SC), (SA) are those of *barycentric algebras* (a.k.a. *abstract convex sets*) due to M. H. Stone [24] where (SC) stands for *skew commutativity* and (SA) for *skew associativity*. (IB) is the *interpolative barycentric axiom* introduced in [14].

5.1 On Metric Spaces

Let $\Pi: \mathbf{Met} \rightarrow \mathbf{Met}$ be the functor assigning to each $X \in \mathbf{Met}$ the metric space $\Pi(X)$ of finitely supported Borel probability measures with Kantorovich metric and acting on morphisms $f \in \mathbf{Met}(X, Y)$ as $\Pi(f)(\mu) = \mu \circ f^{-1}$, for $\mu \in \Pi(X)$.

This functor has a monad structure, with unit $\delta: Id \Rightarrow \Pi$ and multiplication $m: \Pi^2 \Rightarrow \Pi$, given as follows, for $x \in X$, $\Phi \in \Pi^2(X)$, and Borel subset $E \subseteq X$

$$\delta_X(x) = \delta_x, \quad m(\Phi)(E) = \int ev_E d\Phi,$$

where δ_x is the Dirac distribution at $x \in X$, and $ev_E: \Pi(X) \rightarrow [0, 1]$ is the evaluation function, taking $\mu \in \Pi(X)$ to $\mu(E) \in [0, 1]$. This monad is also known as the *finite distribution monad*.

For any $X \in \mathbf{Met}$, one can define a quantitative $\Sigma_{\mathcal{B}}$ -algebra $(\Pi(X), \phi_X)$ as follows, for arbitrary $\mu, \nu \in \Pi X$

$$\phi_X: \Sigma_{\mathcal{B}}\Pi X \rightarrow \Pi X \quad \phi_X(in_{+e}(\mu, \nu)) = e\mu + (1-e)\nu.$$

This quantitative algebra satisfies the interpolative barycentric theory \mathcal{B} [14, Theorem 10.4]) and is universal in the following sense.

Theorem 5.1 ([14, Th. 10.5]). *For any $\Sigma_{\mathcal{B}}$ -algebra (A, α) satisfying \mathcal{B} and non-expansive map $\beta: X \rightarrow A$, there exists a unique homomorphism $h: \Pi X \rightarrow A$ of quantitative $\Sigma_{\mathcal{B}}$ -algebras making the diagram below commute*

$$\begin{array}{ccccc} X & \xrightarrow{\delta_X} & \Pi X & \xleftarrow{\phi_X} & \Sigma_{\mathcal{B}}\Pi X \\ & \searrow \beta & \downarrow h & & \downarrow \Sigma_{\mathcal{B}}h \\ & & A & \xleftarrow{\alpha} & \Sigma_{\mathcal{B}}A \end{array}$$

From this we obtain that Π is isomorphic to the monad $T_{\mathcal{B}}$ on the quantitative theory \mathcal{B} of interpolative barycentric algebras.

Theorem 5.2. *The monads $T_{\mathcal{B}}$ and Π in \mathbf{Met} are isomorphic.*

The proof is given in the appendix.

5.2 On Complete Separable Metric Spaces

Define the functor $\Delta: \mathbf{CSMet} \rightarrow \mathbf{CSMet}$ assigning to each $X \in \mathbf{CSMet}$ the complete separable metric space $\Delta(X)$ of Borel probability measures with Kantorovich metric and acting on morphisms $f \in \mathbf{CSMet}(X, Y)$ as $\Delta(f)(\mu) = \mu \circ f^{-1}$, for $\mu \in \Delta(X)$. This functor has a monad structure, defined similarly to the one for Π . It is known as the *metric Giry monad*.

Note that Cauchy completion preserves separability. Thus the Cauchy completion functor $\mathbb{C}: \mathbf{Met} \rightarrow \mathbf{CMet}$ restricts to separable spaces. By Lemma 2.2 and the fact that $\mathbb{C}\mathbb{C} \cong \mathbb{C}$, one can verify that the canonical monad structure on $\mathbb{C}\Pi$ is isomorphic to the one on Δ in \mathbf{CSMet} . In [14], it has been proven that $T_{\mathcal{B}}$ preserves separability. Hence, by Theorem 5.2, we obtain the following.

Theorem 5.3. *The monads $\mathbb{C}T_{\mathcal{B}}$ and Δ in \mathbf{CSMet} are isomorphic.*

Note that, since \mathcal{B} is axiomatised by a continuous schema of quantitative inferences, the free monad on \mathcal{B} in \mathbf{CSMet} is exactly given by $\mathbb{C}T_{\mathcal{B}}$. Therefore, Theorem 5.3 provides an algebraic characterisation of the metric Giry monad.

6 Algebras of Contractive Operators

A *signature of contractive operators* Σ is a signature of function symbols $f: n \in \Sigma$ with associated *contractive factor* $0 < c < 1$. We write this as $f: \langle n, c \rangle \in \Sigma$.

The *quantitative equational theory of contractive operators* associated to a signature Σ , written $\mathcal{O}(\Sigma)$, is the quantitative equational theory satisfying, for each $f: \langle n, c \rangle \in \Sigma$, the axioms

$$(f\text{-Lip}) \{x_1 =_{\varepsilon} y_1, \dots, x_n =_{\varepsilon} y_n\} \vdash f(x_1, \dots, x_n) \equiv_{\delta} f(y_1, \dots, y_n),$$

for all rationals $\delta \geq c\varepsilon$. The axiom $(f\text{-Lip})$ requires the interpretation of f to be c -Lipschitz continuous.

6.1 On Metric Spaces

For a contractive signature Σ , we define a modification of the signature endofunctor on \mathbf{Met} by:

$$\tilde{\Sigma} = \coprod_{f: \langle n, c \rangle \in \Sigma} c \cdot Id^n$$

where $c \cdot Id$ is the *rescaling functor*, mapping a metric space (X, d_X) to $(X, c \cdot d_X)$.

Next we show that quantitative Σ -algebras satisfying $\mathcal{O}(\Sigma)$ are in one-to-one correspondence with universal $\tilde{\Sigma}$ -algebras; moreover, this correspondence lifts the identity functor on \mathbf{Met} (cf. [12]).

Lemma 6.1. *There exists an isomorphism of categories between $\tilde{\Sigma}\text{-Alg}$ and $\mathbb{K}(\Sigma, \mathcal{O}(\Sigma))$ making the following diagram commute*

$$\begin{array}{ccc} \tilde{\Sigma}\text{-Alg} & \xrightleftharpoons{\cong} & \mathbb{K}(\Sigma, \mathcal{O}(\Sigma)) \\ & \searrow U_{\tilde{\Sigma}} & \swarrow U_{\Sigma} \\ & \mathbf{Met} & \end{array}$$

The proof is given in the appendix.

In virtue of Lemma 6.1, by an abuse of notation, we will denote by the same name the algebras in $\tilde{\Sigma}\text{-Alg}$ and $\mathbb{K}(\Sigma, \mathcal{O}(\Sigma))$.

Next we show that the free monad $T_{\mathcal{O}(\Sigma)}$ is isomorphic to the free monad on $\tilde{\Sigma}$. For this result, we first need some discussion of free and initial algebras and free monads.

Given any endofunctor H on a category \mathbf{C} , we write $(\mu y.Hy, \alpha_H)$ for the initial H -algebra, if it exists. If \mathbf{C} has binary coproducts, the free H -algebra on $X \in \mathbf{C}$ with its unit can be identified with $(\mu y.(Hy + X), \alpha_{H+X})$, and the one exists if and only the other does. These free algebras exist if, for example, \mathbf{C} is locally countably presentable and H has countable rank.

A *free monad* on H is a monad H^* on \mathbf{C} and a natural transformation $\gamma: H \Rightarrow H^*$ that is initial among all such pairs $(S, \lambda: H \Rightarrow S)$. If the forgetful functor from $H\text{-Alg}$ to \mathbf{C} has a left adjoint (equivalently, any \mathbf{C} -object has a free H -algebra with unit), then the resulting monad on \mathbf{C} is free on H and is said to be algebraic [3]. If H^* exists and is algebraic, the category $H^*\text{-Alg}$ of Eilenberg-Moore algebras for the monad H^* is isomorphic to the category $H\text{-Alg}$ of universal algebras for the endofunctor H .

We see from the above that, if \mathbf{C} has binary sums, then H^* can be identified with $\mu y.(Hy + -)$ and the former exists if and only if the other does. We further see that if \mathbf{C} is locally countably

presentable and H has countable rank, then H^* exists and has countable rank [13].

Therefore, since \mathbf{Met} is locally countably presentable [2] and $\tilde{\Sigma}$ has countable rank, the free algebra for $\tilde{\Sigma}$ exists and so does the free monad $\tilde{\Sigma}^*$. Let $\kappa_X : \tilde{\Sigma}\tilde{\Sigma}^*X \rightarrow \tilde{\Sigma}^*X$ be the free $\tilde{\Sigma}$ -algebra on $X \in \mathbf{Met}$ with unit $\chi : Id \Rightarrow \tilde{\Sigma}^*$.

Then, the next result follows by Lemma 6.1 and freeness of $\tilde{\Sigma}^*$.

Corollary 6.2. *Let Σ be a signature of contractive operators. Then, for any quantitative Σ -algebra (A, α) satisfying $O(\Sigma)$ and non-expansive map $\beta : X \rightarrow A$, there exists a unique homomorphism $h : \tilde{\Sigma}^*X \rightarrow A$ of quantitative Σ -algebras making the diagram below commute*

$$\begin{array}{ccccc} X & \xrightarrow{\chi_X} & \tilde{\Sigma}^*X & \xleftarrow{\kappa_X} & \Sigma\tilde{\Sigma}^*X \\ & \searrow \beta & \downarrow h & & \downarrow \Sigma h \\ & & A & \xleftarrow{\alpha} & \Sigma A \end{array}$$

By freeness of $T_{O(\Sigma)}$ and Corollary 6.2, the following holds:

Theorem 6.3. *The monads $T_{O(\Sigma)}$ and $\tilde{\Sigma}^*$ in \mathbf{Met} are isomorphic.*

6.2 On Complete Metric Spaces

The category \mathbf{CMet} has coproducts and finite products. Moreover, since rescaling a metric by a factor $0 < c < 1$ preserves Cauchy completeness, the rescaling functor $c \cdot Id$ can be restricted to an endofunctor on \mathbf{CMet} . Hence, for any contractive signature Σ , the endofunctor $\tilde{\Sigma} = \coprod_{f: \langle n, c \rangle \in \Sigma} c \cdot Id^n$ is well defined in \mathbf{CMet} .

By repeating the construction in Lemma 6.1 we get the following.

Lemma 6.4. *There exists an isomorphism of categories between $\tilde{\Sigma}\text{-Alg}$ and $\mathbb{K}(\Sigma, O(\Sigma))$ making the following diagram commute*

$$\begin{array}{ccc} \tilde{\Sigma}\text{-Alg} & \xrightarrow{\cong} & \mathbb{K}(\Sigma, O(\Sigma)) \\ U_{\tilde{\Sigma}} \swarrow & & \swarrow U_{\Sigma} \\ & \mathbf{CMet} & \end{array}$$

Because $O(\Sigma)$ is a continuous quantitative theory, by Theorem 3.4, the free monad on $O(\Sigma)$ in \mathbf{CMet} is given by $\mathbb{C}T_{O(\Sigma)}$.

Note that, also \mathbf{CMet} is locally countably presentable [2], and since $\tilde{\Sigma}$ is of countable rank, we have that the free monad on $\tilde{\Sigma}$ exists in \mathbf{CMet} too. Therefore, by repeating the same argument we used before, by Lemma 6.4 and freeness of $\tilde{\Sigma}$ and $\mathbb{C}T_{O(\Sigma)}$ we obtain:

Theorem 6.5. *The monads $\mathbb{C}T_{O(\Sigma)}$ and $\tilde{\Sigma}^*$ in \mathbf{CMet} are isomorphic.*

6.3 On Complete Separable Metric Spaces

The category \mathbf{CSMet} has countable coproducts and finite products. Moreover, since the operation of rescaling a metric by a constant factor $0 < c < 1$, preserves both Cauchy completeness and separability, the endofunctor $c \cdot Id$ can be restricted to \mathbf{CSMet} .

Hence, provided that the contractive signature Σ consists of only a countable set of operators, $\tilde{\Sigma} = \coprod_{f: \langle n, c \rangle \in \Sigma} c \cdot Id^n$ is a well defined endofunctor on \mathbf{CSMet} . Hereafter, we assume the signature Σ to be countable.

Unlike \mathbf{CMet} , the category \mathbf{CSMet} is not locally countably presentable, because is not cocomplete (it does not have uncountable coproducts). However, \mathbf{CSMet} has all \aleph_1 -filtered colimits, and since every separable space in \mathbf{CMet} is a countably presentable (or \aleph_1 -presentable) object [2, Corollary 2.9], \mathbf{CSMet} is \aleph_1 -accessible. Since $\tilde{\Sigma}$ is of countable rank (i.e., \aleph_1 -accessible), by [9, Lemma 3.4] the free monad $\tilde{\Sigma}^*$ on $\tilde{\Sigma}$ exists in \mathbf{CSMet} and is algebraic.

The functor Σ^* in \mathbf{Met} preserves separability of the metric spaces. Thus, by Theorem 6.3, so does $T_{O(\Sigma)}$. Moreover, since the quantitative equational theory $O(\Sigma)$ is continuous, by Theorem 3.4, the free monad on $O(\Sigma)$ in \mathbf{CSMet} is given by $\mathbb{C}T_{O(\Sigma)}$. Thus, by freeness of $\tilde{\Sigma}^*$ and Lemma 6.1, the following holds:

Theorem 6.6. *The monads $\mathbb{C}T_{O(\Sigma)}$ and $\tilde{\Sigma}^*$ in \mathbf{CSMet} are isomorphic.*

7 Interpolative Barycentric Algebras with Contractive Operators

In this section we study a variation of interpolative barycentric quantitative algebras where we add operations from a contractive signature Σ assumed to be disjoint from $\Sigma_{\mathcal{B}}$.

These are quantitative algebras for the signature

$$\Sigma_{\mathcal{B}} + \Sigma = \{+e : 2 \mid e \in [0, 1]\} \cup \Sigma,$$

satisfying the disjoint union of the axioms of interpolative barycentric quantitative theory, namely (B1), (B2), (SC), (SA), and (IB), and, for each $f \in \Sigma$, the axiom (f -Lip) from the quantitative theory of contractive operators.

The quantitative equational theory induced by these axioms will be called *interpolative barycentric theory with contractive operators* in Σ , and it coincides with the disjoint union $\mathcal{B} + O(\Sigma)$ of theories.

7.1 On Metric Spaces

We have already noted that the quantitative theories \mathcal{B} and $O(\Sigma)$ are simple. Thus, by Theorem 4.4, the free monad $T_{\mathcal{B}+O(\Sigma)}$ on \mathbf{Met} induced by $\mathcal{B} + O(\Sigma)$ is the sum of $T_{\mathcal{B}}$ and $T_{O(\Sigma)}$. Moreover, by Theorems 5.2 and 6.3, it is also isomorphic to $\Pi + \tilde{\Sigma}^*$.

In the following we will prove an alternative characterisation of $T_{\mathcal{B}+O(\Sigma)}$, that will eventually reveal the connection between interpolative barycentric algebras with operators and Markov processes.

Depending on the type of monads, several specific conditions of existence and constructions appear in the literature [1] for the sum of monads. One of these, due to Hyland, Plotkin, and Power [10], recalled below for convenience, characterises the sum of a monad with a free one.

Theorem 7.1 ([10, Theorem 4]). *Given an endofunctor F and a monad T on a category \mathcal{C} , if the free monads F^* and $(FT)^*$ exist and are algebraic, then the sum of monads $T + F^*$ exists in the category of monads over \mathcal{C} and is given by a canonical monad structure on the composite $T(FT)^*$.*

In terms of the above result, given that $T_{\mathcal{B}+O(\Sigma)}$ is isomorphic to $\Pi + \tilde{\Sigma}^*$, if we prove that the free monad $(\tilde{\Sigma}\Pi)^*$ in \mathbf{Met} exists and is algebraic, then $T_{\mathcal{B}+O(\Sigma)}$ would be also isomorphic to $\Pi(\tilde{\Sigma}\Pi)^*$. For this reason, in the following we characterise (and hence prove the existence of) the free algebra on $\tilde{\Sigma}\Pi$. relevant For arbitrary $X \in \mathbf{Met}$, let P_X be the smallest set such that

- if $x \in X$, then $\underline{x} \in P_X$;
- if $\mu_1, \dots, \mu_n \in \Pi(P_X)$ and $f : n \in \Sigma$, then $f\langle \mu_1, \dots, \mu_n \rangle \in P_X$.

We define the metric $d_{P_X} : P_X \times P_X \rightarrow [0, 1]$ by induction on the complexity on the structure of the elements in P_X as follows², for arbitrary $x, x' \in X$, $\mu_1, \dots, \mu_n, \nu_1, \dots, \nu_k \in \Pi(P_X)$, and distinct operators $f, g \in \Sigma$ of arity $f : n$ and $g : k$ such that $n \leq k$, where

²The symmetric cases are omitted and defined as expected.

we assume $f : \langle n, c \rangle \in \Sigma$.

$$d_{P_X}(x, x') = d_X(x, x'),$$

$$d_{P_X}(f\langle \mu_1, \dots, \mu_n \rangle, f\langle \nu_1, \dots, \nu_n \rangle) = c \cdot \max_{i=1}^n \mathcal{K}(d_{P_X})(\mu_i, \nu_i),$$

$$d_{P_X}(f\langle \mu_1, \dots, \mu_n \rangle, g\langle \nu_1, \dots, \nu_k \rangle) = 1,$$

$$d_{P_X}(x, f\langle \mu_1, \dots, \mu_n \rangle) = 1.$$

Proposition 7.2. *For any $X \in \mathbf{Met}$, d_{P_X} is a well defined metric.*

For $X \in \mathbf{Met}$, define $\varrho_X : X \rightarrow P_X$ and $\vartheta_X : \tilde{\Sigma}\Pi P_X \rightarrow P_X$ as follows, for arbitrary $x \in X$, $f : n \in \Sigma$, and $\mu_1, \dots, \mu_n \in \Pi(P_X)$

$$\varrho_X(x) = x, \quad \vartheta_X(\inf(\mu_1, \dots, \mu_n)) = f\langle \mu_1, \dots, \mu_n \rangle.$$

By definition of d_{P_X} , it is straightforward to show that both ϱ_X and ϑ_X are non-expansive maps (actually, are isometric injections), thus they are morphisms in \mathbf{Met} . In particular, ϑ_X is a universal $\tilde{\Sigma}\Pi$ -algebra on P_X in \mathbf{Met} .

Theorem 7.3 (Free Algebra). *For any $X \in \mathbf{Met}$, $\tilde{\Sigma}\Pi$ -algebra (A, α) , and non-expansive map $\beta : X \rightarrow A$, there exists a unique $\tilde{\Sigma}\Pi$ -homomorphism $h : P_X \rightarrow A$ making the diagram below commute*

$$\begin{array}{ccc} X & \xrightarrow{\varrho_X} & P_X & \xleftarrow{\vartheta_X} & \tilde{\Sigma}\Pi P_X \\ & \searrow \beta & \downarrow h & & \downarrow \tilde{\Sigma}\Pi h \\ & & A & \xleftarrow{\alpha} & \tilde{\Sigma}\Pi A \end{array}$$

The proof is given in the appendix.

Theorem 7.3 states that ϑ_X is the free $\tilde{\Sigma}\Pi$ -algebra for $X \in \mathbf{Met}$, or equivalently, that the forgetful functor from the category of $\tilde{\Sigma}\Pi$ -algebras to \mathbf{Met} has a left adjoint. Thus, the free monad $(\tilde{\Sigma}\Pi)^*$ on $\tilde{\Sigma}\Pi$ in \mathbf{Met} exists and is algebraic. Moreover, it acts on objects $X \in \mathbf{Met}$ as $(\tilde{\Sigma}\Pi)^*X = P_X$.

Corollary 7.4. *The monads $T_{\mathcal{B}+O(\Sigma)}$ and $\Pi(\tilde{\Sigma}\Pi)^*$ in \mathbf{Met} are isomorphic.*

Proof. Direct consequence of Theorems 4.4, 5.2, 6.3, 7.1, and 7.3. \square

As observed in [10], the monad $T(FT)^*$ of Theorem 7.1 is simply another form of the generalised resumptions monad transformer of Cenciarelli and Moggi [5], sending T to $\mu y.T(Fy + -)$. Hence, by the characterisation above and guided by the same observations that lead to [10, Corollary 2], we obtain the following isomorphism.

Theorem 7.5. *The monad $T_{\mathcal{B}+O(\Sigma)}$ in \mathbf{Met} is isomorphic to the canonical monad structure on $\mu y.\Pi(\tilde{\Sigma}y + -)$.*

Proof. By [23, Proposition 5.3], it is easy to show that $\mu y.\Pi(\tilde{\Sigma}y + -)$ exists if and only if $(\tilde{\Sigma}\Pi)^*$ does, and that $\Pi(\tilde{\Sigma}\Pi)^*$ and $\mu y.\Pi(\tilde{\Sigma}y + -)$ are then isomorphic. \square

7.2 On Complete Separable Metric Spaces

We would like to apply Corollary 4.6 to provide a characterisation of the free monad on $\mathcal{B} + O(\Sigma)$ in the category \mathbf{CSMet} as the sum of monads $\mathcal{CT}_{\mathcal{B}} + \mathcal{CT}_{O(\Sigma)}$.

To this end we have to verify that the conditions required by the corollary are satisfied. We already noted that the quantitative theories \mathcal{B} and $O(\Sigma)$ are continuous and that the functors $T_{\mathcal{B}}$ and $T_{O(\Sigma)}$ preserve separability of the metric spaces. We are only left to prove that also $T_{\mathcal{B}+O(\Sigma)}$ preserves separability.

Lemma 7.6. *The functor $T_{\mathcal{B}+O(\Sigma)}$ in \mathbf{Met} preserves separability.*

The proof is given in the appendix.

Thus, we can apply Corollary 4.6 to obtain the desired result.

Corollary 7.7. *The monad $\mathcal{CT}_{\mathcal{B}+O(\Sigma)}$ in \mathbf{CSMet} is the sum of $\mathcal{CT}_{\mathcal{B}}$ and $\mathcal{CT}_{O(\Sigma)}$.*

An immediate consequence of the characterisation above and Theorems 5.3 and 6.6 is the following.

Corollary 7.8. *The monad $\mathcal{CT}_{\mathcal{B}+O(\Sigma)}$ in \mathbf{CSMet} is isomorphic to $\Delta + \tilde{\Sigma}^*$.*

According to the above and similarly to what we have done in Section 7.1, we would like to prove for the free monad $\mathcal{CT}_{\mathcal{B}+O(\Sigma)}$ on $\mathcal{B} + O(\Sigma)$ in \mathbf{CSMet} corresponding results to Corollary 7.4 and Theorem 7.5.

We will proceed again by using Theorem 7.1. Thus, as we did in Section 7.1, we need to show that the free monad on $\tilde{\Sigma}\Delta$ in \mathbf{CSMet} exists and is algebraic. We already noted that the category \mathbf{CSMet} is accessible and that $\tilde{\Sigma}$ is an accessible endofunctor on it. Moreover, by [26, Corollary 22], we also have that Δ on \mathbf{CSMet} is accessible, so that their composition $\tilde{\Sigma}\Delta$ is accessible too. Therefore, by [9, Lemma 3.4] the free monad $(\tilde{\Sigma}\Delta)^*$ exists and is algebraic.

This observation, then leads to the desired characterisations.

Corollary 7.9. *The monad $\mathcal{CT}_{\mathcal{B}+O(\Sigma)}$ in \mathbf{CSMet} is isomorphic to the monads $\Pi(\tilde{\Sigma}\Pi)^*$ and $\mu y.\Delta(\tilde{\Sigma}y + -)$ with the canonical monad structures.*

8 The Algebras of Markov Processes

In this section we show how interpolative barycentric quantitative theories with operators can be used to axiomatise the probabilistic bisimilarity distance of Desharnais et al. [7] over Markov processes.

For any $0 < c < 1$, we define the finite signature of contractive operators

$$\mathcal{M}_c = \{0 : \langle 0, c \rangle\} \cup \{\diamond : \langle 1, c \rangle\},$$

consisting of one constant symbol 0 , representing termination, and a unary operator $\diamond(t)$ expressing the capability of doing a transition to t . Both operators are associated with the same contractive factor.

The interpolative barycentric quantitative theory with operators in \mathcal{M}_c , given as the disjoint union $\mathcal{B} + O(\mathcal{M}_c)$ of theories is generated by the following set of axioms

$$(B1) \vdash x +_1 y \equiv_0 x,$$

$$(B2) \vdash x +_e x \equiv_0 x,$$

$$(SC) \vdash x +_e y \equiv_0 y +_{1-e} x,$$

$$(SA) \vdash (x +_e y) +_{e'} z \equiv_0 x +_{ee'} (y +_{\frac{e'-ee'}{1-ee'}} z), \text{ for } e, e' \in [0, 1),$$

$$(IB) \{x \equiv_\varepsilon y, x' \equiv_{\varepsilon'} y'\} \vdash x +_e x' \equiv_\delta y +_e y', \text{ for } \delta \geq e\varepsilon + (1-e)\varepsilon',$$

$$(\diamond\text{-Lip}) \{x =_\varepsilon y\} \vdash \diamond(x) \equiv_\delta \diamond(y), \text{ for } \delta \geq c\varepsilon.$$

Note that, the constant 0 has no explicit associated axiom, since it is derivable from (Refl).

8.1 Markov Processes over Metric Spaces

We briefly recall the definitions of Markov processes over metric spaces and discounted probabilistic bisimilarity distance on them, presented following the pattern proposed in [26, Section 6].

Definition 8.1. A (sub-probabilistic) *Markov process* over a metric space is a tuple (X, τ) consisting of a metric space X of *states* and non-expansive *Markov kernel* $\tau : X \rightarrow \Delta(1 + X)$.

It is clear that these structures correspond to the coalgebras for the (sub-probabilistic) Giry functor $\Delta(1 + Id)$ in **Met**.

In [26], van Breugel et al. proved that the final coalgebra for $\Delta(1 + Id)$ in **Met** exists and they characterised the probabilistic bisimilarity distance on Markov processes as the pseudometric induced by the unique homomorphism to the final coalgebra.

We will do the same here slightly extended their approach for dealing with the case when the probabilistic bisimilarity distance is discounted by a factor $0 < c < 1$. Explicitly, the only difference consists in considering coalgebras for the functor $\Delta(1 + c \cdot Id)$ in **Met**. For simplicity we call these structures c -Markov processes. Note that Markov processes is a proper subclass: one can turn any Markov process into a c -Markov process as $(X, \Delta(1 + id_X^c) \circ \tau)$, where $id^c : Id \Rightarrow c \cdot Id$ is the obvious natural transformation acting as the identity on the elements of the metric space and allowing for the change of “type”.

The final coalgebra (Z_c, ω_c) for $\Delta(1 + c \cdot Id)$ exists by similar arguments of [26, Section 6]. Then, for an arbitrary c -Markov process (X, τ) , the c -discounted probabilistic bisimilarity pseudometric on (X, τ) is defined as the function $d_\tau^c : X \times X \rightarrow [0, 1]$ given as

$$d_\tau^c(x, x') = d_{Z_c}(h(x), h(x')),$$

where $h : X \rightarrow Z_c$ is the unique homomorphism of coalgebras from (X, τ) to (Z_c, ω_c) .

Since terminal objects are unique up to isomorphism, the definition of the distance function d_τ^c does not depend on which terminal $\Delta(1 + c \cdot Id)$ -coalgebra is chosen. Clearly, since d_{Z_c} is a 1-bounded metric, then d_τ^c is a well defined 1-bounded pseudometric.

Proposition 8.2 ([26]). *Let (X, τ) be a c -Markov process. Then, for all $x, x' \in X$, $d_\tau^c(x, x') = 0$, if and only if, x and x' are probabilistically bisimilar.*

We can show that this distance has a characterisation as the least fixed point of a monotone function on a complete lattice of 1-bounded pseudometrics.

Theorem 8.3 ([26]). *The c -discounted probabilistic bisimilarity pseudometric d_τ^c on (X, τ) is the least fixed point of the following operator on the complete lattice of 1-bounded pseudometrics d on X with point-wise order \sqsubseteq , such that $d \sqsubseteq d_X$,*

$$\Psi_\tau^c(d)(x, x') = \sup_{f \in \Phi_{1+c \cdot X}} \left| \int f \, d\tau(x) - \int f \, d\tau(x') \right|,$$

with supremum ranging over the set $\Phi_{1+c \cdot X}$ of non-expansive positive 1-bounded real valued functions $f : 1 + c \cdot X \rightarrow [0, 1]$.

8.2 On Metric Spaces

In this section we want to relate c -Markov processes and their c -discounted probabilistic bisimilarity pseudometric with the free monad arising from the quantitative theory $\mathcal{B} + \mathcal{O}(\mathcal{M}_c)$ in **Met**.

First note that the functor associated to the signature \mathcal{M}_c is

$$\tilde{\mathcal{M}}_c = 1 + c \cdot Id.$$

where 1 is the terminal object in **Met** (i.e., the singleton metric space)³. Thus, by Theorem 7.5, the free monad on $\mathcal{B} + \mathcal{O}(\mathcal{M}_c)$ corresponds to the canonical monad structure on $\mu y. \Pi(1 + c \cdot y + -)$.

³Here we are implicitly applying the isomorphism $1 \cong c \cdot 1$.

Explicitly, this means that, the free monad on $\mathcal{B} + \mathcal{O}(\mathcal{M}_c)$ assigns to an arbitrary metric space $M \in \mathbf{Met}$ the *initial solution* of the following functorial equation in **Met**

$$FP_M \cong \Pi(1 + c \cdot FP_M + M).$$

In particular, when $M = 0$ is the empty metric space (i.e., the initial object) the above corresponds to the isomorphism on the initial $\Pi(1 + c \cdot Id)$ -algebra. This gives rise to a $\Pi(1 + c \cdot Id)$ -coalgebra structure on FP_0 , which in turn can be converted into a c -Markov process via a post-composition with the subspace inclusion $\Pi(-) \hookrightarrow \Delta(-)$. Let us write this c -Markov process as

$$(FP, \alpha : FP \rightarrow \Delta(1 + c \cdot FP)).$$

The following states that the metric on FP corresponds to the c -discounted probabilistic bisimilarity (pseudo)metric on (FP, α) .

Lemma 8.4. $d_{FP} = d_\alpha^c$

Proof. By Theorem 8.3 we need to prove that d_{FP} is the least fixed point of Ψ_α^c . This follows trivially by definition of the functor $\Pi(1 + c \cdot Id)$ and because (FP, α^{-1}) is the initial $\Pi(1 + c \cdot Id)$ -algebra. \square

Next we would like to give a more explicit characterisation of the elements in FP . By recalling the characterisation of the metric term monad in [14], the elements in FP can be represented by equivalence classes of terms generated by the following grammar

$$f ::= \mathbf{0} \mid \diamond(f) \mid f +_e f. \quad \text{for } e \in [0, 1]$$

with respect to the kernel of the distance. In this specific case the distance corresponds to d_α^c , with transition probability function α defined as follows:

$$\alpha(\mathbf{0}) = \delta_\perp, \quad \alpha(\diamond(f)) = \delta_f, \quad \alpha(f +_e g) = \alpha(f) +_e \alpha(g).$$

Thus, by Lemma 8.2, we can interpret the element in FP as pointed (or rooted) Markov processes constructed over the above grammar and quotiented by bisimilarity. It is not difficult to see that these structures correspond to the class of *rooted acyclic finite Markov processes* from [6].

8.3 On Complete Separable Metric Spaces

We would like to relate c -Markov processes and the c -discounted probabilistic bisimilarity pseudometric with the free monad in **CSMet**.

By Corollary 7.9, we know that the free monad on $\mathcal{B} + \mathcal{O}(\mathcal{M}_c)$ in **CSMet** corresponds to the canonical monad structure on $\mu y. \Delta(1 + c \cdot y + -)$.

Explicitly, this means that, for the case of complete metric spaces the free monad on $\mathcal{B} + \mathcal{O}(\mathcal{M}_c)$ assigns to any arbitrary metric space $M \in \mathbf{CSMet}$ the *initial solution* of the following functorial equation in **CSMet**

$$MP_M \cong \Delta(1 + c \cdot MP_M + M).$$

Observe that the map $\omega_M : MP_M \rightarrow \Delta(1 + c \cdot MP_M + M)$ arising from the above isomorphism is a coalgebra structure for the functor $\Delta(1 + c \cdot Id + M)$. Next we show that (MP_M, ω_M) is actually the final coalgebra in **CSMet**.

We will do this by using the following result from [25, Section 7].

Theorem 8.5 ([25]). *Every locally contractive endofunctor H on **CSMet** has a unique fixed point which is both an initial algebra and a final coalgebra for H .*

Note that if the fixed point lies in a subcategory of \mathbf{CMet} , it is unique also in that subcategory. Hence, our goal is to prove that, for any $M \in \mathbf{CMet}$, the functor $\Delta(1 + c \cdot Id + M)$ is locally contractive.

In \mathbf{CMet} the homsets $\mathbf{CMet}(X, Y)$ are themselves complete separable metric spaces, with distance

$$d_{X \rightarrow Y}(f, g) = \sup_{x \in X} d_Y(f(x), g(x)).$$

An endofunctor H on \mathbf{CMet} is *locally c -Lipschitz continuous* if for all $X, Y \in \mathbf{CMet}$ and non-expansive maps $f, g: X \rightarrow Y$,

$$d_{HX \rightarrow HY}(H(f), H(g)) \leq c \cdot d_{X \rightarrow Y}(f, g).$$

We say that H is *locally non-expansive* if it is locally 1-Lipschitz continuous, and *locally contractive* if it is locally c -Lipschitz continuous, for some $0 \leq c < 1$.

Examples of locally contractive functors are the constant functors and the rescaling functor $c \cdot Id$, for $0 \leq c < 1$. Locally contractiveness is preserved by products and coproducts and composition. Moreover, if H is locally non-expansive and G is locally contractive, then FG is locally contractive.

Lemma 8.6. *The endofunctor Δ on \mathbf{CMet} is locally non-expansive.*

Thus, for the free monad on $\mathcal{B} + \mathcal{O}(\mathcal{M}_c)$ in \mathbf{CSMet} , the following holds.

Theorem 8.7. *For every $M \in \mathbf{CSMet}$, (MP_M, ω_M) is the final coalgebra of the functor $\Delta(1 + c \cdot Id + M)$ in \mathbf{CSMet} .*

Proof. This is a direct consequence of Theorem 8.5 and Lemma 8.6, since, $1 + c \cdot Id + M$ is locally contractive and the composition of a locally contractive functor with a locally non-expansive one is locally contractive. \square

Note that, when $M = 0$ is the empty metric space, the coalgebras of this functor correspond to the final c -Markov process we have used in Section 8.1 to characterise the c -discounted probabilistic bisimilarity distance. When M is not the empty space we have a kind of Markov process that terminates in the states in M ; one can view the states of M as absorbing states.

Hence, in the light of the Theorem 8.7, we have shown that $\mathcal{B} + \mathcal{O}(\mathcal{M}_c)$, for the case of complete metric spaces, axiomatises the c -probabilistic bisimilarity distance on the final Markov process.

9 Conclusions

The main contribution of this paper was extending the notion of “sum of theories” from [10] to the quantitative setting. This, we feel opens the way to developing combinations of quantitative effects just as [10] did for combining effects in the ordinary sense. The Markov process example developed in this paper is of interest in its own right as it is the underlying operational semantics for probabilistic programming languages.

A significant novelty of this paper is a treatment of Markov processes that presents them both as algebras and as coalgebras. The algebra structure arises by combining the quantitative equational theory of probability distributions equipped with the Kantorovich metric whereas the coalgebra structure corresponds to the final coalgebra equipped with the discounted probabilistic bisimilarity distance. Such algebra-coalgebra duality has been used before [4] for automata but much more could be done and in future work we hope to use this connection to reason about properties of probabilistic programs.

Acknowledgments

This research was supported by grants from the Danish Research Council and from NSERC (Canada).

References

- [1] Jiri Adámek, Stefan Milius, Nathan Bowler, and Paul Blain Levy. 2012. Coproducts of Monads on Set. In *LICS 2012*. IEEE Computer Society, 45–54. <https://doi.org/10.1109/LICS.2012.16>
- [2] Jiri Adámek, Stefan Milius, and Lawrence S. Moss. 2012. On Finitary Functors and Their Presentations. In *CMCS 2012 (Lecture Notes in Computer Science)*, Vol. 7399. Springer, 51–70. https://doi.org/10.1007/978-3-642-32784-1_4
- [3] Michael Barr. 1970. Coequalizers and free triples. *Mathematische Zeitschrift* 116, 4 (1970), 307–322. <https://doi.org/10.1007/BF0111838>
- [4] Filippo Bonchi, Marcello M. Bonsangue, Helle Hvid Hansen, Prakash Panangaden, Jan J. M. M. Rutten, and Alexandra Silva. 2014. Algebra-coalgebra duality in Brzozowski’s minimization algorithm. *ACM Trans. Comput. Log.* 15, 1 (2014), 3:1–3:29. <https://doi.org/10.1145/2490818>
- [5] Pietro Cenciarelli and Eugenio Moggi. 1993. *A Syntactic Approach to Modularity in Denotational Semantics*. Technical report. CWI. Proc. 5th. Biennial Meeting on Category Theory and Computer Science.
- [6] Josee Desharnais, Vineet Gupta, Radha Jagadeesan, and Prakash Panangaden. 2003. Approximating labelled Markov processes. *Inf. Comput.* 184, 1 (2003), 160–200. [https://doi.org/10.1016/S0890-5401\(03\)00051-8](https://doi.org/10.1016/S0890-5401(03)00051-8)
- [7] Josee Desharnais, Vineet Gupta, Radha Jagadeesan, and Prakash Panangaden. 2004. Metrics for labelled Markov processes. *Theoretical Computer Science* 318, 3 (2004), 323–354.
- [8] J. Dugundji. 1966. *Topology*. J. Wiley.
- [9] Neil Ghani, Christoph Lüth, Federico De Marchi, and John Power. 2001. Algebras, Coalgebras, Monads and Comonads. *Electr. Notes Theor. Comput. Sci.* 44, 1 (2001), 128–145. [https://doi.org/10.1016/S1571-0661\(04\)80905-8](https://doi.org/10.1016/S1571-0661(04)80905-8)
- [10] Martin Hyland, Gordon D. Plotkin, and John Power. 2006. Combining effects: Sum and tensor. *Theor. Comput. Sci.* 357, 1–3 (2006), 70–99. <https://doi.org/10.1016/j.tcs.2006.03.013>
- [11] Martin Hyland and John Power. 2007. The Category Theoretic Understanding of Universal Algebra: Lawvere Theories and Monads. *Electronic Notes in Theor. Comp. Sci.* 172 (2007), 437–458.
- [12] Peter T. Johnstone. 1975. Adjoint Lifting Theorems for Categories of Algebras. *Bulletin of the London Mathematical Society* 7, 3 (1975), 294–297. <https://doi.org/10.1112/blms/7.3.294>
- [13] G.M. Kelly. 1980. A unified treatment of transfinite constructions for free algebras, free monoids, colimits, associated sheaves, and so on. *Bulletin of the Australian Mathematical Society* 22, 1 (1980), 1–83. <https://doi.org/10.1017/S0004972700006353>
- [14] Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. 2016. Quantitative Algebraic Reasoning. In *LICS*. ACM, 700–709. <https://doi.org/10.1145/2933575.2934518>
- [15] Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. 2017. On the axiomatizability of quantitative algebras. In *LICS 2017*. IEEE Computer Society, 1–12. <https://doi.org/10.1109/LICS.2017.8005102>
- [16] Eugenio Moggi. 1988. *The partial lambda calculus*. Ph.D. Dissertation. University of Edinburgh. College of Science and Engineering. School of Informatics.
- [17] Eugenio Moggi. 1991. Notions of computation and monads. *Information and computation* 93, 1 (1991), 55–92.
- [18] Prakash Panangaden. 2009. *Labelled Markov Processes*. Imperial College Press.
- [19] Gordon Plotkin and John Power. 2001. Semantics for algebraic operations. *Electronic Notes in Theoretical Computer Science* 45 (2001), 332–345.
- [20] Gordon Plotkin and John Power. 2002. Notions of computation determine monads. In *Foundations of Software Science and Computation Structures*. Springer, 342–356.
- [21] Dana Scott. 1970. *Outline of a mathematical theory of computation*. Technical Monograph PRG-2. Oxford University Computing Laboratory.
- [22] Dana Scott. 1972. Lattice Theory, Data Types and Semantics. In *NYU Symposia in Computer Science*, Randall Rustin (Ed.), 65–106.
- [23] Alex K. Simpson and Gordon D. Plotkin. 2000. Complete Axioms for Categorical Fixed-Point Operators. In *LICS 2000*. IEEE Computer Society, 30–41. <https://doi.org/10.1109/LICS.2000.855753>
- [24] Marshall H. Stone. 1949. Postulates for the barycentric calculus. *Annali di Matematica Pura ed Applicata* 29, 1 (1949), 25–30.
- [25] Daniele Turi and Jan J. M. M. Rutten. 1998. On the Foundations of Final Coalgebra Semantics. *Mathematical Structures in Computer Science* 8, 5 (1998), 481–540.
- [26] Franck van Breugel, Claudio Hermida, Michael Makkai, and James Worrell. 2007. Recursively defined metric spaces without contraction. *Theor. Comput. Sci.* 380, 1–2 (2007), 143–163. <https://doi.org/10.1016/j.tcs.2007.02.059>
- [27] Cédric Villani. 2008. *Optimal transport: old and new*. Springer-Verlag.

A Proofs from Section 4

Here is the proof of Proposition 4.1.

Proposition A.1. $\mathbb{K}(\Sigma + \Sigma', \mathcal{U} + \mathcal{U}') \cong \mathbb{K}((\Sigma, \mathcal{U}) \oplus (\Sigma', \mathcal{U}'))$.

Proof. The isomorphism is given by the following pair of functors

$$\mathbb{K}(\Sigma + \Sigma', \mathcal{U} + \mathcal{U}') \begin{array}{c} \xrightarrow{H} \\ \xleftarrow{K} \end{array} \mathbb{K}((\Sigma, \mathcal{U}) \oplus (\Sigma', \mathcal{U}'))$$

defined, for an arbitrary quantitative $(\Sigma + \Sigma')$ -algebra (A, γ) satisfying $\mathcal{U} + \mathcal{U}'$ and a $\langle \mathcal{U}, \mathcal{U}' \rangle$ -bialgebra (B, α, β) , respectively as

$$H(A, \gamma) = (A, \gamma \circ in_l, \gamma \circ in_r), \quad K(B, \alpha, \beta) = (B, [\alpha, \beta]),$$

where $[\alpha, \beta]$ is the unique map induced by α and β by couniversality of the coproduct $\Sigma A + \Sigma' A$. On morphisms both functors map a morphism to itself; it is easy to see that a homomorphism in one sense is also a homomorphism in the other.

The fact that the functors are inverses is clear: $H \circ K = Id$ and $K \circ H = Id$ follow immediately from the couniversal property of coproducts. We are done, provided we show that the functors are indeed well defined. In order to show that the functors are well defined, we need to prove that the functors preserve the relevant quantitative equations.

To show that H is well defined we need to prove that whenever (A, γ) satisfies $\mathcal{U} + \mathcal{U}'$, then $(A, \gamma \circ in_l)$ and $(A, \gamma \circ in_r)$ satisfy \mathcal{U} and \mathcal{U}' , respectively. We will prove only that $(A, \gamma \circ in_l)$ satisfies \mathcal{U} , since the other follows similarly. Let $\Gamma \vdash t \equiv_\varepsilon s \in \mathcal{U}$ and $\iota: X \rightarrow A$ be an arbitrary assignment of the variables. Since $\mathcal{U} \subseteq \mathcal{U} + \mathcal{U}'$, we have:

$$\begin{array}{c} \text{(for all } t' \equiv_{\varepsilon'} s' \in \Gamma, d_A(i^\sharp(t'), i^\sharp(s')) \leq \varepsilon') \\ \text{implies} \\ d_A(i^\sharp(t), i^\sharp(s)) \leq \varepsilon, \end{array} \quad (3)$$

where $i^\sharp: \mathbb{T}(\Sigma + \Sigma', X) \rightarrow A$ is the homomorphic extension of ι on (A, γ) .

Note that, by definition of coproduct of functors and homomorphic extension, we have that the following diagram commutes

$$\begin{array}{ccccc} X & \xrightarrow{\eta_X^\Sigma} & \mathbb{T}(\Sigma, X) & \xleftarrow{\psi_X^\Sigma} & \Sigma \mathbb{T}(\Sigma, X) \\ & \searrow \eta_X^{\Sigma+\Sigma'} & \downarrow i & & \downarrow \Sigma i \\ & & \mathbb{T}(\Sigma+\Sigma', X) & \xleftarrow{\psi_{\Sigma+\Sigma'}} & (\Sigma+\Sigma')\mathbb{T}(\Sigma+\Sigma', X) & \xleftarrow{in_l} & \Sigma \mathbb{T}(\Sigma+\Sigma', X) \\ & & \downarrow i^\sharp & & \downarrow (\Sigma+\Sigma')i^\sharp & & \downarrow \Sigma i^\sharp \\ & & A & \xleftarrow{\gamma} & (\Sigma+\Sigma')A & \xleftarrow{in_l} & \Sigma A \end{array}$$

where i is the canonical inclusion of Σ -terms in $\mathbb{T}(\Sigma + \Sigma', X)$. The above implies also that $i^\sharp \circ i$ is the homomorphic extension of ι on $(A, \gamma \circ in_l)$. Recall that \mathcal{U} is of type Σ . Thus in $\Gamma \vdash t \equiv_\varepsilon s$ can occur only terms in $\mathbb{T}(\Sigma, X)$. Therefore, (3) implies that $(A, \gamma \circ in_l)$ satisfies $\Gamma \vdash t \equiv_\varepsilon s$. This argument is general so it applies to the whole theory \mathcal{U} .

For K , we need to show that whenever (A, α) satisfies \mathcal{U} and (A, β) satisfies \mathcal{U}' , then $(A, [\alpha, \beta])$ satisfies $\mathcal{U} + \mathcal{U}'$. By the definition of the disjoint union of quantitative theories, it suffices to prove that $(A, [\alpha, \beta])$ is a model for both \mathcal{U} and \mathcal{U}' . We show the former case, since the other follows similarly. Let $\Gamma \vdash t \equiv_\varepsilon s \in \mathcal{U}$

and $\iota: X \rightarrow A$ be an arbitrary assignment of the variables. Since (A, α) satisfies \mathcal{U} , we have that

$$\begin{array}{c} \text{(for all } t' \equiv_{\varepsilon'} s' \in \Gamma, d_A(i^\sharp(t'), i^\sharp(s')) \leq \varepsilon') \\ \text{implies} \\ d_A(i^\sharp(t), i^\sharp(s)) \leq \varepsilon, \end{array} \quad (4)$$

where $i^\sharp: \mathbb{T}(\Sigma, X) \rightarrow A$ is the homomorphic extension of ι on (A, α) . Note that, by definition of coproduct of functors and homomorphic extension, we have that the following diagram commutes

$$\begin{array}{ccccc} X & \xrightarrow{\eta_X^\Sigma} & \mathbb{T}(\Sigma, X) & \xleftarrow{\psi_X^\Sigma} & \Sigma \mathbb{T}(\Sigma, X) \\ & \searrow \eta_X^{\Sigma+\Sigma'} & \downarrow i & & \downarrow \Sigma i \\ & & \mathbb{T}(\Sigma+\Sigma', X) & \xleftarrow{\psi_{\Sigma+\Sigma'}} & (\Sigma+\Sigma')\mathbb{T}(\Sigma+\Sigma', X) & \xleftarrow{in_l} & \Sigma \mathbb{T}(\Sigma+\Sigma', X) \\ & & \downarrow i^\flat & & \downarrow (\Sigma+\Sigma')i^\flat & & \downarrow \Sigma i^\flat \\ & & A & \xleftarrow{[\alpha, \beta]} & (\Sigma+\Sigma')A & \xleftarrow{in_l} & \Sigma A \end{array}$$

where i is the canonical inclusion of Σ -terms in $\mathbb{T}(\Sigma + \Sigma', X)$, and i^\flat is the homomorphic extension of ι on $(A, [\alpha, \beta])$. Since $[\alpha, \beta] \circ in_l = \alpha$, the above implies also that $i^\flat \circ i = i^\sharp$. Since \mathcal{U} is of type Σ , then $\Gamma \vdash t \equiv_\varepsilon s$ contains only terms in $\mathbb{T}(\Sigma, X)$. Therefore, (4) implies that $(A, [\alpha, \beta])$ satisfies $\Gamma \vdash t \equiv_\varepsilon s$; again this implies the result for all of \mathcal{U} . \square

This is the proof of Theorem 4.5

Theorem A.2. Let $\mathcal{U}, \mathcal{U}'$ be continuous quantitative theories. Then, the monad $\mathbb{C}T_{\mathcal{U}+\mathcal{U}'}$ in \mathbf{CMet} is the sum of $\mathbb{C}T_{\mathcal{U}}$ and $\mathbb{C}T_{\mathcal{U}'}$.

Proof. By Theorem 3.4, the monads $\mathbb{C}T_{\mathcal{U}+\mathcal{U}'}$, $\mathbb{C}T_{\mathcal{U}}$, and $\mathbb{C}T_{\mathcal{U}'}$ are, respectively, the free monads on $\mathcal{U} + \mathcal{U}'$, \mathcal{U} , and \mathcal{U}' in \mathbf{CMet} .

Similarly to Corollary 4.3, one obtains that $\mathbb{C}\mathbb{K}(\Sigma + \Sigma', \mathcal{U} + \mathcal{U}')$ and $\langle \mathbb{C}T_{\mathcal{U}}, \mathbb{C}T_{\mathcal{U}'} \rangle$ -**biAlg** are isomorphic. Thus, by Theorem 3.4 the forgetful functor from $\langle \mathbb{C}T_{\mathcal{U}}, \mathbb{C}T_{\mathcal{U}'} \rangle$ -**biAlg** to \mathbf{Met} has a left adjoint, and the monad generated by this adjunction is isomorphic to $\mathbb{C}T_{\mathcal{U}+\mathcal{U}'}$. Thus, by [13] (cf. also [1, Proposition 2.8]), $\mathbb{C}T_{\mathcal{U}+\mathcal{U}'}$ is the sum of $\mathbb{C}T_{\mathcal{U}}$ and $\mathbb{C}T_{\mathcal{U}'}$. \square

Lemma A.3. Let \mathcal{U} be a quantitative theory of type Σ . If Σ is countable, then $T_{\mathcal{U}}$ preserves separability of the metric spaces.

Proof. Let \mathcal{U}_0 be the quantitative theory induced without extra axioms (i.e., satisfying only the meta-axioms of quantitative equational theories). Hence, any quantitative Σ -algebra is a model of \mathcal{U}_0 , that is, $\mathbb{K}(\Sigma, \mathcal{U}_0) = \Sigma\text{-Alg}$.

Since \mathbf{Met} is locally countably presentable [2] and Σ has countable rank the free monad Σ^* exists and is algebraic. In particular, $\Sigma^*\text{-Alg} \cong \Sigma\text{-Alg}$. By Theorem 4.2, $T_{\mathcal{U}_0}\text{-Alg} \cong \mathbb{K}(\Sigma, \mathcal{U}_0)$. Thus the monads $T_{\mathcal{U}_0}$ and Σ^* coincide. Since Σ is countable, it preserves separability of the metric spaces. Thus, $T_{\mathcal{U}_0}$ does it too.

Let \mathcal{U} be any quantitative theory. Since $\mathcal{U}_0 \subseteq \mathcal{U}$, any model of \mathcal{U} is of course a model of \mathcal{U}_0 . Thus, for any $X \in \mathbf{Met}$, since $(T_{\mathcal{U}}X, \psi_X^{\mathcal{U}}) \in \mathbb{K}(\Sigma, \mathcal{U}_0)$, by Theorem 3.3, there exist a unique homomorphism $h: T_{\mathcal{U}_0}X \rightarrow T_{\mathcal{U}}X$ such that the following diagram commutes

$$\begin{array}{ccccc} X & \xrightarrow{\eta_X^{\mathcal{U}_0}} & T_{\mathcal{U}_0}X & \xleftarrow{\psi_X^{\mathcal{U}_0}} & \Sigma T_{\mathcal{U}_0}X \\ & \searrow \eta^{\mathcal{U}} & \downarrow h & & \downarrow \Sigma h \\ & & T_{\mathcal{U}}X & \xleftarrow{\psi_X^{\mathcal{U}}} & \Sigma T_{\mathcal{U}}X \end{array}$$

In particular, h is non-expansive (hence, continuous).

If X is a separable metric space, then since $T_{\mathcal{U}_0}$ preserves separability, there exists a countable dense subset $D \subseteq T_{\mathcal{U}_0}X$. Because h is a continuous function, then $h(D)$ is a dense subset in $T_{\mathcal{U}}X$. Clearly, $h(D)$ is countable, because D is so. Thus, $T_{\mathcal{U}}$ preserves separability of metric spaces. \square

B Proofs from Sections 5, 7 and 8

This is the proof of Theorem 5.2.

Theorem B.1. *The monads $T_{\mathcal{B}}$ and Π in \mathbf{Met} are isomorphic.*

Proof. In the following we write $\Sigma_{\mathcal{B}}$ and $T_{\mathcal{B}}$ simply as Σ and T , respectively. Similarly, we drop the superscript \mathcal{B} for the relevant natural maps. We need to show that there exist a natural isomorphism $\sigma: T \Rightarrow \Pi$ such that $\sigma \circ \eta = \delta$ and $\sigma \circ \mu = m \circ \sigma$. We define σ at $X \in \mathbf{Met}$ as the unique map σ_X that, by Theorem 3.3, makes the following diagram commute

$$\begin{array}{ccccc} X & \xrightarrow{\eta_X} & TX & \xleftarrow{\psi_X} & \Sigma TX \\ & \searrow \delta_X & \downarrow \sigma_X & & \downarrow \Sigma \sigma_X \\ & & \Pi X & \xleftarrow{\phi_X} & \Sigma \Pi X \end{array}$$

That σ is an isomorphism follows by Theorem 3.3 and Theorem 5.1. The equality $\sigma \circ \eta = \delta$ follows by definition of σ . The equality $\sigma \circ \mu = m \circ \sigma$ follows by Theorem 3.3 and the commutativity of the following diagrams

$$\begin{array}{ccc} \begin{array}{ccccc} T & \xrightarrow{\eta T} & TT & \xleftarrow{\psi T} & \Sigma TT \\ & \searrow id & \downarrow \mu & & \downarrow \Sigma \mu \\ & & T & \xleftarrow{\psi} & \Sigma T \\ \sigma \swarrow & & \downarrow \sigma & & \downarrow \Sigma \sigma \\ & & \Pi & \xleftarrow{\phi} & \Sigma \Pi \end{array} & & \begin{array}{ccccc} T & \xrightarrow{\eta T} & TT & \xleftarrow{\psi T} & \Sigma TT \\ \sigma \downarrow & & \downarrow \sigma \sigma & & \downarrow \Sigma \mu \\ \Pi & \xrightarrow{\delta \Pi} & \Pi \Pi & \xleftarrow{\phi \Pi} & \Sigma \Pi \Pi \\ & \searrow id & \downarrow m & & \downarrow \Sigma m \\ & & \Pi & \xleftarrow{\phi} & \Sigma \Pi \end{array} \end{array}$$

All but one of the squares above commute by definition or by the unit law of monads, whereas the equality $m \circ \phi \Pi = \phi \circ \Sigma m$ follows by definitions of m and ϕ and linearity of Lebesgue integral. \square

This is the proof of Theorem 7.3

Theorem B.2 (Free Algebra). *For any $X \in \mathbf{Met}$, $\tilde{\Sigma}\Pi$ -algebra (A, α) , and non-expansive map $\beta: X \rightarrow A$, there exists a unique $\tilde{\Sigma}\Pi$ -homomorphism $h: P_X \rightarrow A$ making the diagram below commute*

$$\begin{array}{ccccc} X & \xrightarrow{\varrho_X} & P_X & \xleftarrow{\vartheta_X} & \tilde{\Sigma}\Pi P_X \\ & \searrow \beta & \downarrow h & & \downarrow \tilde{\Sigma}\Pi h \\ & & A & \xleftarrow{\alpha} & \tilde{\Sigma}\Pi A \end{array}$$

Proof. Let $X \in \mathbf{Met}$. Given a $\tilde{\Sigma}\Pi$ -algebra (A, α) and the non-expansive map $\beta: X \rightarrow A$, we define the function $h: P_X \rightarrow A$ by induction of the complexity of the structure of the elements in P_X as follows, for arbitrary $x \in X$, $f: n \in \Sigma$, and $\mu_1, \dots, \mu_n \in \Pi(X)$,

$$h(\underline{x}) = \beta(x),$$

$$h(f\langle \mu_1, \dots, \mu_n \rangle) = \alpha(\text{inf}_f(\mu_1 \circ h^{-1}, \dots, \mu_n \circ h^{-1})).$$

The equality $h \circ \varrho_X = \beta$ follows by definition, because $\varrho_X(x) = \underline{x}$.

For arbitrary $f: n \in \Sigma$ and $\mu_1, \dots, \mu_n \in \Pi(P_X)$, we have

$$\begin{aligned} h \circ \vartheta_X(\text{inf}_f(\mu_1, \dots, \mu_n)) &= \\ &= h(f\langle \mu_1, \dots, \mu_n \rangle) && \text{(def. } \vartheta_X) \\ &= \alpha(\text{inf}_f(\mu_1 \circ h^{-1}, \dots, \mu_n \circ h^{-1})) && \text{(def. } h) \\ &= \alpha(\text{inf}_f(\Pi(h)(\mu_1), \dots, \Pi(h)(\mu_n))) && \text{(def. } \Pi) \end{aligned}$$

By couniversality of the coproduct, from the above we get the required equality $h \circ \vartheta_X = \alpha \circ \tilde{\Sigma}\Pi h$. If $h': P_X \rightarrow A$ is another homomorphism such that $h' \circ \varrho_X = \beta$, then the following equalities hold, for arbitrary $x \in X$, $f: n \in \Sigma$, and $\mu_1, \dots, \mu_n \in \Pi(P_X)$,

$$h'(\underline{x}) = \beta(x),$$

$$h'(f\langle \mu_1, \dots, \mu_n \rangle) = \alpha(\text{inf}_f(\Pi(h')(\mu_1), \dots, \Pi(h')(\mu_n))).$$

Then, by an easy induction on the complexity of the structure of the elements in P_X we get that $h = h'$: the base case is trivial; the inductive case follows by

$$\begin{aligned} h'(f\langle \mu_1, \dots, \mu_n \rangle) &= \\ &= \alpha(\text{inf}_f(\Pi(h')(\mu_1), \dots, \Pi(h')(\mu_n))) && \text{(} h' \text{ homomorphism)} \\ &= \alpha(\text{inf}_f(\mu_1 \circ (h')^{-1}, \dots, \mu_n \circ (h')^{-1})) && \text{(def. } \Pi) \\ &= \alpha(\text{inf}_f(\mu_1 \circ h^{-1}, \dots, \mu_n \circ h^{-1})) && \text{(by inductive hp.)} \\ &= h(f\langle \mu_1, \dots, \mu_n \rangle) && \text{(def. } h) \end{aligned}$$

It only remains to prove that h is non-expansive. We proceed by induction. The base case follows trivially by definition h and d_{P_X} , and non-expansiveness of β . As for the inductive case, by definition of d_{P_X} , as the metric d_A is 1-bounded, the only interesting case to check is the following, for $f: \langle n, c \rangle \in \Sigma$

$$\begin{aligned} d_{P_X}(f\langle \mu_1, \dots, \mu_n \rangle, f\langle \nu_1, \dots, \nu_n \rangle) &= \\ &= c \cdot \max_{i=1}^n \mathcal{K}(d_{P_X})(\mu_i, \nu_i) && \text{(def. } d_{P_X}) \\ &\geq c \cdot \max_{i=1}^n \mathcal{K}(d_{P_X})(\mu_i \circ h^{-1}, \nu_i \circ h^{-1}) && \text{(inductive hp.)} \\ &\geq d_A(\alpha(\text{inf}_f(\mu_1, \dots, \mu_n)), \alpha(\text{inf}_f(\nu_1, \dots, \nu_n))) && (\alpha, \text{inf}_f \text{ non-exp.)} \\ &= d_A(h(f\langle \mu_1, \dots, \mu_n \rangle), h'(f\langle \nu_1, \dots, \nu_n \rangle))) && \text{(def. } h) \end{aligned}$$

This concludes the proof. \square

This is the proof of Lemma 7.6 from Subsection 7.2.

Lemma B.3. *The the functor $T_{\mathcal{B}+O(\Sigma)}$ in \mathbf{Met} preserves separability.*

Proof. It is known that Π with Kantorovich metric preserves separability (cf. [27, Theorem 6.18]). For convenience, we briefly recall the argument here. Let X be a separable metric space and $D \subseteq X$ a countable dense subset of X . For any subset $A \subseteq X$, written $\Theta(A)$ the set of Borel probability distributions expressed as finite convex sums of the form $\sum_{i=1}^n q_i \cdot \delta_{a_i}$, where δ_{a_i} is the Dirac distribution at $a_i \in D$, and $\sum_{i=1}^n q_i = 1$ for $q_i \in \mathbb{Q}_{\geq 0}$. Clearly, $\Theta(A) \subseteq \Pi(A)$ and, if A is countable, so is $\Theta(A)$. By a standard limiting argument, if A is dense in X , so is $\Theta(A)$ in $\Pi(X)$ w.r.t. the Kantorovich metric $\mathcal{K}(d_X)$. Therefore $\Theta(D)$ is a countable dense subset of $\Pi(X)$.

Next we show that also $(\tilde{\Sigma}\Pi)^*$ preserves separability. Let X be a separable metric space and $D \subseteq X$ a countable dense subset of X . From Section 7.1, we know that $(\tilde{\Sigma}\Pi)^*X = P_X$. We define the subsets $S_X^n \subseteq P_X$, by induction on $n \in \mathbb{N}$ as follows

- if $d \in D$, then $\underline{d} \in S_X^0$;
- if $\mu_1, \dots, \mu_n \in \Theta(S_X^n)$ and $f: n \in \Sigma$, then $f\langle \mu_1, \dots, \mu_n \rangle \in S_X^{n+1}$,

where $\Theta(-)$ is defined as above. By an easy induction on $n \in \mathbb{N}$, one proves that S_X^n is countable. The base case is trivial. Assume that S_X^n is countable. Then $\Theta(S_X^n)$ is countable and since the signature Σ is assumed to be countable, we have that also S_X^{n+1} is countable.

Consequently, the set $S_X = \bigcup_n S_X^n$ is countable. Moreover, by an easy induction on the complexity of the structure of the elements in P_X , the one shows that any element in P_X is arbitrarily close to some element in S_X w.r.t. the metric d_{P_X} . Hence, $(\tilde{\Sigma}\Pi)^*$ preserves separability.

By Corollary 7.4, $T_{\mathcal{B}+O(\Sigma)} \cong \Pi(\tilde{\Sigma}\Pi)^*$. Thus, $T_{\mathcal{B}+O(\Sigma)}$ preserves separability of the metric spaces. \square

Here is the proof of Lemma 8.6.

Lemma B.4. *The endofunctor Δ on \mathbf{CMet} is locally non-expansive.*

Proof. We need to check that for all $f, g \in \mathbf{CMet}(X, Y)$,

$$\sup_{x \in X} d_Y(f(x), g(x)) \geq \sup_{\mu \in \Delta(X)} \mathcal{K}(d_Y)(\Delta f(\mu), \Delta g(\mu)). \quad (5)$$

Write Φ_Y for the set of non-expansive functions $k: Y \rightarrow [0, 1]$, i.e., those functions such that $\forall y, y'. |k(y) - k(y')| \leq d_Y(y, y')$. Then, for any $\mu \in \Delta(X)$,

$$\begin{aligned} \mathcal{K}(d_Y)(\Delta f(\mu), \Delta g(\mu)) &= \\ &= \sup_{k \in \Phi_Y} \left| \int k \, d\Delta f(\mu) - \int k \, d\Delta g(\mu) \right| \quad (\text{def. } \mathcal{K}(d_Y)) \\ &= \sup_{k \in \Phi_Y} \left| \int k \, d(\mu \circ f^{-1}) - \int k \, d(\mu \circ g^{-1}) \right| \quad (\text{def. } \Delta) \\ &= \sup_{k \in \Phi_Y} \left| \int k \circ f \, d\mu - \int k \circ g \, d\mu \right| \quad (\text{change of var.}) \\ &= \sup_{k \in \Phi_Y} \left| \int (k \circ f) - (k \circ g) \, d\mu \right| \quad (\text{linearity of } \int) \\ &\leq \sup_{k \in \Phi_Y} \int |(k \circ f) - (k \circ g)| \, d\mu \quad (\text{subadd. of } |\cdot|) \\ &\leq \int d_Y \circ \langle f, g \rangle \, d\mu \quad (\text{monotonicity of } \int) \\ &\leq \int \sup_{x \in X} d_Y(f(x), g(x)) \, d\mu \quad (\text{monotonicity of } \int) \\ &= \sup_{x \in X} d_Y(f(x), g(x)). \quad (\mu \text{ probability measure}) \end{aligned}$$

By the above, $\sup_{x \in X} d_Y(f(x), g(x))$ is an upper bound of the set $\{\mathcal{K}(d_Y)(\Delta f(\mu), \Delta g(\mu)) \mid \mu \in \Delta(X)\}$. Hence, this implies (5). \square