

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Authors: Sarkis Manoukian¹, Sally Stewart², Nicholas Graves³, Helen Mason¹, Chris Robertson⁴, Sharon Kennedy⁵, Jiafeng Pan⁴, Kimberley Kavanagh⁴, Lynne Haahr², Mahmood Adil⁶, Stephanie J. Dancer^{7,8}, Brian Cook⁹, Jacqui Reilly²

¹ Yunus Centre for Social Business and Health, Glasgow Caledonian University, Glasgow, UK

² Safeguarding Health through Infection Prevention Research Group, Glasgow Caledonian University, Glasgow, UK

³ Duke-NUS Medical School, Singapore

⁴ Department of Mathematics and Statistics, University of Strathclyde, Glasgow, UK

⁵ HPS Stats Support, Public Health Scotland, Glasgow, UK

⁶ Public Health Scotland, Edinburgh, UK

⁷ Department of Microbiology, Hairmyres Hospital, NHS Lanarkshire, Scotland, UK

⁸ School of Applied Science, Edinburgh Napier University, Edinburgh, UK

⁹ Departments of Anaesthesia and Critical Care, Royal Infirmary of Edinburgh, Edinburgh, UK

Corresponding author:

Sally Stewart
Glasgow Caledonian University
Cowcaddens Road
Glasgow
G4 0BA
Sally.Stewart@gcu.ac.uk

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Structured Summary

Background

Healthcare associated infection (HAI) is associated with increased morbidity and mortality resulting in excess costs.

Aim

To investigate the impact of all types of HAI on the inpatient cost of HAI using different approaches.

Methods

The incidence, types of HAI and excess LOS were estimated using data collected as part of the Evaluation of Cost of Nosocomial Infection (ECONI) study. Scottish NHS reference costs were used to estimate unit costs for bed-days. Variable (cash) costs associated with infection prevention and control (IPC) measures and treatment were calculated for each HAI type and overall. The inpatient cost of HAI is presented in terms of bed-days lost, bed-day costs and cash costs.

Findings

In Scotland 58,010 (95% CI: 41,730-74,840) bed-days were estimated to be lost to HAI during 2018/19 costing £46.4 million (95% CI: £19-£129). The total annual cost in the UK is estimated to be £774 million (95% CI: £328-£2,192). Bloodstream infection and pneumonia were the most costly HAI types per case. Cash costs are a small proportion of the total cost of HAI contributing 2.4% of total costs.

Conclusion

Reliable estimates of the cost burden of HAI management are important for assessing the cost-effectiveness of IPC programmes. This unique study presents robust economic data, demonstrating that HAI remains a burden to the UK NHS and bed-days capture the majority of inpatient costs. These findings can be used to inform the economic evaluation and decision analytic modelling of competing IPC programmes at local and national level.

Keywords:

122:Bloodstream infections

230:Costs

360:Epidemiology

466:Health economics

510:Hospital-acquired infection (HAI)

560:Infection prevention and control

890:Respiratory infections

973:Surgical site infections

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

1000:Survey

1080:Urinary tract infection (UTI)

50: Antibiotics

354: Environmental cleaning

Additional MESH terms

Cross infection

Length of stay

Introduction

Healthcare associated infection (HAI) represents a major issue for health services, patients and public authorities [1]. These potentially avoidable events are associated with increased morbidity and mortality resulting in excess management costs [1-3]. The Evaluation of Cost of Nosocomial Infection (ECONI) study estimated HAI incidence of 0.76% (95% CI: 0.72%-0.81%) with 7,437 (95% CI: 7,021-7,849) HAI cases in teaching and large general hospitals in NHS Scotland in 2018/19 [4].

Many studies have examined the costs associated with HAIs but comparisons among estimates can be challenging due to the differences between settings, patient groups and methods of attributing costs [5]. Few studies have examined the whole hospital incidence and all HAI types. The last study to do this for the UK, Plowman et al [6], found HAI to be a considerable burden, equivalent of 9.1% of the total 1994/5 inpatient acute, obstetric and geriatric programme budget in England. These cost estimates arose from methods that did not account for the time-varying nature of HAI and are likely to have been overestimated [7]. In the intervening period the UK and other countries established national infection prevention and control (IPC) programmes focused on HAI prevention [8, 9]. Since the 1990s there have been significant changes in the age and frailty of the hospital population, complexity of treatments, use of invasive devices, antimicrobial resistance (AMR) and targeted IPC measures, all of which have affected the incidence and impact of HAI. Accurate and current information is needed to optimise decision making when selecting between competing IPC programmes to ensure the most efficient programmes are implemented [10, 11].

The aim of this study was to estimate the costs of different types of HAI in acute hospitals in NHS Scotland using a variety of approaches and extrapolate to the whole of the UK. One of the major costs of HAI can be shown by the number of bed-days released by its prevention. Released hospital beds can be valued in monetary terms by examining how much they historically cost the hospital to supply, the accounting cost. Bed-days include a high proportion of fixed costs and some variable costs [11]. When HAI cases are prevented, releasing bed-days, resources represented by fixed costs can be redeployed for other uses but not recovered while variable costs can be recovered as cash savings. The cost of HAI is

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

presented as natural units of bed-days lost due to HAI and bed-days valued using accounting costs. The accounting costs represent valuations of resources for alternative uses. The cash or variable costs related to the treatment of HAI in NHS hospitals are also calculated separately. The cash costs of treating HAI represent potential savings if HAI cases were prevented [11].

This study presents new information for healthcare decision makers which can inform economic decision making for competing IPC programmes locally and nationally.

Methods

To estimate the inpatient cost of HAI on the health service the incidence rates and estimates of excess length of stay (LOS) derived from the Evaluation of Cost of Nosocomial Infection (ECONI) study were combined [4, 12]. ECONI was a two centre, prospective observational incidence study with a nested case-control study [13]. ECONI took place in one teaching hospital and one large general hospital in NHS Scotland in 2018/2019. The cost of HAI to the NHS acute care sector was estimated by taking the following steps. First, the expected number of annual HAI cases were identified. Second, the extra resources used to treat the average case were identified. Third, those resources used for the treatment of HAI were valued in monetary terms. The costs from the ECONI study were extrapolated to produce an estimate for the UK.

Expected annual HAI cases

The expected number and types of annual HAIs were estimated using data collected as part of the ECONI study [4]. ECONI reported the incidence of HAI using standard case definitions of the European centre for disease prevention and control (ECDC) [13]. Incidence rates of HAI per 100,000 acute-occupied bed-days (AOBDs) in the study hospitals were used to estimate HAI incidence in NHS Scotland. National estimates of HAIs were derived by applying the incidence within each specialty group to the total annual overnight admissions within NHS Scotland [4]. The nested case-control sample [14] included approximately 5% of the HAI cases and was used to calculate the cash cost of antibiotic treatment only (see appendix).

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Resources used to manage the average HAI case

HAI cost is mainly driven by excess LOS in the hospital [15]. During these extra days in hospital HAI patients consume a mix of resources represented by fixed and variable costs. Fixed costs do not change with the number of patients treated and, in the NHS, represent buildings and staff, variable costs change with the number of patients treated and include consumables such as drugs and disposable equipment. Excess LOS due to HAI estimates were produced for each HAI type and all HAI using a multistate modelling approach which took account of time-varying exposures and the competing risks of death and discharge [16, 17]. The multistate model estimated excess LOS, for each HAI type, that could have been spent in any clinical area inclusive of intensive care unit (ICU), high dependency unit (HDU) or other wards. The methods and results for the multistate model are described in detail elsewhere [12]. Estimates of annual bed-days lost to HAI depend on both estimated incidence and excess LOS due to HAI [4, 12].

Valuing resources used in the treatment of HAI in monetary terms

To value bed-days lost due to HAI, the “total” and “direct” accounting unit costs were calculated. These are mostly fixed costs and are based on a top-down valuation approach with gross-costing methodology [18]. Total costs include capital, overheads, staff, pharmacy, and laboratory costs. The total and direct costs were calculated using routinely reported historical data from NHS Scotland [19]. Direct costs include staff and consumable costs outlined above and exclude capital and overhead costs. Direct costs can be directly attributed to patient care and have been used previously to estimate the costs of treating HAI [20-25]. Unit cost estimates were weighted by the total number of discharges across all specialties from April 2017 to March 2018 in Scottish teaching hospitals and large general hospitals (see appendix). Unit costs were aggregated across ICUs, HDUs and other wards to match the estimates of the multistate model [12]. Unit cost raw data were assumed to be log-normally distributed. In cost estimation confidence intervals were calculated by taking 5,000 Monte Carlo draws from log-normal distributions fitted to the unit cost data. Bed-day unit costs were based on Scottish reference costs [26] and are presented in Table I. Standard deviations in Table I reflect the substantial variability in bed-day costs across specialties and hospitals in Scotland.

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

The excess variable or cash costs incurred for each type of HAI was also calculated separately. These cash costs excluded fixed staff costs and were grouped into diagnostic tests, IPC consumables (including cleaning materials and antibiotic drugs). Diagnostic tests were defined based on the HAI type and causative organisms reported in the ECONI study [4]. IPC and cleaning consumable costs were calculated for each level of transmission-based precautions (TBP) required for treatment. This included mapping the proportion of each HAI type which required use of TBP based on the Scottish National IPC manual and consultation with IPC experts [27]. The unit costs of IPC consumables were based on a mix of existing literature and expert opinion (see appendix for full details).

Data on antibiotic prescribing in hospital was collected from the nested case-control sample of the ECONI study. The British National Formulary (BNF) was used to cost antibiotics [28]. The costs were recorded in the week after data collection was completed. Cash costs were multiplied by the estimated annual number of HAIs to calculate cash treatment by HAI type and a total. All cash costs were calculated in 2019 Pounds (£), more detail is presented in the appendix.

UK Extrapolation of bed-day HAI costs

The ECONI study estimated incidence 0.76% (95% CI:0.72%-0.81%) was applied to the UK total elective and emergency patient admissions in the financial year April 2017 to March 2018 to produce national estimates of costs [29-32]. This extrapolation assumes that the two hospitals of the study were representative of other acute hospitals in Scotland and the UK in terms of specialties, excess LOS and bed-day unit costs [19].

Estimation of lost bed-days and costs due to HAI

Estimates of annual bed-days lost, for each HAI type and overall HAI, along with bias corrected confidence intervals, using 5,000 non-parametric bootstrap samples, were based on incidence and excess LOS estimated elsewhere [4, 12]. In this study the annual and per case cost of HAI is defined as the product of vectors of unit costs and excess resources consumed due to HAI. Point estimates of annual and per case bed-day costs, for each HAI type and overall HAI, were calculated using annual and per case bed-days lost multiplied by bed-day unit costs and excess LOS multiplied by bed-day unit costs, respectively. To

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

estimate 95% confidence intervals of annual and per case costs 5,000 Monte Carlo simulations were drawn from log-normal distributions fitted to the data shown in Table I [17].

Cash costs were calculated separately using a different analytical process (see appendix). Due to the way cash costs were calculated point estimates of the unit costs were used in the calculation of cash costs. Therefore, no confidence intervals are provided for cash costs related to IPC and treatment of HAI.

Results

The incidence rate and average excess LOS from the ECONI study and total bed-days lost by HAI type for teaching and general hospitals in Scotland are shown in Table I Bed-day unit costs

Unit cost	Mean	SD ¹	Distribution	Source
1) Bed-day total cost	£799.17	£535.37	Log-normal	ISD[26]
2) Bed-day direct cost	£519.38	£344.27	Log-normal	ISD[26]
Notes 1 Standard deviation				

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table II. These are based on the total annual admissions to teaching and general hospitals in NHS Scotland in 2018/19.

Bloodstream infection (BSI) was the HAI type with the greatest total annual impact in terms of bed-days lost followed by pneumonia (PN) and surgical site infection (SSI). Urinary tract infection (UTI) was shown to have a negative impact on LOS in the multistate model. In practice this means that on average UTI did not increase LOS and it was assumed that there were zero additional days for these patients. In total there were 58,010 bed-days (95% CI: 41,730-74,840) lost in a year due to all HAI types.

Cost per case and annual inpatient costs in Scotland are shown in Table I [Bed-day unit costs](#)

Unit cost	Mean	SD ¹	Distribution	Source
3) Bed-day total cost	£799.17	£535.37	Log-normal	ISD[26]
4) Bed-day direct cost	£519.38	£344.27	Log-normal	ISD[26]
Notes 1 Standard deviation				

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table II: Incidence, average excess LOS and bed-days lost by HAI type in Scotland

HAI	Incidence Rates per 100,000 AOBs (95% CI) ¹	Estimated annual number of HAI cases (95% CI) ¹	Average Excess LOS ² per HAI (95% CI) ¹	Total annual bed-days lost to HAI ³ (N)	(95% CI) ¹
BSI	35.0 (31.3-39.5)	1,389 (1,245-1,570)	11.4 (5.8-17.0)	15,830	(7,550-23,950)
GI	31.6 (28.2-36.3)	1,256 (1,122-1,440)	6.0 (-0.7-12.7)	7,540	(0 -16,100)
LRI	26.2 (3.6-30.7)	1,041 (937-1,218)	7.3 (1.8-12.7)	7,600	(1,300 -13,540)
PN	15.9 (13.7 -19.2)	630 (544-764)	16.3 (7.5-25.5)	10,270	(4,170-16,380)
SSI	25.8 (22.8-30.5)	1,023 (904-1,210)	9.8 (4.5-15.0)	10,030	(4,190-15,900)
UTI	41.0 (36.6- 46.2)	1,628 (1,454-1,836)	-1.0 (-4.3-2.3)	0	(0-4,180)
Other	12.0 (9.5-15.8)	475 (378-628)	14.0 (-3.9-31.8)	6,650	(0-16,360)
All HAI	187.2 (176.8-197.6)	7,437 (7,021-7,849)	7.8 (5.7-9.9)	58,010	(41,730-74,840)

Notes: (BSI) Blood Stream Infection; (GI) Gastrointestinal Infection; (LRI) Lower Respiratory tract Infection; (PN) Pneumonia; (SSI) Surgical Site Infection; (UTI) Urinary Tract Infection; Other are small number infections – includes (SST) Skin Soft Tissue; (BJ) Bone and Joint; (CV) Cardiovascular; (EENT) Eye, Ear, Nose and Throat; and (SI) systemic infections.
1 95% confidence interval.
2 Excess LOS is expressed in additional days per case.
3 NHS Scotland acute hospitals.

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table III. The total cost attributable to HAI treatment annually in NHS Scotland is estimated to be £46.35 (95% CI: £19.4-£128.8) million pounds. The direct cost due to HAI is estimated to be £30.11 (95% CI: £14.1-£74.4) million pounds per year.

The extrapolation of the costs for the whole of the UK is shown in Table I Bed-day unit costs

Unit cost	Mean	SD ¹	Distribution	Source
5) Bed-day total cost	£799.17	£535.37	Log-normal	ISD[26]
6) Bed-day direct cost	£519.38	£344.27	Log-normal	ISD[26]
Notes 1 Standard deviation				

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table II: Incidence, average excess LOS and bed-days lost by HAI type in Scotland

HAI	Incidence Rates per 100,000 AOBs (95% CI) ¹	Estimated annual number of HAI cases (95% CI) ¹	Average Excess LOS ² per HAI (95% CI) ¹	Total annual bed-days lost to HAI ³ (N)	(95% CI) ¹
BSI	35.0 (31.3-39.5)	1,389 (1,245-1,570)	11.4 (5.8-17.0)	15,830	(7,550-23,950)
GI	31.6 (28.2-36.3)	1,256 (1,122-1,440)	6.0 (-0.7-12.7)	7,540	(0 -16,100)
LRI	26.2 (3.6-30.7)	1,041 (937-1,218)	7.3 (1.8-12.7)	7,600	(1,300 -13,540)
PN	15.9 (13.7 -19.2)	630 (544-764)	16.3 (7.5-25.5)	10,270	(4,170-16,380)
SSI	25.8 (22.8-30.5)	1,023 (904-1,210)	9.8 (4.5-15.0)	10,030	(4,190-15,900)
UTI	41.0 (36.6- 46.2)	1,628 (1,454-1,836)	-1.0 (-4.3-2.3)	0	(0-4,180)
Other	12.0 (9.5-15.8)	475 (378-628)	14.0 (-3.9-31.8)	6,650	(0-16,360)
All HAI	187.2 (176.8-197.6)	7,437 (7,021-7,849)	7.8 (5.7-9.9)	58,010	(41,730-74,840)

Notes: (BSI) Blood Stream Infection; (GI) Gastrointestinal Infection; (LRI) Lower Respiratory tract Infection; (PN) Pneumonia; (SSI) Surgical Site Infection; (UTI) Urinary Tract Infection; Other are small number infections – includes (SST) Skin Soft Tissue; (BJ) Bone and Joint; (CV) Cardiovascular; (EENT) Eye, Ear, Nose and Throat; and (SI) systemic infections.
1 95% confidence interval.
2 Excess LOS is expressed in additional days per case.
3 NHS Scotland acute hospitals.

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table III: Cost per case for each HAI type and annual cost of HAI in NHS Scotland

HAI	Cost per case for each HAI type and overall				Annual cost in NHS Scotland (Million)			
	Total cost per case		Direct cost per case		Total cost		Direct cost	
	Mean	(95% CI) ¹	Mean	(95% CI) ¹	Mean	(95% CI) ¹	Mean	(95% CI) ¹
BSI	£9,109	(£3,511-£28,210)	£5,917	(£2,552-£15,438)	£12.65	(£4.82-£38.96)	£8.22	(£3.45-£21.94)
GI	£4,794	(£445-£19,835)	£3,114	(£192-£10,401)	£6.02	(£0.52-£24.83)	£3.91	(£0.24-£14.11)
LRI	£5,833	(£1,729-£20,019)	£3,789	(£1,234-£11,684)	£6.07	(£1.66-£21.25)	£3.94	(£1.17-£11.84)
PN	£13,024	(£4,808-£45,061)	£8,460	(£3,432-£23,548)	£8.20	(£2.99-£25.88)	£5.33	(£2.01-£14.44)
SSI	£7,830	(£2,987-£24,993)	£5,086	(£2,095-£14,433)	£8.01	(£2.95-£27.02)	£5.20	(£1.96-£13.45)
UTI	£0	(£0-£2,109)	£0	(£0-£1,304)	£0	(£0-£3.63)	£0	(£0-£2.21)
Other	£11,186	(£0-£45,319)	£7,266	(£0-£26,523)	£5.31	(£0.05-£24.45)	£3.45	(£0-£12.77)
All HAI	£6,232	(£2,733-£18,181)	£4,048	(£1,927-£9,591)	£46.35	(£19.43-£128.81)	£30.11	(£14.12-£74.46)

Notes: (BSI) Blood Stream Infection; (GI) Gastrointestinal Infection; (LRI) Lower Respiratory tract Infection; (PN) Pneumonia; (SSI) Surgical Site Infection; (UTI) Urinary Tract Infection; Other are small number infections – includes (SST) Skin Soft Tissue; (BJ) Bone and Joint; (CV) Cardiovascular; (EENT) Eye, Ear, Nose and Throat; and (SI) systemic infections.
¹ 95% confidence interval.

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table IV. Total costs in the UK based on total admissions are estimated to be £774 million (95% CI: £328 million-£2.2 billion) annually. Direct costs are estimated to be £503 million (95% CI: £236 million-£1.2 billion) per year.

The costs of consumable items which could be saved in the short term are shown in Table I
Bed-day unit costs

Unit cost	Mean	SD ¹	Distribution	Source
7) Bed-day total cost	£799.17	£535.37	Log-normal	ISD[26]
8) Bed-day direct cost	£519.38	£344.27	Log-normal	ISD[26]
Notes 1 Standard deviation				

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table II: Incidence, average excess LOS and bed-days lost by HAI type in Scotland

HAI	Incidence Rates per 100,000 AOBs (95% CI) ¹	Estimated annual number of HAI cases (95% CI) ¹	Average Excess LOS ² per HAI (95% CI) ¹	Total annual bed-days lost to HAI ³ (N)	(95% CI) ¹
BSI	35.0 (31.3-39.5)	1,389 (1,245-1,570)	11.4 (5.8-17.0)	15,830	(7,550-23,950)
GI	31.6 (28.2-36.3)	1,256 (1,122-1,440)	6.0 (-0.7-12.7)	7,540	(0 -16,100)
LRI	26.2 (3.6-30.7)	1,041 (937-1,218)	7.3 (1.8-12.7)	7,600	(1,300 -13,540)
PN	15.9 (13.7 -19.2)	630 (544-764)	16.3 (7.5-25.5)	10,270	(4,170-16,380)
SSI	25.8 (22.8-30.5)	1,023 (904-1,210)	9.8 (4.5-15.0)	10,030	(4,190-15,900)
UTI	41.0 (36.6- 46.2)	1,628 (1,454-1,836)	-1.0 (-4.3-2.3)	0	(0-4,180)
Other	12.0 (9.5-15.8)	475 (378-628)	14.0 (-3.9-31.8)	6,650	(0-16,360)
All HAI	187.2 (176.8-197.6)	7,437 (7,021-7,849)	7.8 (5.7-9.9)	58,010	(41,730-74,840)

Notes: (BSI) Blood Stream Infection; (GI) Gastrointestinal Infection; (LRI) Lower Respiratory tract Infection; (PN) Pneumonia; (SSI) Surgical Site Infection; (UTI) Urinary Tract Infection; Other are small number infections – includes (SST) Skin Soft Tissue; (BJ) Bone and Joint; (CV) Cardiovascular; (EENT) Eye, Ear, Nose and Throat; and (SI) systemic infections.
1 95% confidence interval.
2 Excess LOS is expressed in additional days per case.
3 NHS Scotland acute hospitals.

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table III: Cost per case for each HAI type and annual cost of HAI in NHS Scotland

HAI	Cost per case for each HAI type and overall				Annual cost in NHS Scotland (Million)			
	Total cost per case		Direct cost per case		Total cost		Direct cost	
	Mean	(95% CI) ¹	Mean	(95% CI) ¹	Mean	(95% CI) ¹	Mean	(95% CI) ¹
BSI	£9,109	(£3,511-£28,210)	£5,917	(£2,552-£15,438)	£12.65	(£4.82-£38.96)	£8.22	(£3.45-£21.94)
GI	£4,794	(£445-£19,835)	£3,114	(£192-£10,401)	£6.02	(£0.52-£24.83)	£3.91	(£0.24-£14.11)
LRI	£5,833	(£1,729-£20,019)	£3,789	(£1,234-£11,684)	£6.07	(£1.66-£21.25)	£3.94	(£1.17-£11.84)
PN	£13,024	(£4,808-£45,061)	£8,460	(£3,432-£23,548)	£8.20	(£2.99-£25.88)	£5.33	(£2.01-£14.44)
SSI	£7,830	(£2,987-£24,993)	£5,086	(£2,095-£14,433)	£8.01	(£2.95-£27.02)	£5.20	(£1.96-£13.45)
UTI	£0	(£0-£2,109)	£0	(£0-£1,304)	£0	(£0-£3.63)	£0	(£0-£2.21)
Other	£11,186	(£0-£45,319)	£7,266	(£0-£26,523)	£5.31	(£0.05-£24.45)	£3.45	(£0-£12.77)
All HAI	£6,232	(£2,733-£18,181)	£4,048	(£1,927-£9,591)	£46.35	(£19.43-£128.81)	£30.11	(£14.12-£74.46)

Notes: (BSI) Blood Stream Infection; (GI) Gastrointestinal Infection; (LRI) Lower Respiratory tract Infection; (PN) Pneumonia; (SSI) Surgical Site Infection; (UTI) Urinary Tract Infection; Other are small number infections – includes (SST) Skin Soft Tissue; (BJ) Bone and Joint; (CV) Cardiovascular; (EENT) Eye, Ear, Nose and Throat; and (SI) systemic infections.
¹ 95% confidence interval.

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table IV Annual total inpatient cost of HAI in the UK

Annual total cost in the UK (£Million per Year) ²		Annual direct cost in the UK (£Million per Year) ²	
Mean	(95% CI) ¹	Mean	(95% CI) ¹
£774	(£328-£2,192)	£503	(£236-£1,217)
Notes: 1 (95% confidence interval). 2 Annual UK total and direct costs are based on ECONI HAI incidence: 0.76% (95% CI: 0.72%-0.81%).			

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table V. The total cost of these items varies by HAI type. These costs are small in comparison to staff time and other fixed resources. The total annual estimate is approximately £1.1 million pounds across acute hospitals in the Scottish NHS. Cash costs are approximately 2.4% of the estimated total cost of HAI and 3.7% of the direct cost of HAI in Scotland.

Discussion

The ECONI study is the first in the UK to report whole hospital incidence of HAI since Plowman et al which was based on data now over 20 years old [6]. It was estimated that in Scotland 58,010 (95% CI: 41,730-74,840) bed-days were lost due to HAI each year. These occupied beds are the main economic cost arising from HAI. The impact of lost bed-days in a system like the NHS results in an increase in waiting lists which means potential patients are delayed in accessing services. This size of the loss equates to 159 beds or a small general hospital of 180 beds (assuming average bed occupancy of 88%) being occupied for a whole year with HAI cases. Alternatively, the impact of HAI can be shown by the number of elective patients who could have been treated after a 10% reduction in HAI rates. Given that LOS for elective patients in NHS Scotland is on average 3.4 days [33] then 1,700 additional elective patients could be treated annually if 10% of HAI is prevented. Estimated lost bed-days represent the opportunity cost of HAI expressed in natural units. The opportunity cost of a bed-day can be defined as the forgone benefit of the best alternative use of the resource such as treating another patient. In the NHS high demand means a released bed is very valuable since a new patient will always occupy it but in situations of spare capacity the value would be lower if a bed remains empty. Bed-days are a unit of currency familiar to healthcare decision makers who manage resources with a high proportion of fixed costs.

The total cost valuation of the lost bed-days due to HAI was £46.35 million (95% CI: £19m-£129m) and the direct cost was £30.11 million (95% CI: £14m-£75m). In 2018-19 the total acute inpatient expenditure in Scotland was approximately £4 billion [34]; this sets the total cost of HAI at around 1.1% of the total acute inpatient expenditure in NHS Scotland. These are mostly fixed costs that cannot be recovered as savings if HAI is prevented since they are based on historical accounting unit costs. However, it is still important to consider costs in this way so that comparisons can be made with other NHS spending that is reported in

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

similar terms. These costs also represent a monetary valuation of benefits from alternative uses of these resources when HAI is prevented. Direct costs are a representation of staff and consumable resources that could have been used to treat other patients if HAI rates were reduced.

The top three HAI types which had the greatest cost impact were BSI, pneumonia and SSI (Table III). HAI cost is estimated as a combination of incidence and excess LOS and the rank order of HAI type changes according to the impact on either incidence or LOS. For example, according to ECONI incidence, UTIs are the most common infections in Scotland but these are usually easily treated and thus have minimal impact on inpatient resources. This does not mean IPC teams should disinvest in UTI prevention since these high incidence infections frequently require antibiotic treatment, thus contributing to AMR. UTI were also found to be one of the most common primary infection types leading to secondary BSI [4]. BSI is the second most common HAI type and has a relatively large impact on LOS (11.4 additional days on average). BSI is very costly and targeting investment in IPC measures to reduce this type of HAI is likely to be cost-effective. Pneumonias also have serious consequences for the patients and can be difficult to treat increasing LOS by more than 16 days on average, but they were less frequent than other HAIs. Nevertheless, IPC measures to prevent pneumonias could represent a good use of resources given the high cost per case of these infections. Outbreaks that would cause an increase in viral respiratory tract infection can be very costly for the health service and vigilance against these is required. SSIs have a lower cost per case than pneumonias but are more frequent making these the second most costly HAI type annually in Scotland.

The cost of HAI was extrapolated to the whole of the UK and is estimated to be at £774 million (95% CI: £328m-£2,192m) and direct costs at £503 million (95% CI: £236m-£1,217m) annually. The total cost of HAI is estimated to be at 0.62% of the NHS total spending in the four UK nations in 2018 and is associated with approximately 968,000 lost bed-days annually. This is much lower than the 9.1% and 3.64 million lost bed-days estimated by Plowman [6] more than 20 years ago. Since the 1990s changes in the hospital population, treatments, use of devices, antimicrobial resistance and IPC measures have affected the incidence and impact of HAI on the health service. The extrapolation results in this study

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

should be interpreted with caution since they assume a similar mix of hospitals, specialties and severity of illness across the UK.

While total costs may represent historical spending to supply these beds, they do not necessarily represent their value in terms of opportunity costs especially in situations of spare capacity. Opportunity costs are a more useful measure for decision making under conditions of scarce resources but estimating these accurately can be difficult [35]. There have been attempts to estimate the opportunity cost of a bed-day using willingness-to-pay (WTP) surveys in other countries [36, 37]. For example, Page et al [37] estimated opportunity costs using a WTP survey, for bed-days released by IPC programmes, elicited from CEOs of acute hospitals in Australia. These estimates suggest that the total accounting cost to opportunity cost ratio is approximately equal to 3.7 which would imply an economic value of £216 for each bed-day in this study. A bed-day cost at this level would result in yearly opportunity costs in Scotland of up to £16 million which would fall at the lower end of the estimated direct cost confidence interval (95% CI: £14-£75 million). The true economic value or opportunity cost of the occupied bed-day in teaching and general hospitals in the UK is unknown and using WTP estimates from another country or other types of hospitals would introduce sources of uncertainty. WTP can over or underestimate the cost depending on demand for hospital beds and estimates would be much higher in an infection outbreak or pandemic. In a WTP study in European tertiary hospitals Stewardson et al recognised that a limitation of this approach is that WTP measures purchase intention not actual cost [36]. Additionally, they found that WTP values depended on the type of hospital reimbursement [36]. This is also reflected in the findings of other studies who found that the hospital remuneration process had implications in how hospital executives view infection prevention [38, 39].

This study values resource lost to HAI in total costs and direct costs using estimates of HAI incidence and excess LOS that are based on methodologies that minimise bias [4, 12]. Most UK hospitals are operating close to full capacity [40, 41] with long waiting lists which suggests that when HAI is prevented resources could be readily redeployed to treat other patients. This means the valuation of HAI cost in this study represents resources with alternative uses. In addition, this study allows a comparison of the cost of different HAI types and at the same time contrasts incidence with resource use which is relevant to IPC

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

planning. Another strength of this study is the probabilistic methodology that combined estimates of HAI incidence, excess LOS and bed-day unit costs and accounted for all types of uncertainty in the parameters. Previously in this literature studies have reported costs without fully accounting for all sources of uncertainty [38, 42-45].

This study has several limitations. The study was not conducted using a bottom-up micro-costing approach [18]. An example of this approach is the patient level information costing system (PLICS) but this was not routinely available during the study period. PLICS is a patient-level approach which can be used to standardise the method of reporting cost information across the health service. PLICS has the potential to be more accurate than gross-costing methods but is still limited by unit costs that do not reflect opportunity costs and staff costing may occasionally be based on averages resembling a gross-costing methodology. A study in an English NHS hospital using PLICS to report the economic burden of SSI reported an attributable cost of SSI of £5,239 (95% CI: £4,622-£6,719) [38]. This study estimates that the attributable direct hospital cost of SSI was £5,086 (95% CI 2,095-14,433). Another limitation of this study is that the multistate model estimates of excess LOS do not distinguish between facilities that have completely different cost profiles. For example, if 10% of the excess LOS took place in an ICU bed this would increase estimated costs by approximately 14%. Heister et al in a study of *C. difficile* show that multistate models may also underestimate excess LOS which means the results of this study could be a conservative estimate of the true cost of HAI [46]. Furthermore, since antibiotic costs are based on a smaller sample these costs may not be directly comparable with the rest of the results.

Bed-day costs reported previously in the literature mostly relate to capital costs, overheads and fixed contracts that do not change in the short term [11]. Since the cost of facilities and staff are fixed under normal circumstances there is small potential for any annual saving which could be achieved by preventing HAI. Any savings would relate to the consumable items which are purchased as they are consumed. Gastrointestinal infections (GI) and pneumonias had the highest IPC and laboratory cash costs per case with GIs having the greatest annual impact. This study shows that pneumonias are very resource intensive both in terms of fixed costs and variable costs. While the results of this study have shown that Scottish annual cash costs for all HAI types are relatively small at approximately £1.1 million per year, this money could eventually be reinvested if HAI cases were prevented. Plowman

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

et al [6] suggested that cash costs are 11% of the total costs. The results of this study suggested that cash costs are a much smaller proportion of costs at 2.4% of the total and 3.8% of direct costs. This means that any expected cash savings from HAI prevention would be very small. Preventing HAI may only release relatively small funds as cash savings but there is an additional benefit, which is not included in these monetary valuations, of reducing antibiotic consumption [47]. The human cost of HAI should also be taken into account when evaluating HAI prevention since it is also not included in these monetary costs.

A recent meta-analysis concluded that HAI reduction of between 35%-55% may be possible with the implementation of multifaceted interventions [48]. However, preventable proportions of HAI may decrease over time as IPC initiatives are successfully implemented. Reductions shown in some of the earlier studies within this meta-analysis would not be achievable today as many of the proposed interventions have been implemented in some way. It is likely that realistic reductions in overall HAI numbers in Scotland would be lower than Schreiber et al [48] report especially given the low proportion of surgical and device related HAIs identified in the ECONI study [4]. Exaggerated estimates of the cost of HAI and the potential benefits from prevention of HAI may lead policymakers to be disappointed when IPC programmes turn out to be less efficient than expected.

While considerable progress has been made in evaluating IPC measures with cost-effectiveness modelling approaches, these analyses rely heavily on unbiased information [49]. When seeking funding, “attention grabbing” costs are frequently quoted in order to convince policymakers to invest in IPC measures among other interventions. Within the literature there are also attempts to compare the cost of HAI with other conditions, e.g. cancers, with the intention to focus policymakers’ attention on IPC and HAI prevention. However, as discussed elsewhere, [50] it is patients with these conditions who are most at risk of developing HAI. Rather than this competition for resources among patient groups, studies should include integration of IPC measures in order to maximise the efficacy of existing treatments for those most at risk of developing HAI. This study reports the cost of HAI using a variety of measures that can reliably inform future economic evaluations of IPC programmes.

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Conflict of interest statement

None declared.

Funding source

This work was supported by National Services Scotland NHS Health Protection Scotland Programme (RIE reference 14-154) from October 2015 to January 2020. The work was also pump primed by the Scottish Healthcare Associated Infection Prevention Institute (SHAIPi), which has been set up with Scottish Government funding via the Scottish Infection Research Network.

Author contributions

SM contributed to aspects of the study design and led data collection for economic analysis, calculated the total and direct costs and prepared the manuscript. HM contributed to the study design, led on health economic aspects of the study design and contributed to the manuscript. SS led the study design, wrote study protocols and ethics and PBPP approvals, contributed to the development of statistical analysis, the manuscript and costed the cash treatment costs. NG prepared the manuscript and contributed to the study design. CR contributed to the concept of the study, study design, statistical analysis plan and the manuscript. KK, SK, JP undertook the statistical analysis and prepared results. MA and LH contributed to the final manuscript. SD and BC are the Principal Investigators at the recruiting sites and contributed to the final manuscript. JR conceived the study, is Chief Investigator for the study and contributed to the manuscript.

Non Author Collaborators

The ECONI Steering Committee. MA represented the funder, AL, RD, AM, MS and JI represented the Scottish Government HAI policy unit on the Steering Committee. ER and LR represented Infection Prevention Society (IPS). MW, LB and MR were lay representatives on the Steering Committee and MW and LB contributed to the development of the patient facing materials for the study.

Professor Jacqui Reilly (JR)

Professor Mahmood Adil (MA)

Dr Helen Mason (HM)

Professor Chris Robertson (CR)

Professor Nick Graves (NG)

Josephine Ives (JI)

Margaret Syme (MS)

Rachel Dunk (RD)

Abigail Mullings (AM)

Elaine Ross (ER)

Professor Stephanie Dancer (SD)

Dr Brian Cook (BC)

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Professor Alistair Leonard (AL)

Moira Whyte (MW)

Maureen Rodgers (MR)

Louise Brown (LB)

Sally Stewart (SS)

Acknowledgements: Data collection could not have been completed without the enormous effort from the ECONI research nurses: Norma Williamson, Helen Thain, Rosalind Boyle, Audrey McAlpine and Claire Beith. Thanks are extended to all members of the Infection Prevention and Control teams, Microbiology departments and Research and Development teams in the study sites for their support in initiating the study. The study was supported by the clinical staff in the study hospitals in particular the infection control teams, laboratory staff, and research and development teams. The authors would like to specifically thank ER and LR for their advice and guidance on the costing of the IPC measures.

Ethics: this study was surveillance and as such was confirmed as ineligible for ethical review (Personal communication A. Bailey¹). The case control study component of the study received a favourable ethical opinion from the Scotland A Research Ethics Committee on 3rd March 2017 (Reference 16/SS/0199). It was approved by national information governance approvals: Public Benefit of Health and Social Care: Incidence study: 1617-0037

¹ Bailey, A., South East Scotland Research Ethics Service Letter, S. Stewart, Editor. 2016.

References

- [1] World Health Organization. Report on the burden of endemic health care-associated infection worldwide. Geneva: World Health Organization; 2011. Available at: <http://apps.who.int/iris/handle/10665/80135> [last accessed 18 March 2020].
- [2] Zimlichman E, Henderson D, Tamir O *et al*. Health care-associated infections: a meta-analysis of costs and financial impact on the US health care system. *JAMA Intern Med* 2013; **173**: 2039-46. doi: 10.1001/jamainternmed.2013.9763
- [3] Umscheid C, Mitchell M, Doshi J, Agarwal R, Williams K, Brennan P. Estimating the Proportion of Healthcare-Associated Infections That Are Reasonably Preventable and the Related Mortality and Costs. *InfectControl HospEpidemiol* 2011; **32**: 101-14. doi: 10.1086/657912
- [4] Stewart S, Robertson C, Pan J *et al*. ECONI: Epidemiology of Healthcare Associated Infection. Is there a need for a new Infection Prevention and Control strategy? *J Hosp Infect*; **In this issue**:
- [5] Manoukian S, Stewart S, Dancer S *et al*. Estimating excess length of stay due to healthcare-associated infections: a systematic review and meta-analysis of statistical methodology. *J Hosp Infect* 2018; **100**: 222-35. doi: 10.1016/j.jhin.2018.06.003
- [6] Plowman R, Graves N, Griffin MA *et al*. The rate and cost of hospital-acquired infections occurring in patients admitted to selected specialties of a district general hospital in England and the national burden imposed. *JHospInfect* 2001; **47**: 198-209. doi: 10.1053/jhin.2000.0881
- [7] Beyersmann J, Kneib T, Schumacher M, Gastmeier P. Nosocomial infection, length of stay, and time-dependent bias. *Infect Control Hosp Epidemiol* 2009; **30**: 273-6. doi: 10.1086/596020
- [8] Public Health England. Healthcare associated infections (HCAI): guidance, data and analysis. 2014. Available at: <https://www.gov.uk/government/collections/healthcare-associated-infections-hcai-guidance-data-and-analysis> [last accessed 27 May 2020].
- [9] Raza S. Healthcare Associated Infections. 2011. Available at: http://www.parliament.scot/ResearchBriefingsAndFactsheets/Factsheets/SB_11-80.pdf.
- [10] Graves N. Make economics your friend. *J Hosp Infect* 2018; **100**: 123-9. doi: 10.1016/j.jhin.2018.07.008
- [11] Graves N, Harbarth S, Beyersmann J, Barnett A, Halton K, Cooper B. Estimating the cost of health care-associated infections: mind your p's and q's. *Clin Infect Dis* 2010; **50**: 1017. doi: 10.1086/651110 [doi]
- [12] Stewart S, Robertson C, Pan J *et al*. ECONI: Impact of Healthcare Associated Infection on Length of Stay. *J Hosp Infect*; **In this issue**:
- [13] European Centre for Disease Prevention and Control. Point prevalence survey of healthcare-associated infections and antimicrobial use in European acute care hospitals – protocol version 5.3. Stockholm: ECDC; 2016. Available at: <http://ecdc.europa.eu/en/publications/Publications/0512-TED-PPS-HAI-antimicrobial-use-protocol.pdf>.
- [14] Stewart S, Robertson C, Manoukian S *et al*. How do we evaluate the cost of nosocomial infection? The ECONI protocol: an incidence study with nested case-

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

- control evaluating cost and quality of life. *BMJ Open* 2019; **9**: e026687. doi: 10.1136/bmjopen-2018-026687
- [15] Barnett AG, Batra R, Graves N, Edgeworth J, Robotham J, Cooper B. Using a longitudinal model to estimate the effect of methicillin-resistant *Staphylococcus aureus* infection on length of stay in an intensive care unit. *American Journal of Epidemiology* 2009; **170**: 1186. doi: 10.1093/aje/kwp249 [doi]
- [16] Wolkewitz M, von Cube M, Schumacher M. Multistate Modeling to Analyze Nosocomial Infection Data: An Introduction and Demonstration. *Infect Control Hosp Epidemiol* 2017; **38**: 953-9. doi: 10.1017/ice.2017.107
- [17] Stewardson AJ, Allignol A, Beyersmann J *et al*. The health and economic burden of bloodstream infections caused by antimicrobial-susceptible and non-susceptible Enterobacteriaceae and *Staphylococcus aureus* in European hospitals, 2010 and 2011: a multicentre retrospective cohort study. *Euro Surveill* 2016; **21**: pii=30319. doi: 10.2807/1560-7917.es.2016.21.33.30319
- [18] Spacirova Z, Epstein D, Garcia-Mochon L, Rovira J, Olry de Labry Lima A, Espin J. A general framework for classifying costing methods for economic evaluation of health care. *Eur J Health Econ* 2020. doi: 10.1007/s10198-019-01157-9
- [19] Information Services Division (ISD) Scotland. R040X -specialty costs and activity - inpatients in all specialties (excluding long stay), by speciality. 2017/18. Available at: <https://www.isdscotland.org/Health-Topics/Finance/Costs/File-Listings-2018.asp> [last accessed 26 July 2019].
- [20] Orsi G, Di Stefano L, Noah N. Hospital-acquired, laboratory-confirmed bloodstream infection: Increased hospital stay and direct costs. *InfectControl HospEpidemiol* 2002; **23**: 190-7. doi: 10.1086/502034
- [21] Graves N, Weinhold D, Tong E *et al*. Effect of healthcare-acquired infection on length of hospital stay and cost. *Infect Control Hosp Epidemiol* 2007; **28**: 280-92. doi: 10.1086/512642
- [22] Arefian H, Hagel S, Heublein S *et al*. Extra length of stay and costs because of health care-associated infections at a German university hospital. *Am J Infect Control* 2016; **44**: 160-6. doi: 10.1016/j.ajic.2015.09.005
- [23] Leistner R, Hirsemann E, Bloch A, Gastmeier P, Geffers C. Costs and prolonged length of stay of central venous catheter-associated bloodstream infections (CVC BSI): a matched prospective cohort study. *Infection* 2014; **42**: 31-6. doi: 10.1007/s15010-013-0494-z
- [24] Schweizer ML, Cullen JJ, Perencevich EN, Vaughan Sarrazin MS. Costs Associated With Surgical Site Infections in Veterans Affairs Hospitals. *JAMA Surg* 2014; **149**: 575-81. doi: 10.1001/jamasurg.2013.4663
- [25] Vrijens F, Hulstaert F, Van de Sande S, Devriese S, Morales I, Parmentier Y. Hospital-acquired, laboratory-confirmed bloodstream infections: linking national surveillance data to clinical and financial hospital data to estimate increased length of stay and healthcare costs. *JHospInfect* 2010; **75**: 158. doi: <http://dx.doi.org/10.1016/j.jhin.2009.12.006>
- [26] Information Services Division (ISD) Scotland. R040: Specialty Group Costs - Inpatients in all Specialties (exc Long Stay) April 2017 - March 2018. 2018. Available at: <https://www.isdscotland.org/Health-Topics/Finance/Costs/File-Listings-2018.asp> [last accessed 12 April 2019].

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

- [27] NHS National Services Scotland. National Infection Prevention and Control Manual - Appendix 13 - Mandatory - NHSScotland Alert organism/Condition list. 2012. Available at: <http://www.nipcm.hps.scot.nhs.uk/> [last accessed 13 November 2018].
- [28] National Institute for Health and Clinical Excellence. British National Formulary. 2019. Available at: <https://bnf.nice.org.uk/>.
- [29] NHS Digital. Hospital Admitted Patient Care and Adult Critical Care Activity in England 2017-18. 2018. Available at: <https://digital.nhs.uk/data-and-information/publications/statistical/hospital-admitted-patient-care-activity/2017-18> [last accessed 8 December 2019].
- [30] NHS Wales. Patient Episode Database for Wales (PEDW) Data for the period 1st April 2017 to 31st March 2018 NHS Hospital Admitted Patients - Headline Figures. 2018. Available at: http://www.infoandstats.wales.nhs.uk/Documents/869/KeyFacts_WelshProviders_2018.pdf [last accessed 8 December 2019].
- [31] Public Health Scotland - Data and intelligence. Summary inpatient/day case activity by NHS Board of residence, 2009/10 - 2018/19p. 2019. Available at: <https://www.isdscotland.org/Health-Topics/Hospital-Care/Publications/2019-09-10/Inpatient-Day-Case-Activity-Health-Board-of-Residence-Sep19.xlsx> [last accessed 8 December 2019].
- [32] Department of Health - Northern Ireland. Release of Northern Ireland Inpatient, Day Case And Outpatient Hospital Statistics For 2017/18. 2018. Available at: <https://www.health-ni.gov.uk/news/release-northern-ireland-inpatient-day-case-and-outpatient-hospital-statistics-201718> [last accessed 8 December 2019].
- [33] Information Services Division (ISD) Scotland. Average-Length-of-Stay-by-Health-Board-and-Specialty-Sep19. 2019. Available at: <https://www.isdscotland.org/Health-Topics/Hospital-Care/Inpatient-and-Day-Case-Activity/>.
- [34] Information Services Division (ISD) Scotland. R025: Hospital Running Costs by Patient Type - NHS Board Level. 2019. Available at: https://www.isdscotland.org/Health-Topics/Finance/Publications/2019-11-19/Costs_D025_2019.xlsx [last accessed January 2020].
- [35] Sandmann FG, Robotham JV, Deeny SR, Edmunds WJ, Jit M. Estimating the opportunity costs of bed-days. *Health Econ* 2017. doi: 10.1002/hec.3613
- [36] Stewardson AJ, Harbarth S, Graves N. Valuation of hospital bed-days released by infection control programs: a comparison of methods. *Infect Control Hosp Epidemiol* 2014; **35**: 1294-7. doi: 10.1086/678063
- [37] Page K, Barnett AG, Graves N. What is a hospital bed day worth? A contingent valuation study of hospital Chief Executive Officers. *BMC Health Serv Res* 2017; **17**: 137. doi: 10.1186/s12913-017-2079-5
- [38] Jenks PJ, Laurent M, McQuarry S, Watkins R. Clinical and economic burden of surgical site infection (SSI) and predicted financial consequences of elimination of SSI from an English hospital. *JHospInfect* 2014; **86**: 24-33. doi: 10.1016/j.jhin.2013.09.012
- [39] Kaier K, Wolkewitz M, Hehn P, Mutters NT, Heister T. The impact of hospital-acquired infections on the patient-level reimbursement-cost relationship in a DRG-based hospital payment system. *Int J Health Econ Manag* 2019. doi: 10.1007/s10754-019-09267-w
- [40] Information Services Division (ISD) Scotland. Annual Trends in Available Beds by Health Board of Treatment and Hospital. 2019. Available at:

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

- <https://www.isdscotland.org/Health-Topics/Hospital-Care/Beds/> [last accessed 6 March 2020].
- [41] NHS England. Bed Availability and Occupancy Data (Beds Timeseries 2010-11 onwards). 2019. Available at: <https://www.england.nhs.uk/statistics/statistical-work-areas/bed-availability-and-occupancy/bed-data-overnight/> [last accessed 6 March 2020].
- [42] Macedo-Viñas M, De Angelis G, Rohner P *et al.* Burden of meticillin-resistant *Staphylococcus aureus* infections at a Swiss University hospital: excess length of stay and costs. *J Hosp Infect* 2013; **84**: 132-7. doi: 10.1016/j.jhin.2013.02.015
- [43] Le Monnier A, Duburcq A, Zahar JR *et al.* Hospital cost of *Clostridium difficile* infection including the contribution of recurrences in French acute-care hospitals. *J Hosp Infect* 2015; **91**: 117-22. doi: 10.1016/j.jhin.2015.06.017
- [44] Baier C, Linke L, Eder M *et al.* Incidence, risk factors and healthcare costs of central line-associated nosocomial bloodstream infections in hematologic and oncologic patients. *PLOS ONE* 2020; **15**: e0227772. doi: 10.1371/journal.pone.0227772
- [45] Guest JF, Keating T, Gould D, Wigglesworth N. Modelling the annual NHS costs and outcomes attributable to healthcare-associated infections in England. *BMJ Open* 2020; **10**: e033367. doi: 10.1136/bmjopen-2019-033367
- [46] Heister T, Wolkewitz M, Hehn P *et al.* Costs of hospital-acquired *Clostridium difficile* infections: an analysis on the effect of time-dependent exposures using routine and surveillance data. *Cost Eff Resour Alloc* 2019; **17**: 16. doi: 10.1186/s12962-019-0184-5
- [47] Naylor NR, Atun R, Zhu N *et al.* Estimating the burden of antimicrobial resistance: a systematic literature review. *Antimicrob Resist Infect Control* 2018; **7**: 58. doi: 10.1186/s13756-018-0336-y
- [48] Schreiber PW, Sax H, Wolfensberger A, Clack L, Kuster SP, Swissnoso. The preventable proportion of healthcare-associated infections 2005-2016: Systematic review and meta-analysis. *Infect Control Hosp Epidemiol* 2018; **39**: 1277-95. doi: 10.1017/ice.2018.183
- [49] Caro JJ, Briggs AH, Siebert U, Kuntz KM. Modeling Good Research Practices—Overview: A Report of the ISPOR-SMDM Modeling Good Research Practices Task Force-1. *Value Health* 2012; **15**: 796-803. doi: <http://doi.org/10.1016/j.jval.2012.06.012>
- [50] Stewart S, Robertson C, Kennedy S *et al.* ECONI: Personalised infection prevention and control – identifying patients at risk of healthcare associated infection. *J Hosp Infect*; **In this issue**:

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table 1 Bed-day unit costs

Unit cost	Mean	SD ¹	Distribution	Source
9) Bed-day total cost	£799.17	£535.37	Log-normal	ISD[26]
10) Bed-day direct cost	£519.38	£344.27	Log-normal	ISD[26]
Notes 1 Standard deviation				

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table II: Incidence, average excess LOS and bed-days lost by HAI type in Scotland

HAI	Incidence Rates per 100,000 AOBs (95% CI) ¹	Estimated annual number of HAI cases (95% CI) ¹	Average Excess LOS ² per HAI (95% CI) ¹	Total annual bed-days lost to HAI ³ (N)	(95% CI) ¹
BSI	35.0 (31.3-39.5)	1,389 (1,245-1,570)	11.4 (5.8-17.0)	15,830	(7,550-23,950)
GI	31.6 (28.2-36.3)	1,256 (1,122-1,440)	6.0 (-0.7-12.7)	7,540	(0 -16,100)
LRI	26.2 (3.6-30.7)	1,041 (937-1,218)	7.3 (1.8-12.7)	7,600	(1,300 -13,540)
PN	15.9 (13.7 -19.2)	630 (544-764)	16.3 (7.5-25.5)	10,270	(4,170-16,380)
SSI	25.8 (22.8-30.5)	1,023 (904-1,210)	9.8 (4.5-15.0)	10,030	(4,190-15,900)
UTI	41.0 (36.6- 46.2)	1,628 (1,454-1,836)	-1.0 (-4.3-2.3)	0	(0-4,180)
Other	12.0 (9.5-15.8)	475 (378-628)	14.0 (-3.9-31.8)	6,650	(0-16,360)
All HAI	187.2 (176.8-197.6)	7,437 (7,021-7,849)	7.8 (5.7-9.9)	58,010	(41,730-74,840)

Notes: (BSI) Blood Stream Infection; (GI) Gastrointestinal Infection; (LRI) Lower Respiratory tract Infection; (PN) Pneumonia; (SSI) Surgical Site Infection; (UTI) Urinary Tract Infection; Other are small number infections – includes (SST) Skin Soft Tissue; (BJ) Bone and Joint; (CV) Cardiovascular; (EENT) Eye, Ear, Nose and Throat; and (SI) systemic infections.
1 95% confidence interval.
2 Excess LOS is expressed in additional days per case.
3 NHS Scotland acute hospitals.

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table III: Cost per case for each HAI type and annual cost of HAI in NHS Scotland

HAI	Cost per case for each HAI type and overall				Annual cost in NHS Scotland (Million)			
	Total cost per case		Direct cost per case		Total cost		Direct cost	
	Mean	(95% CI) ¹	Mean	(95% CI) ¹	Mean	(95% CI) ¹	Mean	(95% CI) ¹
BSI	£9,109	(£3,511-£28,210)	£5,917	(£2,552-£15,438)	£12.65	(£4.82-£38.96)	£8.22	(£3.45-£21.94)
GI	£4,794	(£445-£19,835)	£3,114	(£192-£10,401)	£6.02	(£0.52-£24.83)	£3.91	(£0.24-£14.11)
LRI	£5,833	(£1,729-£20,019)	£3,789	(£1,234-£11,684)	£6.07	(£1.66-£21.25)	£3.94	(£1.17-£11.84)
PN	£13,024	(£4,808-£45,061)	£8,460	(£3,432-£23,548)	£8.20	(£2.99-£25.88)	£5.33	(£2.01-£14.44)
SSI	£7,830	(£2,987-£24,993)	£5,086	(£2,095-£14,433)	£8.01	(£2.95-£27.02)	£5.20	(£1.96-£13.45)
UTI	£0	(£0-£2,109)	£0	(£0-£1,304)	£0	(£0-£3.63)	£0	(£0-£2.21)
Other	£11,186	(£0-£45,319)	£7,266	(£0-£26,523)	£5.31	(£0.05-£24.45)	£3.45	(£0-£12.77)
All HAI	£6,232	(£2,733-£18,181)	£4,048	(£1,927-£9,591)	£46.35	(£19.43-£128.81)	£30.11	(£14.12-£74.46)

Notes: (BSI) Blood Stream Infection; (GI) Gastrointestinal Infection; (LRI) Lower Respiratory tract Infection; (PN) Pneumonia; (SSI) Surgical Site Infection; (UTI) Urinary Tract Infection; Other are small number infections – includes (SST) Skin Soft Tissue; (BJ) Bone and Joint; (CV) Cardiovascular; (EENT) Eye, Ear, Nose and Throat; and (SI) systemic infections.
¹ 95% confidence interval.

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table IV Annual total inpatient cost of HAI in the UK

Annual total cost in the UK (£Million per Year) ²		Annual direct cost in the UK (£Million per Year) ²	
Mean	(95% CI) ¹	Mean	(95% CI) ¹
£774	(£328-£2,192)	£503	(£236-£1,217)
Notes: 1 (95% confidence interval). 2 Annual UK total and direct costs are based on ECONI HAI incidence: 0.76% (95% CI: 0.72%-0.81%).			

ECONI: Bed-days and costs associated with the inpatient burden of healthcare associated infection in the UK

Table V: Annual IPC, laboratory materials and antibiotics used in the treatment of HAI

Cash Treatment Costs in Scotland						
IPC and Laboratory costs					Antibiotic costs ¹	
HAI	Mean IPC cost per HAI case	Mean Laboratory test cost per HAI case (£)	Total IPC and Lab cost per HAI case (£)	Annual cost Scotland (£)	Cost of antibiotics per HAI case (£)	Annual cost in Scotland (£)
BSI	£15.76	£23.13	£38.89	£54,018	£91	£126,399
GI	£67.10	£65.52	£132.62	£166,571	£43	£54,008
LRI	£22.53	£39.10	£61.63	£64,157	£125	£130,125
PN	£24.01	£36.41	£60.42	£38,065	£280	£176,400
SSI	£12.62	£27.10	£39.72	£40,634	£454	£464,442
UTI	£14.96	£19.03	£33.99	£55,336	£7	£11,396
Other	£8.44		£8.44	£4,009	£4	£1,900
All HAI				£422,790	£91	£676,767

Notes
1: Antibiotic costs were calculated from the case-control sample and not directly comparable with other results presented in this study.