

Improving older adults' vaccination uptake: Are existing measures of vaccine hesitancy valid and reliable for older people?

Journal of Health Psychology
1–12

© The Author(s) 2022



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/13591053221089104

journals.sagepub.com/home/hpq



Nicola Cogan¹ , Allyson J Gallant²,
Louise A Brown Nicholls¹, Susan Rasmussen¹, David Young¹
and Lynn Williams¹

Abstract

We sought to establish whether two recently developed measures, the 5C scale and the Vaccination Attitudes Examination (VAX) were reliable and valid for use with older adults. A total of 372 UK-dwelling participants (65–92 years, $M = 70.5$ years, $SD = 4.6$) completed a cross-sectional survey measuring health and socio-demographic characteristics in relation to vaccine uptake for influenza, pneumococcal and shingles. The 5C and VAX scales were administered to test their reliability, validity and dimensionality. Both scales showed good internal reliability and convergent, discriminant and concurrent validity. Their factor structures were also confirmed, supporting their use with older adult populations.

Keywords

ageing, influenza, pneumococcal, psychometrics, shingles, vaccine hesitancy

Introduction

Outbreaks of vaccine-preventable diseases (VPDs) are a growing international concern particularly in the context of the COVID-19 pandemic (Suryadevara, 2021; Gallant et al., 2021). As a result of these outbreaks there is an urgent need to focus on the development and evaluation of screening tools to identify those who may be hesitant about vaccine(s) and to inform interventions to increase vaccine uptake. Vaccine hesitancy (VH) has contributed to outbreaks of VPDs across a range of socio-cultural contexts and populations (Murphy et al., 2021; Sallam, 2021; Wiysonge et al., 2021). Although there is a lack of consensus in terms of how to

define VH, it can be understood as the mental state of holding back, in doubt or indecision regarding vaccination (Bedford et al., 2018). VH includes a broad spectrum of attitudes, beliefs and behaviours including vaccine refusal and intentional vaccine delay (Ryan and Malinga, 2021). While there has been a

¹University of Strathclyde, UK

²Dalhousie University, Canada

Corresponding author:

Nicola Cogan, School of Psychological Sciences and Health, University of Strathclyde, 40 George Street, Glasgow G1 1QE, UK.

Email: nicola.cogan@strath.ac.uk

growing body of research exploring VH among children, parents and diverse adult populations and occupational groups (Alfieri et al., 2021; Noyman-Veksler et al., 2021; Quinn et al., 2019; Saied et al., 2021; Santibanez et al., 2020; Shahar et al., 2017), there is need for further work incorporating the expanding demographic of older adults (aged ≥ 65 years), given that they are recognised as an 'at risk' group for VPDs (Bhagianadh and Arora, 2022; Nicholls et al., 2021; Privor-Dumm et al., 2021; Talbird et al., 2020).

Due to decreased immunity and greater likelihood of existing chronic health conditions, older adults are more susceptible to infectious diseases and have altered immune response to vaccinations (Coll et al., 2020; Russell et al., 2018). Vaccination can support healthy ageing and efforts to increase uptake among older adults seek to reduce morbidity, mortality and loss of quality of life associated with VPDs (Ozawa et al., 2016). Vaccination uptake varies by vaccine (Klett-Tammen et al., 2016); despite the availability of effective vaccines, many countries fail to reach recommended coverage levels (Doherty et al., 2018). Within the UK, older adults (aged ≥ 65 years) are freely offered an annual influenza vaccine, a single-dose pneumococcal vaccine and a single-dose shingles (aged 70–79 years) vaccine. However, uptake has been found to fall below the World Health Organisation's (WHO's) target of 75% (Dios-Guerra et al., 2017; Sheikh et al., 2018). For influenza vaccination, between 2004–05 and 2019–20, uptake has fluctuated between 71% and 75% among older adults (Health Protection Scotland, 2021b). Pneumococcal vaccination uptake is typically around 51%–69% (Briggs et al., 2019; Frank et al., 2020), and shingles uptake rates can be lower than 50% (Health Protection Scotland, 2021a).

Given these uptake rates, it is important to understand the reasons for low uptake and/or VH. A recent population-level cohort study of UK adults (aged ≥ 65 years) found that uptake for influenza, shingles and pneumococcal vaccination are patterned by ethnicity, deprivation, household size and comorbidities (Tan et al.,

2021). In addition, psychosocial factors are increasingly being recognised as providing important insights into the determinants of vaccination behaviour (Betsch et al., 2018, 2020; Schmid et al., 2017). Measuring vaccination-related psychosocial factors is necessary to identify target populations, determine potential reasons for under-vaccination and inform the design of tailored and cost-effective interventions (Butler and MacDonald, 2015; Eitze et al., 2021). Recently, two self-report measures, the 5C scale (Betsch et al., 2018) and the Vaccination Attitudes Examination (VAX) Scale (Martin and Petrie, 2017), have been developed to help identify reasons why people do or do not vaccinate. The 5C scale is a novel measure of five psychological antecedents, or precursors, of vaccination behaviour: confidence, complacency, constraints, calculation and collective responsibility. The VAX scale assesses attitudes that may underlie VH across four domains: mistrust of vaccine benefit, worries about unforeseen future effects, concerns about commercial profiteering, and preference for natural immunity. These measures are useful for predicting vaccination behaviour; however, to date, their reliability and validity have yet to be confirmed in older adults aged ≥ 65 years. While the 5C scale has been tested once with slightly older adults, this sample was middle aged ($M=48$ years). The VAX scale has been tested with young adults only (typically those in their 20s and 30s) and, therefore, it is not known if it can distinguish between vaccinating and non-vaccinating older adults.

Presently, we aimed to test the reliability, validity and dimensionality of the 5C (Betsch et al., 2018) and VAX (Martin and Petrie, 2017) scales with UK-dwelling older adults (aged ≥ 65 years) in relation to VH for influenza, pneumococcal and shingles. We sought to test the psychometric properties of these scales and use confirmatory factor analysis to elucidate their underlying factor structures, thereby supporting their generalisability with an older adult population. Further, we aimed to establish their internal consistency and discriminant and convergent validity. Building upon earlier work

assessing independent predictors of older adults' vaccination uptake (Nicholls et al., 2021), the concurrent validity of the 5C and VAX was also examined.

Methods

Participants

Participants were 372 older adults aged 65–92 years ($M=70.5$ years, $SD=4.6$). Participants were living in the UK, generally in good health, not diagnosed with a neurological condition and living independently in the community (see Table 1 for socio-demographic characteristics).

Procedure

An online, cross-sectional survey was administered using Qualtrics. Data collection took place between 8th February and 17th March 2020, prior to the first COVID-19 lockdown in the UK. Participants were recruited through university participation panels and social media (e.g. Twitter and Facebook).

Measures

The online survey included the 5C (Betsch et al., 2018) and the VAX (Martin and Petrie, 2017). We included other measures related to VH; the Beliefs About Medicines Questionnaire (BMQ; Horne et al., 1999), the Perceived Sensitivity to Medicines scale (PSM; Horne et al., 2013) and the Medical Mistrust Index (MMI; LaVeist et al., 2009). We also included measures that we considered theoretically not to be highly related to VH; the Perceived Stress Scale (PSS; Cohen et al., 1994) and the Interpersonal Support Evaluation List (ISEL-12; Cohen et al., 1985) (see Supplemental Table for description of measures). Participants were asked their age, gender, marital status, ethnicity, self-reported overall health, education and socio-economic status. Participants were also asked whether they had received the influenza vaccination in the past 12 months and if they had ever received the pneumococcal vaccination. Those aged 70

Table 1. Participants' socio-demographic data.

Variables	<i>n</i> (372 total)
Age	$M=70.5$ ($SD=4.6$)
Gender	
Female	184 (49.7%)
Male	184 (49.7%)
Prefer not to say	2 (0.5%)
Marital status	
Married	248 (66.8%)
Widowed	47 (12.7%)
Separated/Divorced	35 (9.4%)
Co-habiting	21 (5.7%)
Single	20 (5.4%)
Ethnicity	
White-British	360 (97%)
White-Other	9 (2.4%)
Asian	1 (0.3%)
Mixed/Multiple	1 (0.3%)
Self-rated overall health	
Very good	125 (33.6%)
Good	201 (54%)
Fair	41 (11%)
Poor	5 (1.3%)
Education	
High school	65 (17.5%)
College	114 (30.6%)
University	109 (29.3%)
Postgraduate	84 (22.6%)
Deprivation quintile	
1 (most deprived)	57 (16.3%)
2	83 (23.8%)
3	69 (19.8%)
4	61 (17.5%)
5 (least deprived)	79 (22.6%)

NB: % calculations exclude missing data.

and older were asked about previous uptake of the shingles vaccination. Self-reported vaccine status has been found to be a sensitive and fairly specific indicator of actual vaccine status (Irving et al., 2009).

Data analyses

Analyses were performed using SPSS version 23. First, internal consistency using MIIC rather than Cronbach's α was used due to the low number of items in each subscale. Next, to

verify the structural validity of the 5C and VAX scales, confirmatory factor analysis (CFA) was performed. The following indices of model fit were considered: comparative fit index (CFI), the Tucker Lewis Index (TLI), root mean square error of approximation (RMSEA) and standardised root mean square error of approximation and MIIC. CFI and TLI values between 0.90 and 0.95 and RMSEA values between 0.05 and 0.08 are indicative of acceptable model fit (Chen, 2007; Hu and Bentler, 1999; Kline, 2005). Convergent validity was examined using the correlations between the 5C and VAX with the BMQ, PSM and MMI. Discriminant validity was assessed by examining the correlations between the 5C and VAX with the PSS and IESL-12. Finally, concurrent validity was assessed using logistic regressions (correct classification rates) within each scale, to determine if they successfully predicted vaccination behaviour (dependent variable: vaccinated yes/no for influenza, pneumococcal and shingles). All tests were two-tailed, and $p < 0.05$ was considered statistically significant. The current study includes the complete raw data set collected in the study including the syntax file and log file for analysis. All of the data files will be uploaded to the Figshare repository. The quantitative data that support the findings of this study are currently openly available in Open Science Framework (<https://doi.org/10.17605/OSF.IO/JYPN9>).

Results

Internal reliability

For the 5C, good internal consistency was found across all subscales (with MIIC > 0.15): confidence, MIIC=0.74; complacency, MIIC=0.30; constraints, MIIC=0.47, calculation MIIC=0.58 and collective responsibility MIIC=0.32. For the VAX, a total mean score MIIC=0.45 was obtained with subscales: *mistrust of vaccine benefits*, MIIC=0.74; *worries over unforeseen future effects*, MIIC=0.49; *concerns about commercial profiteering*, MIIC=0.73; and *preference for natural*

immunity, MIIC=0.67; all demonstrated good internal consistency. The 5C and VAX were found to be reliable measures for use with an older adult population.

Factor structure of 5C and VAX

The factor structure of the 5C and VAX was assessed using CFA. CFA was run on all 5C subscales and the 12 VAX items grouped into four subscales (three items per subscale). For the tested models, RMSEA values were reasonable (values near to 0.08). The CFI and TLI were above the acceptable value of 0.90 (5C: CFI=0.959, TLI=0.946; VAX: CFI=0.978, TLI=0.970) suggesting good fit and verifies the structural validity of the scales when tested with an older adult population. The 5C and VAX were found to have sound factor structures, when tested with an older adult population.

Convergent and discriminant validity

Table 2 shows the correlations measuring associations between the mean 5C subscale scores, VAX total score and MMI, BMQ and PSM total means for the sample. Mean 5C subscale scores and VAX total scores correlated positively with MMI, BMQ and PSM, showing good convergent validity. Discriminant validity was assessed by examining the correlations between the mean 5C subscale scores and VAX total score with PSS and IESL-12. As expected, these correlations were found to be weaker than those observed with the BMQ, PSM and MMI. Both the 5C and VAX showed good convergent and discriminant validity, supporting their use with older adult populations.

Concurrent validity

Concurrent validity was assessed using logistic regression to determine if the 5C and VAX scales successfully predicted vaccination behaviour (vaccinated yes/no) for influenza, pneumococcal and shingles. Multivariate logistic models were constructed to determine the

Table 2. Correlation matrix of associations between mean VAX total and 5C scores and relevant questionnaires to test convergent and discriminant validity.

Measure	1	2	3	4	5	6	7	8	9	10	11
1. 5C Confidence	-	-0.338***	-0.250***	-0.095	0.405***	-0.584***	-0.078	0.073	-0.204***	-0.349***	-0.260***
2. 5C Complacency			0.341***	0.156**	-0.443***	0.583***	0.071	-0.083	0.209***	0.335***	0.195***
3. 5C Constraints				0.091**	-0.265***	0.374***	0.265***	-0.160**	0.232***	0.233***	0.220***
4. 5C Calculation					-0.069	0.216***	0.030	0.071	0.096	0.140**	0.112*
5. 5C Collective Responsibility						-0.597***	-0.006	0.062	-0.161**	-0.277***	-0.162**
6. VAX total							0.104*	-0.128	0.389***	0.593***	0.298***
7. PSS total								-0.409***	0.142**	0.135*	0.171***
8. IESL-12 total									-0.126*	-0.157**	-0.075
9. MMI total										0.438***	0.202***
10. BMQ total											0.367***
11. PSM total											-

BMQ: Beliefs About Medicines Questionnaire-General; IESL-12: Interpersonal Support Evaluation List; MMI: Medical Mistrust Index; PSM: Perceived Sensitivity to Medicines Scale; PSS: Perceived Stress Scale; VAX: Vaccination Attitudes Examination Scale.

*** $p < 0.001$. ** $p < 0.01$. * $p < 0.05$.

independent predictors of lack of vaccine uptake for each vaccine. Results (see Table 3) showed that both the 5C and VAX scales were able to correctly classify vaccinators and non-vaccinators across each of the vaccines, thus showing good concurrent validity.

Discussion

To our knowledge, our study is the first to establish the reliability and validity of the 5C (Betsch et al., 2018) and VAX (Martin and Petrie, 2017) scales, two psychosocial measures of vaccine uptake, in an older adult population. Our study revealed satisfactory psychometric properties for both measures as the results were indicative of statistically sound factor models. The 5C and VAX were shown to have good internal consistency and good levels of convergent and discriminant validity. As reported in earlier work, psychosocial factors independently predicted older adults' hesitancy towards the influenza, pneumococcal and shingles vaccines (Nicholls et al., 2021). Presently, these findings additionally show that the two scales have good concurrent validity.

The 5C and VAX contribute to metrics which aim to assess VH (Betsch et al., 2018; Huza, 2020; Martin and Petrie, 2017; Wood et al., 2019) through demonstrating that both these measures are reliable and valid for use with an older adult (aged 65+) population. Such measures can be used to inform targeted public health action to increase vaccine uptake with older adults, using appropriate strategies, policies and interventions to reduce VH, and during mass screening and vaccination campaigns (Frank et al., 2020; Jarrett et al., 2015; Nicholls et al., 2021). Given the pressures on healthcare systems during the COVID-19 pandemic, having sound measures of VH with older adults, who may be more vulnerable to adverse outcomes, has come into strong focus (Roller-Wirnsberger et al., 2021). This is pertinent given that recent longitudinal work suggests that the pandemic has led older adults to re-evaluate the importance of vaccination and may have increased their willingness

to protect others; however, concerns about vaccine side effects were found to have significantly increased during the pandemic (Gallant et al., 2021). These findings are also consistent with previous research relating to COVID-19 vaccines among diverse populations, where participants have raised concerns about vaccine side effects (Domnich et al., 2020; Pogue et al., 2020; Williams et al., 2020). Understanding and measuring ongoing structural and psychosocial barriers to vaccination is essential (Baraniuk, 2021).

Future studies are recommended to determine the relationship between psychosocial and other factors associated with older adults' vaccine uptake using the 5C and VAX. Given that participants in the current study were high functioning and were living independently at home with minimal assistance, the findings may not be generalisable to older adults with lower functional abilities, for example, those who are experiencing cognitive impairment or limitations in everyday functioning (Nicholls et al., 2021). Further work could therefore explore the psychometric properties and feasibility of the 5C and VAX in more diverse populations of older adults (e.g. ethnic minorities, LGBTQ+, varied functional levels) and socio-cultural contexts (e.g. low-income countries); This is essential given increasing evidence that minority and/or vulnerable populations face challenges due to inequitable access, discrimination and systemic suppression which impacts on vaccine uptake (Corbie-Smith, 2021; Garg et al., 2021). Longitudinal studies are also required to establish their ability to predict outcomes such as objective measurements of vaccination uptake relative to the total number of recommended dosages. Furthermore, it is possible that participants may report less VH face-to-face than through an anonymous online survey, which may encourage self-disclosure on sensitive items (Hollier et al., 2017). It would be useful for future work to evaluate possible differences in VH using the 5C and VAX with online surveys, telephone surveys and face-to-face assessments. Given that older adults are at increased risk of disease morbidity, including VPDs, targeting

Table 3. Multivariate logistic regression analyses within the 5C and VAX Scales for predicting older adults' non-uptake of influenza, pneumococcal, and shingles vaccines.

Vaccination	Concurrent validity	Independent predictors	OR (95% CI)	Correct classification rate (%)
Influenza	5C	Complacency ($p < 0.001$ ***)	1.834 (1.359, 2.474)	89.6
		Calculation ($p = 0.003$ ***)	1.568 (1.161, 2.118)	
Pneumococcal	VAX	Collective responsibility ($p < 0.001$ ***)	0.375 (0.276, 0.509)	91.0
		Mistrust ($p < 0.001$ ***)	2.562 (1.567, 4.188)	
		Worries ($p = 0.011$ *)	1.954 (1.163, 3.284)	
		Natural immunity ($p < 0.001$ ***)	2.583 (1.571, 4.247)	
		Complacency ($p = 0.007$ ***)	1.350 (1.088, 1.675)	
Shingles	5C	Collective responsibility ($p = 0.001$ ***)	0.685 (0.552, 0.849)	68.6
		Mistrust ($p = 0.004$ ***)	1.624 (1.164, 2.265)	
		Concerns ($p = 0.002$ ***)	1.613 (1.190, 2.187)	
		Complacency ($p = 0.033$ *)	1.385 (1.026, 1.871)	
		Collective responsibility ($p = 0.004$ ***)	0.623 (0.453, 0.856)	
Shingles	VAX	Worries ($p = 0.039$ *)	1.656 (1.026, 2.673)	67.0
		Concerns ($p = 0.002$ ***)	1.985 (1.281, 3.076)	

OR: Odds Ratio.

*** $p < 0.001$. ** $p < 0.01$. * $p < 0.05$.

research to inform intervention programmes to increase uptake, using psychometrically validated measures of VH, is of paramount importance. This is particularly pertinent given the global challenges associated with the dynamic, complex and changing nature of COVID-19, highlighting the need for ongoing research on VH (Holeva et al., 2022; Karafillakis et al., 2022; Reno et al., 2021). Measuring and monitoring VH among older adults and using this information to inform policy (Sprengholz et al., 2021) and intervention developments to increase vaccine uptake is essential (Higgins et al., 2021; Motta et al., 2021). Developing a better understanding of older adults' perceived risks versus benefits of vaccines not only involves measuring attitudes and beliefs concerning vaccines but also engaging with health practitioners, media, governmental bodies and community organisations to increase awareness of the burden of VPDs (Van Buynder and Woodward, 2019) and the risks associated with VH (Dube et al., 2013). Indeed, a recent longitudinal randomised controlled study conducted with older adults, who were especially vulnerable to primary and secondary diseases, recommended that interventions aiming to reduce VH might be more effective in specific contexts and locations such as doctors' clinics (Eitze et al., 2021). Using the 5C and VAX in conjunction with other measures of VH such as the Multidimensional Vaccine Hesitancy Scale (Howard, 2021) may be particularly useful in capturing the changeable nature of VH and help inform the development of adaptive interventions. Given the heterogeneity in VH and the diversity of situations and contexts in which it can arise; screening tools and interventions aiming to address VH may benefit from being both context and problem-specific (Ryan and Malinga, 2021), as well as seeking to understand population-specific challenges (Kristensen et al., 2016; Tan et al., 2022).

Conclusion

Globally, in developed and developing countries alike, a demographic shift towards an ageing population is underway. Many VPDs inflict

a disproportionate burden of disease in older adults but may be prevented or attenuated by vaccination. Vaccine uptake by this population is generally low and must be substantially improved if the potential of vaccines to reduce the morbidity, mortality, loss of quality of life and health and social care costs caused by VPDs is to be realised. Importantly, the available evidence indicates that VH in older adults can be reduced. Having reliable and valid measures of VH is essential, however, in order to better understand and develop tailored and targeted interventions to address VH among this population. In our work, the 5C and VAX scales were found to be reliable and valid measures of VH with a UK-dwelling, older adult (65+) population. Future research could use these scales to evaluate the impact of psychological antecedents of vaccine uptake, and how concerns about vaccination may be challenged and reversed in older adults. Improved measurement and targeted education around VPDs, disease risk and vaccine benefits is required to increase vaccine coverage. It is hoped that these findings will influence future research and intervention development with older adults aimed at achieving this. The 5C and VAX scales are therefore useful to help understand the health and vaccination attitudes that promote and deter older adults' vaccination behaviours. Behavioural and psychosocial research can support the design and measurement of effective vaccination interventions directed towards older adults, aiming to reduce VH and increase uptake. A multi-dimensional approach measuring VH using the 5C and VAX, along with ensuring access to vaccines, will be important. The lessons learned from the COVID-19 vaccination drive will also aid in the successful implementation of subsequent vaccinations and in efforts to reach those older adults who may be most resistant to vaccines. Further work is needed in order to design evidence-informed responses to VH appropriate to the setting, context and hesitant subgroup. Given the ageing population and corresponding increases in healthcare resource utilisation required to treat VPDs, we

recommend particular attention be directed towards research focussing on 'at risk' older adult populations.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Data sharing

The quantitative data that support the findings of this study are openly available in Open Science Framework (<https://doi.org/10>).

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the Chief Scientist Office, Scottish Government [grant number CGA/19/52].


Ethical approval

Ethical approval (34/26/11/2019/Staff) was granted by the School of Psychological Sciences and Health Ethics Committee (SEC19/20: Williams et al., 2020). Approved on 8th January 2020. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent

Informed consent was obtained from all individual participants included in the study.

ORCID iD

Nicola Cogan  <https://orcid.org/0000-0003-0861-5133>

Supplemental material

Supplemental material for this article is available online.

References

- Alfieri NL, Kusma JD, Heard-Garris N, et al. (2021) Parental COVID-19 vaccine hesitancy for children: Vulnerability in an urban hotspot. *BMC Public Health* 21(1): 1–9.
- Baraniuk C (2021) Covid-19: How the UK vaccine rollout delivered success, so far. *BMJ* 372: n421.
- Bedford H, Attwell K, Danchin M, et al. (2018) Vaccine hesitancy, refusal and access barriers: The need for clarity in terminology. *Vaccine* 36(44): 6556–6558.
- Betsch C, Bach Habersaat K, Deshevoi S, et al. (2020) Sample study protocol for adapting and translating the 5C scale to assess the psychological antecedents of vaccination. *BMJ Open* 10(3): e034869.
- Betsch C, Schmid P, Heinemeier D, et al. (2018) Beyond confidence: Development of a measure assessing the 5C psychological antecedents of vaccination. *PLoS One* 13(12): e0208601.
- Bhagianadh D and Arora K (2022) COVID-19 vaccine hesitancy among community-dwelling older adults: The role of Information Sources. *Journal of Applied Gerontology* 41(1): 4–11.
- Briggs L, Fronek P, Quinn V, et al. (2019) Perceptions of influenza and pneumococcal vaccine uptake by older persons in Australia. *Vaccine* 37(32): 4454–4459.
- Butler R and MacDonald NE (2015) Diagnosing the determinants of vaccine hesitancy in specific subgroups: The Guide to Tailoring Immunization Programmes (TIP). *Vaccine* 33(34): 4176–4179.
- Chen FF (2007) Sensitivity of goodness of fit indexes to lack of measurement invariance. *Structural Equation Modeling: A Multidisciplinary Journal* 14(3): 464–504.
- Cohen S, Kamarck T and Mermelstein R (1994) Perceived stress scale. *Measuring Stress: A Guide for Health and Social Scientists* 10: 1–2.
- Cohen S, Mermelstein R, Kamarck T, et al. (1985) Measuring the functional components of social support. In: Sarason IG and Sarason BR (eds) *Social Support: Theory, Research and Applications*. Dordrecht: Springer Netherlands, pp.73–94.
- Coll PP, Costello VW, Kuchel GA, et al. (2020) The prevention of infections in older adults: Vaccination. *Journal of the American Geriatrics Society* 68(1): 207–214.
- Corbie-Smith G (2021) Vaccine hesitancy is a scapegoat for structural racism. *JAMA Health Forum* 2(3): e210434.
- Dios-Guerra C, Carmona-Torres JM, López-Soto PJ, et al. (2017) Prevalence and factors associated with influenza vaccination of persons over 65 years old in Spain (2009–2014). *Vaccine* 35(51): 7095–7100.

- Doherty TM, Connolly MP, Del Giudice G, et al. (2018) Vaccination programs for older adults in an era of demographic change. *European Geriatric Medicine* 9(3): 289–300.
- Domnich A, Cambiaggi M, Vasco A, et al. (2020) Attitudes and beliefs on influenza vaccination during the COVID-19 pandemic: Results from a Representative Italian Survey. *Vaccines* 8(4): 711.
- Dubé, E., Laberge, C., Guay, M., Bramadat, P., Roy, R., & Bettinger, J. A. (2013). Vaccine hesitancy: An overview. *Human Vaccines & Immunotherapeutics* 9(8): 1763–1773.
- Eitze S, Heinemeier D, Schmid-Küpke NK, et al.; Vaccination60+ Study Group (2021) Decreasing vaccine hesitancy with extended health knowledge: Evidence from a longitudinal randomized controlled trial. *Health Psychology* 40(2): 77–88.
- Frank O, De Oliveira Bernardo C, González-Chica DA, et al. (2020) Pneumococcal vaccination uptake among patients aged 65 years or over in Australian general practice. *Human Vaccines & Immunotherapeutics* 16(4): 965–971.
- Gallant AJ, Nicholls LAB, Rasmussen LS, et al. (2021) Changes in attitudes to vaccination as a result of the COVID-19 pandemic: A longitudinal study of older adults in the UK. *PLoS One* 16(12): e0261844.
- Garg I, Hanif H, Javed N, et al. (2021) COVID-19 vaccine hesitancy in the LGBTQ+ population: A systematic review. *Infectious Disease Reports* 13(4): 872–887.
- Health Protection Scotland (2021a) HPS Website—Shingles. Available at: <https://www.hps.scot.nhs.uk/a-to-z-of-topics/shingles/> (accessed October 2021).
- Health Protection Scotland (2021b) Vaccine-Update-dashboard. TableauSoftware. Available at: https://public.tableau.com/views/Vaccine-Update-dashboard/Story2?:embed=y&:showVizHome=no&:host_url=https%3A%2F%2Fpublic.tableau.com%2F&:tabs=no&:toolbar=yes&:animate_transition=yes&:display_static_image=no&:display_spinner=no&:display_overlay=yes&:display_count=yes&publish=yes&:loadOrderID=0 (accessed October 2021).
- Higgins ST, Klemperer EM and Coleman SRM (2021) Looking to the empirical literature on the potential for financial incentives to enhance adherence with COVID-19 vaccination. *Preventive Medicine* 145: 106421.
- Holeva V, Parlapani E, Nikopoulou VA, et al. (2022) COVID-19 vaccine hesitancy in a sample of Greek adults. *Psychology Health & Medicine* 27: 113–119.
- Hollier LP, Pettigrew S, Slevin T, et al. (2017) Comparing online and telephone survey results in the context of a skin cancer prevention campaign evaluation. *Journal of Public Health* 39(1): 193–201.
- Horne R, Faasse K, Cooper V, et al. (2013) The perceived sensitivity to medicines (PSM) scale: An evaluation of validity and reliability. *British Journal of Health Psychology* 18(1): 18–30.
- Horne R, Weinman J and Hankins M (1999) The beliefs about Medicines Questionnaire: The development and evaluation of a new method for assessing the cognitive representation of medication. *Psychology and Health* 14(1): 1–24.
- Howard MC (2021) A more comprehensive measure of vaccine hesitancy: Creation of the Multidimensional Vaccine Hesitancy Scale (MVHS). *Journal of Health Psychology*. Epub ahead of print 2 September 2021. DOI: 10.1177/13591053211042062.
- Hu L and Bentler PM (1999) Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal* 6(1): 1–55.
- Huza G (2020) The psychometric properties of a Romanian version of the Vaccination Attitudes Examination (VAX) scale. *International Journal of HIV/AIDS Prevention, Education and Behavioural Science* 6(1): 25.
- Irving SA, Donahue JG, Shay DK, et al. (2009) Evaluation of self-reported and registry-based influenza vaccination status in a Wisconsin cohort. *Vaccine* 27(47): 6546–6549.
- Jarrett C, Wilson R, O’Leary M, et al.; SAGE Working Group on Vaccine Hesitancy (2015) Strategies for addressing vaccine hesitancy – A systematic review. *Vaccine* 33(34): 4180–4190.
- Karafillakis E, Van Damme P, Hendrickx G, et al. (2022) COVID-19 in Europe: New challenges for addressing vaccine hesitancy. *Lancet* 399: 699–701.
- Klett-Tammen CJ, Krause G, Seefeld L, et al. (2016) Determinants of tetanus, pneumococcal and influenza vaccination in the elderly: A representative cross-sectional study on knowledge,

- attitude and practice (KAP). *BMC Public Health* 16(1): 121–129.
- Kline RB (2005) *Principles and Practice of Structural Equation Modeling*, 2nd edn. New York, NY: The Guilford Press.
- Kristensen M, van Lier A, Eilers R, et al. (2016) Burden of four vaccine preventable diseases in older adults. *Vaccine* 34(7): 942–949.
- LaVeist TA, Isaac LA and Williams KP (2009) Mistrust of health care organizations is associated with underutilization of health services. *Health Services Research* 44(6): 2093–2105.
- Martin LR and Petrie KJ (2017) Understanding the dimensions of anti-vaccination attitudes: The Vaccination Attitudes Examination (VAX) scale. *Annals of Behavioral Medicine* 51(5): 652–660.
- Motta M, Sylvester S, Callaghan T, et al. (2021) Encouraging COVID-19 vaccine uptake through effective health communication. *Frontiers in Political Science* 3: 630133.
- Murphy J, Vallières F, Bentall RP, et al. (2021) Psychological characteristics associated with COVID-19 vaccine hesitancy and resistance in Ireland and the United Kingdom. *Nature Communications* 12(1): 29.
- Nicholls LAB, Gallant AJ, Cogan N, et al. (2021) Older adults' vaccine hesitancy: Psychosocial factors associated with influenza, pneumococcal, and shingles vaccine uptake. *Vaccine* 39(26): 3520–3527.
- Noyman-Veksler G, Greenberg D, Grotto I, et al. (2021) Parents' malevolent personification of mass vaccination solidifies vaccine hesitancy. *Journal of Health Psychology* 26(12): 2164–2172.
- Ozawa S, Portnoy A, Getaneh H, et al. (2016) Modeling the economic burden of adult vaccine-preventable diseases in the United States. *Health Affairs* 35(11): 2124–2132.
- Pogue K, Jensen JL, Stancil CK, et al. (2020) Influences on attitudes regarding potential COVID-19 vaccination in the United States. *Vaccines* 8(4): 582.
- Privor-Dumm LA, Poland GA, Barratt J, et al. (2021) A global agenda for older adult immunization in the COVID-19 era: A roadmap for action. *Vaccine* 39: 5240–5250.
- Quinn SC, Jamison AM, An J, et al. (2019) Measuring vaccine hesitancy, confidence, trust and flu vaccine uptake: Results of a national survey of White and African American adults. *Vaccine* 37(9): 1168–1173.
- Reno C, Maietti E, Fantini MP, et al. (2021) Enhancing COVID-19 vaccines acceptance: Results from a survey on vaccine hesitancy in northern Italy. *Vaccines* 9(4): 378.
- Roller-Wirmsberger R, Lindner S, Kolosovski L, et al. (2021) The role of health determinants in the influenza vaccination uptake among older adults (65+): A scope review. *Aging Clinical and Experimental Research* 33: 2123–2132.
- Russell K, Chung JR, Monto AS, et al. (2018) Influenza vaccine effectiveness in older adults compared with younger adults over five seasons. *Vaccine* 36(10): 1272–1278.
- Ryan J and Malinga T (2021) Interventions for vaccine hesitancy. *Current Opinion in Immunology* 71: 89–91.
- Saied SM, Saied EM, Kabbash IA, et al. (2021) Vaccine hesitancy: Beliefs and barriers associated with COVID-19 vaccination among Egyptian medical students. *Journal of Medical Virology* 93(7): 4280–4291.
- Sallam M (2021) COVID-19 vaccine hesitancy worldwide: A concise systematic review of vaccine acceptance rates. *Vaccines* 9(2): 160.
- Santibanez TA, Nguyen KH, Greby SM, et al. (2020) Parental vaccine hesitancy and childhood influenza vaccination. *Pediatrics* 146(6): e2020007609.
- Schmid P, Rauber D, Betsch C, et al. (2017) Barriers of influenza vaccination intention and behavior - A systematic review of influenza vaccine hesitancy, 2005 - 2016. *PLoS One* 12(1): e0170550.
- Shahar G, Noyman-Veksler G, Itamar S, et al. (2017) Benevolent personification of the MoH increases compliance with an emergency polio vaccination. *Vaccine* 35(37): 5006–5010.
- Sheikh S, Biundo E, Courcier S, et al. (2018) A report on the status of vaccination in Europe. *Vaccine* 36(33): 4979–4992.
- Sprengholz P, Felgendreiff L, Böhm R, et al. (2021) Vaccination policy reactance: Predictors, consequences, and countermeasures. *Journal of Health Psychology*. Epub ahead of print 6 September 2021. DOI: 10.1177/13591053211044535.
- Suryadevara M (2021) Vaccine-preventable disease outbreaks. In: Domachowske J and Suryadevara M (eds) *Vaccines: A Clinical Overview and Practical Guide*. Cham: Springer International Publishing, pp.419–439.

- Talbird SE, La EM, Carrico J, et al. (2020) Impact of population aging on the burden of vaccine-preventable diseases among older adults in the United States. *Human Vaccines & Immunotherapeutics* 17(2): 332–343.
- Tan M, Straughan PT and Cheong G (2022) Information trust and COVID-19 vaccine hesitancy amongst middle-aged and older adults in Singapore: A latent class analysis Approach. *Social Science & Medicine* 296: 114767.
- Tan PS, Patone M, Clift AK, et al. (2021) Influenza, shingles and pneumococcal vaccine uptake, offer and refusal in adult populations at high-risk for COVID-19: A UK population-based Cohort Study. *SSRN Electronic Journal*. DOI: 10.2139/ssrn.3783784.
- Van Buynder P and Woodward M (2019) Importance and challenges of vaccination in older people. Available at: <https://medicinetoday.com.au/2019/february/feature-article/importance-and-challenges-vaccination-older-people> (accessed 17 May 2021).
- Williams L, Gallant AJ, Rasmussen S, et al. (2020) Towards intervention development to increase the uptake of COVID-19 vaccination among those at high risk: Outlining evidence-based and theoretically informed future intervention content. *British Journal of Health Psychology* 25(4): 1039–1054.
- Wysong CS, Ndwandwe D, Ryan J, et al. (2021) Vaccine hesitancy in the era of COVID-19: Could lessons from the past help in divining the future? *Human Vaccines & Immunotherapeutics* 18: 1–3.
- Wood L, Smith M, Miller CB, et al. (2019) The internal consistency and validity of the vaccination attitudes examination scale: A replication study. *Annals of Behavioral Medicine* 53(1): 109–114.