

1
2
3 **Q fever through consumption of unpasteurised milk and**
4 **milk products – a risk profile and exposure assessment**
5
6
7

8 Paul Gale¹, Louise Kelly¹, Rebecca Mearns², Jackie Duggan³, and Emma L. Snary¹
9

10 ¹Department of Epidemiological Sciences, Animal and Plant Health Agency (APHA),
11 Weybridge.
12

13 ²Animal and Plant Health Agency (APHA), Penrith.
14

15 ³Public Health England (PHE), Porton Down.
16
17
18
19
20
21

22 **Abbreviated Title:** *Coxiella burnetii* exposures through raw milk
23

24 **Correspondence:** Dr Paul Gale, Animal and Plant Health Agency, Weybridge, New
25 Haw, Addlestone, Surrey, KT15 3NB, UK. E-mail: paul.gale@apha.gsi.gov.uk
26
27
28
29
30

31 **Keywords:** *Coxiella burnetii*, risk assessment, exposure, unpasteurized milk
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Q fever is a zoonotic disease caused by the bacterium *Coxiella burnetii* which is endemic in cattle, sheep and goats in much of the world, including the United Kingdom (UK). There is some epidemiological evidence that a small proportion of cases in the developed world may arise from consumption of unpasteurised milk with less evidence for milk products such as cheese. Long maturation at low pH may give some inactivation in hard cheese and viable *C. burnetii* are rarely detected in unpasteurised cheese compared to unpasteurised milk. Simulations presented here predict that the probability of exposure per person to one or more *C. burnetii* through the daily cumulative consumption of raw milk in the UK is 0.4203. For those positive exposures, the average level of exposure predicted is high at 1,266 guinea pig intraperitoneal infectious dose 50% units (GP_IP_ID₅₀) per person per day. However, in the absence of human dose-response data, the case is made that the GP_IP_ID₅₀ unit represents a very low risk through the oral route. The available evidence suggests that the risks from *C. burnetii* through consumption of unpasteurised milk and milk products (including cheese) are not negligible but they are lower in comparison to transmission via inhalation of aerosols from parturient products and livestock contact.

INTRODUCTION

Q fever is a widespread, zoonotic disease caused by the bacterium *Coxiella burnetii* which is endemic in livestock including cattle, sheep and goats in much of the world including the United Kingdom (UK) (Maurin and Raoult 1999; Cutler *et al.* 2006).

The clinical manifestations of Q fever in humans are variable, ranging from asymptomatic to serious. Acute Q fever in humans usually manifests as an asymptomatic or mild flu-like disease with spontaneous recovery (Maurin and Raoult 1999). However, a small minority of patients present with more serious disease which can lead to serious complications and mortality. In some people, the disease can lead to a chronic infection that can manifest years later, even in the absence of primary, acute Q fever symptoms. Large community outbreaks of Q fever with over 3500 notified cases occurred in the Netherlands between spring 2007 until the end of 2009 (Schimmer *et al.* 2011). The aerosol route (inhalation of infected fomites) is considered to be the primary mode of human infection with *C. burnetii*. Infection via *C. burnetii* aerosols may occur from direct contact with the excretions and secretions from infected animals. These include milk, urine, faeces, vaginal mucus, semen and parturient fluids, which may contaminate newborn animals, placenta, or wool (Maurin and Raoult 1999).

Viable *C. burnetii* can be shed in milk from infected livestock including cattle (Bell *et al.* 1949; Enright *et al.* 1957) and have been detected (by passage in mice) in commercial unpasteurised milk samples (Loftis *et al.* 2010). However, the viability in those milk samples was demonstrated by intraperitoneal challenge rather than oral challenge. Indeed, the link between infection and clinical disease in humans through consumption of unpasteurised milk and milk products is unclear (EFSA 2010).

Maurin and Raoult (1999) in their review of Q fever conclude that although milk may

1
2
3 contain large amounts of *C. burnetii*, it is probably a minor route of Q fever
4
5 acquisition.

6
7 The aim of this paper is to assess the risks of *C. burnetii* infection through
8
9 consumption of unpasteurised milk and milk products. The paper first reviews the
10
11 epidemiological evidence for routes of transmission to humans and then addresses the
12
13 feasibility of developing a quantitative risk assessment for milk and milk products.
14
15 The availability of data limits this study to predicting human exposures to viable *C.*
16
17 *burnetii* through consumption of unpasteurised cows' milk. The risks from these
18
19 exposures are interpreted on the basis of infectivity through the oral route and placed
20
21 in context against the risks through other routes of transmission, in particular
22
23 inhalation of aerosols from livestock birth products. The potential risks of
24
25 unpasteurised milk products, namely cheeses, relative to milk are also considered.
26
27
28

29 **EPIDEMIOLOGICAL DATA ON ROUTES OF TRANSMISSION OF *C.***

30 ***BURNETII* TO HUMANS**

31
32 There are a number of routes identified by epidemiological studies for transmission of
33
34 *C. burnetii* to humans.
35
36

37 **Aerosols and direct contact with livestock**

38
39 The main routes of transmission are from livestock and companion mammals either
40
41 through the environment or through direct contact (Langley *et al.* 1988; Connolly *et*
42
43 *al.* 1990; Thomas *et al.* 1995; Schimmer *et al.* 2011). In this respect aerosolisation and
44
45 inhalation appear to be important (Maurin and Raoult 1999). Indeed outbreaks
46
47 associated with windborne transmission from farms and slaughter houses and within
48
49 meat processing plants are well-documented (Brouqui *et al.* 2004; Tissot-Dupont *et*
50
51 *al.* 2004; Wilson *et al.* 2010). The resistance of *C. burnetii* promotes its transmission
52
53 through aerosols, and there are suggestions of outbreaks of Q fever arising from *C.*
54
55
56
57
58
59
60

1
2
3 *burnetii* sources many years after release from an infected host (van Woerden *et al.*
4
5 2004).
6

7 **Consumption of unpasteurised milk and milk products**

8
9
10 There is epidemiological evidence, from the developed world, that cases of Q fever
11
12 have occurred where consumption of unpasteurised milk was the most likely cause.
13
14 The most recent of these was in Michigan (USA) in 2011 and involved five
15
16 individuals (Signs *et al.* 2012). However, suspected milk borne outbreaks are rare in
17
18 the UK. Unpasteurised cows' milk purchased by a patient was thought to be
19
20 responsible for an outbreak of Q fever in a London hospital in 1950 (Marmion and
21
22 Harvey 1956) and it was concluded that raw milk was responsible for an outbreak of
23
24 Q fever in a boys' detention centre in Staffordshire in April 1967 (Brown *et al.* 1968).
25
26 Although these studies are highly suggestive of the consumption of unpasteurised
27
28 milk being the source of the outbreak, there is still uncertainty associated with this
29
30 link (EFSA 2010). An epidemiological study of Q fever cases in the UK from 1984-
31
32 1994 has reported that, out of 1,117 cases of Q fever investigated, three cases were
33
34 reported to have consumed unpasteurised milk (Pebody *et al.* 1996). With the possible
35
36 exception of an outbreak in France (Raoult *et al.* 2000) where unpasteurised milk was
37
38 also consumed, there have been no outbreaks reported due to the consumption of milk
39
40 products (such as cheese) made from unpasteurised milk, so if cases are occurring
41
42 they are likely to be sporadic in nature.
43
44
45
46

47 **PROPERTIES OF THE *C. BURNETII* ORGANISM WITH RELEVANCE TO** 48 49 **ASSESSING THE RISKS THROUGH UNPASTEURISED MILK AND MILK** 50 51 **PRODUCTS**

52
53
54 **Infectious *C. burnetii* have been isolated from milk**
55
56
57
58
59
60

1
2
3 There is much evidence that *C. burnetii* is viable to some degree in unpasteurised
4 milk. Thus, Loftis *et al.* (2010) have confirmed the viability by passage in mice of *C.*
5 *burnetii* in at least two and maybe four of six PCR-positive commercial,
6 unpasteurised milk samples. Experiments conducted in the 1940s and 1950s showed
7 that naturally infected cows' milk can infect guinea pigs and mice (Bell *et al.* 1949;
8 Enright *et al.* 1957) albeit through intraperitoneal challenge. Levels of viable *C.*
9 *burnetii* in milk are often expressed in units of "guinea pig intraperitoneal infectious
10 dose 50%" or GP_IP_ID₅₀. This is the dose which, when given to each and every
11 member of a group of guinea pigs through intraperitoneal challenge, results in 50%
12 being infected (Enright *et al.* 1957).
13
14
15
16
17
18
19
20
21
22
23
24

25 ***C. burnetii* will not multiply in the milk**

26
27 *C. burnetii* is an obligate intracellular bacterium that relies exclusively on a
28 eukaryotic cell for growth (Omsland and Heinzen 2011). This has direct relevance to
29 assessing the risks through food and environmental routes because *C. burnetii* does
30 not grow outside the intracellular environment of the host cell. Thus for the purpose
31 of risk assessment it is assumed that multiplication of the pathogen in milk and milk
32 products does not occur.
33
34
35
36
37
38
39
40

41 **The environmental morphotype is highly resistant**

42
43 The organism has a two stage development cycle, with two distinct morphological
44 variants, or morphotypes namely the large cell variant (LCV) and the small cell
45 variant (SCV) (Minnick and Reghavan 2012). Unlike other obligate intracellular
46 bacteria, *C. burnetii* has spore-like environmental stability due to the resistance of the
47 SCV (Oyston and Davies 2011). Indeed *C. burnetii* can potentially survive for years in
48 the environment, being highly resistant to chemical and physical stresses, including
49 disinfectants, desiccation, UV light, sonication and osmotic stress (Oyston and Davies
50
51
52
53
54
55
56
57
58
59
60

1
2
3 2011). Monocytes and macrophages are the major targets of *C. burnetii* (Amara *et al.*
4 2012) and spread around much of the body. The placenta is a tissue rich in
5
6 macrophages and placental macrophages harbour *C. burnetii* (Amara *et al.* 2012).
7
8

9 **The unit of *C. burnetii* infectivity in milk**

10
11 Macrophages occur in bovine milk (Paape *et al.* 2003). Within the macrophage, a high
12
13 density mixture of LCVs and SCVs exists in the parasitophorous vacuole (Minnick
14
15 and Reghavan 2011). The LCV is very fragile (Minnick and Reghavan 2011). While
16
17 the long survival of *C. burnetii* infectivity in milk (Combiesco *et al.* 1953; Zubkova
18
19 1957) supports the case for SCVs being present in milk, there is no information on the
20
21 relative proportions of LCV to SCV in macrophages in fresh milk. PCR would detect
22
23 DNA from both SCVs and LCVs in milk, with the SCV representing a higher risk to
24
25 human health due to its greater chance of surviving not only in the milk environment
26
27 but also in the digestive tract during initiation of infection after consumption of
28
29 infected milk.
30
31
32

33 *Estimating the number of C. burnetii bacteria comprising a GP_IP_ID₅₀ in milk.*

34
35 Ideally the exposure units for a quantitative risk assessment should be in terms of the
36
37 number of viable bacteria such that the outputs can be used directly in a dose-response
38
39 model should one become available (see below). Thus, expressing *C. burnetii*
40
41 exposures in terms of the numbers of GP_IP_ID₅₀ raises the question of how many *C.*
42
43 *burnetii* bacteria comprise GP_IP_ID₅₀. Guatteo *et al.* (2007) used a PCR method to
44
45 estimate titres in cows' milk by comparison of PCR results with those from solutions
46
47 with a known *C. burnetii* concentration obtained by serial dilution of an external
48
49 positive control. Comparison of quantitative PCR results of Guatteo *et al.* (2007) for
50
51 *C. burnetii* in dairy milk with the numbers of GP_IP_ID₅₀ recorded in milk by Enright
52
53 *et al.* (1957) suggest there could be between 2 and 112 *C. burnetii* organisms per
54
55
56
57
58
59
60

1
2
3 GP_IP_ID₅₀ in milk. This is calculated as follows. The distribution of GP_IP_ID₅₀ in
4
5 milk from 18 naturally infected and shedding dairy cows appears to be log-Normal
6
7 with a mean of 98.75 per ml (Enright *et al.* 1957). The averaged median and averaged
8
9 maximum (for n = 5 cows) number of *C. burnetii* per ml of milk (quantified by PCR
10
11 in Guatteo *et al.* (2007)) were 213 and 11,073, respectively. Since the mean of a log-
12
13 Normal distribution is between the median and the maximum, the mean number of *C.*
14
15 *burnetii* is between 213 and 11,073 per ml of milk which is equivalent to 98.75
16
17 GP_IP_ID₅₀. This suggests there are between $(213/98.75)^2$ and $(11,073/98.75)^2$ 112 *C.*
18
19 *burnetii* organisms per GP_IP_ID₅₀ in milk.
20
21

22 23 **FEASIBILITY OF DEVELOPING A QUANTITATIVE RISK ASSESSMENT** 24 25 **FOR *C. BURNETII* INFECTION THROUGH MILK AND MILK PRODUCTS**

26
27 Milk products include cheese, yoghurt, butter and cream. Milk and milk products may
28
29 be sourced from cattle, sheep and goats. Thus data are needed for each of these
30
31 species although in terms of consumption patterns, the use of cows' and goats' milk is
32
33 more common than for sheep's milk and unpasteurised cheese and yoghurt are
34
35 normally made from cows' or goats' milk in England.
36
37

38 **Availability of dose-response data for infection of *C. burnetii* through the oral** 39 40 **route in humans**

41
42 *C. burnetii* is highly infectious through inhalation with the risk of infection from a
43
44 single bacterium estimated to be as high as 0.9 in guinea pigs (Jones *et al.* 2006).
45
46 There are insufficient data for a dose-response model for the oral route in humans.
47
48 Indeed, transmission by the oral route of *C. burnetii* is controversial (Eldin *et al.*
49
50 2013) and Cerf and Condron (2006) challenge the designation of *C. burnetii* as a
51
52 foodborne pathogen. This suggests *C. burnetii* may not be very infectious through the
53
54 oral route. Over a period of one month, Krumbieoel and Wisniewski (1970) gave 34
55
56
57
58
59
60

1
2
3 volunteers unpasteurised milk that was naturally infected with *C. burnetii*. The
4
5 volunteers consumed an average of 4.5 litres of milk under supervision during the
6
7 month of the trial. None of the volunteers developed any clinical symptoms even after
8
9 12 years and serum samples taken 1 month and 2 months after initial ingestion
10
11 showed no evidence of seroconversion. The authors concluded that either the milk
12
13 may have contained a strain that is not infectious to humans or that an inapparent
14
15 infection without serological response had occurred. Two of 11 patients in an asylum
16
17 in Portugal given *C. burnetii* in food showed signs of seroconversion by complement
18
19 fixation assay and none developed clinical symptoms (Fonseca *et al.*, 1949). The doses
20
21 administered in that study were not specified and it is unlikely there will ever be
22
23 sufficient dose-response data for *C. burnetii* infection in humans through the oral
24
25 route to undertake a quantitative risk assessment. Even if a foodborne outbreak could
26
27 be detected, calibration of a dose-response would currently be difficult because of the
28
29 lack of a straightforward enumeration method for viable organism.
30
31
32

33 34 **Availability of data for estimation of exposure through consumption of** 35 36 **unpasteurised milk**

37
38 The data required for predicting levels of exposure through consumption of
39
40 unpasteurised milk are set out in the exposure pathway in Figure 1.
41

42
43 *Prevalence of C. burnetii in livestock in UK.* *C. burnetii* is endemic in UK dairy cattle
44
45 herds which, in the case of dairy herds in Northern Ireland at least, have higher
46
47 prevalences than beef cattle herds (McCaughey *et al.* 2010). Reported prevalences in
48
49 bulk tank milk (BTM) samples from dairy cattle herds in England and Wales range
50
51 from 22% (ELISA) to 69.7% (PCR) (Paiba *et al.* 1999; Valergakis *et al.* 2012).
52
53 McCaughey *et al.* (2010) present data for between herd and within herd prevalence
54
55 according to herd size. There are fewer data for sheep and goats in England and Wales
56
57
58
59
60

1
2
3 with unpublished estimates of individual animal prevalences in sheep and goats of
4 0.92% and 0.78%, respectively, by ELISA (Lambton *et al.* unpublished) although data
5
6
7 have been published for Northern Ireland (McCaughey *et al.* 2010). The advent of
8
9
10 PCR has enabled detection of *C. burnetii* DNA and even quantification of *C. burnetii*
11
12 DNA in milk as used by Valergakis *et al.* (2012) for BTM from dairy cattle in south-
13
14 west England. However, the problem with PCR is that it gives no information on the
15
16 viability of the organism. ELISA techniques, as used by Lambton *et al.* (unpublished)
17
18 and Paiba *et al.* (1999), may over-estimate prevalence because animals may be sero-
19
20 positive for life, but not actively infected with the bacteria, although some may later
21
22 convert from sero-positive to sero-negative.
23

24
25 *Probability of infected livestock shedding C. burnetii.* Shedding of *C. burnetii* differs
26
27 among ruminant species, milk being the primary route of shedding in cattle and goats
28
29 (Rodolakis *et al.* 2007). Sheep shed mainly in the faeces and vaginal mucus and to a
30
31 lesser extent in milk (Rodolakis *et al.* 2007). Indeed, for infected goats, 31 – 38%
32
33 shed in milk (Rousset *et al.* 2009). Roest *et al.* (2012) reported that all *Coxiella*-
34
35 inoculated goats excreted *C. burnetii* DNA in milk post partum. Guatteo *et al.* (2012)
36
37 give data on the number of infected cows which were shedding at days 14, 21 and 28
38
39 post abortion due to *C. burnetii*.
40
41

42
43 *Levels of viable C. burnetii in milk of shedding livestock.* Enright *et al.* (1957) used a
44
45 guinea pig bioassay approach to measure *C. burnetii* in unpasteurised cows' milk. The
46
47 great advantage (for the purpose of data for risk assessment) of guinea pig bioassay
48
49 over PCR is that it determines viable pathogen in terms of the numbers of
50
51 GP_IP_ID₅₀. Enright *et al.* (1957) reported that milk from 18 of 137 individual cows
52
53 in a naturally-infected dairy herd contained viable *C. burnetii*. Titration of those
54
55 positive milk samples showed levels of 1,000 (n = 3), 100 (n = 5), 10 (n = 5) and one
56
57
58
59
60

1
2
3 (n = 5) GP_IP_ID₅₀ per 2 ml. The mean number of *C. burnetii* is therefore 98.8
4
5 GP_IP_ID₅₀ per ml of milk from a shedding cow. Bell *et al.* (1949) reported a
6
7 maximum of 10⁵ GP_IP_ID₅₀ (presumably per ml) in milk from a cow with mastitis.
8
9 This value is excluded from the analysis here because the mastitis may have increased
10
11 the measured densities of *C. burnetii* in milk by two mechanisms; namely i) by
12
13 increasing the number of *C. burnetii*-infected macrophages actually in the milk (Paape
14
15 *et al.* 2003) and ii) by decreasing the volume of milk produced. From a risk
16
17 assessment perspective, the milk from cows with mastitis would not enter the food
18
19 chain. Similarly data, including a maximum of 10,000 GP_IP_ID₅₀ per 2 ml of milk,
20
21 recorded from a dairy cow (Enright *et al.* 1957) were excluded here because that cow
22
23 was experimentally (as opposed to naturally) infected by introduction via the teat
24
25 canal.
26
27
28
29 There are no quantitative data on levels of viable *C. burnetii* in sheep and goats' milk.
30
31 *Duration of shedding in milk.* Shedding of *C. burnetii* DNA in milk from infected
32
33 goats stopped 38 days post partum (Roest *et al.* 2012), although Arricau-Bouvery *et*
34
35 *al.* (2003) detected *C. burnetii* DNA in goats' milk 52 days after abortion. Guatteo *et*
36
37 *al.* (2012) write that three infected cows were identified as persistent shedders in that
38
39 they were shedding relatively high levels at 14, 21 and 28 days post abortion.
40
41
42 Unfortunately Guatteo *et al.* (2012) do not give data for more than two weeks (albeit
43
44 one month after abortion). Enright *et al.* (1957) give data showing that infected cattle
45
46 can shed in milk for periods of more than one year. They found that the milk of four
47
48 positive cows was still positive 205 days after each had calved, and one of the animals
49
50 was found to be still shedding 1,000 GP_IP_ID₅₀ per 2 ml of milk. Serologic evidence
51
52 indicated that this animal was infected at least 405 days prior to the second milk
53
54
55
56
57
58
59
60

1
2
3 sampling. *C. burnetii* could not be found in the milk of the other three cows at the
4
5 time point of the second milk sampling.
6

7 *Levels of consumption of unpasteurised milk in the UK.* There are no data available on
8
9 consumption of unpasteurised milk in the UK. The mean consumption for milk has
10
11 been estimated as 0.127 kg/person/day (Department of Health 2011). This is for
12
13 (pasteurised presumably) whole milk (3.8% fat) among the 19 to 64 year old age
14
15 group, and includes males and females and, importantly, consumers only.
16
17

18 **Availability of data for estimation of exposure through consumption of** 19 20 **unpasteurised milk products** 21

22
23 Recently some papers have been published reporting results of PCR studies for
24
25 detection of *C. burnetii* DNA in unpasteurised cheeses. As an example, Capuano *et al.*
26
27 (2012) reported 21.3% of cheeses made in Southern Italy from unpasteurised milk
28
29 were PCR-positive. Hirai *et al.* (2012) reported 7 of 41 commercial cheeses made
30
31 from unpasteurised milk were PCR-positive, compared to 20 of 96 made from
32
33 pasteurised milk. To date, however, no published paper has been found giving counts
34
35 of viable *C. burnetii* in cheese with which to compare with the unpasteurised milk
36
37 data of Enright *et al.* (1957).
38
39

40 *Proportion of C. burnetii removed with the whey during cheese-making.* Removal of
41
42 the whey during cheese production could eliminate a considerable proportion of the
43
44 pathogen, although there are no specific data for *C. burnetii* in this respect. Anon
45
46 (2013) present the relative proportions of milk components that remain in the whey or
47
48 partition into the cheese. The data show there are two exclusive outcomes. Thus,
49
50 around 95% of the water soluble components (namely water, lactose and non-
51
52 precipitated proteins) remain in the whey and are removed with the whey, while 95%
53
54 of the water-insoluble components, namely fat and precipitated casein proteins go into
55
56
57
58
59
60

1
2
3 the cheese. Thus it could be that either 95% of the *C. burnetii* are lost with the whey,
4
5 or alternatively that 95% of the *C. burnetii* precipitate into the curds which go on to
6
7 become cheese. The amount of whey is to some extent affected by salt content and
8
9 impacts on moisture content of the cheese which differ between soft and hard cheeses.
10
11 For *C. burnetii*, this could be addressed by considering the physical properties of the
12
13 small cell variant at low pH or after rennet treatment.
14

15
16 *Survival of C. burnetii in milk and milk products with time.* Much of the data on *C.*
17
18 *burnetii* survival was obtained from experiments in the 1940s and 1950s. Although *C.*
19
20 *burnetii* is inactivated by pasteurisation, there is little evidence that any of the
21
22 processes used for unpasteurised cheese, cream or butter production would
23
24 significantly inactivate *C. burnetii*. Jellison *et al.* (1948) reported the presence and
25
26 persistence of infectious *C. burnetii* in butter made from naturally infected milk and
27
28 *C. burnetii* survived in milk (dried 37°C) for 30 – 60 days and in cheese made from
29
30 infected milk for 17 – 46 days (Babudieri and Moscovici 1950). *C. burnetii* survived
31
32 in sterile milk at room temperature for 125 days (Zubkova 1957). There is one study
33
34 where viable pathogen has been detected in a cottage-type cheese after 42 days (Sipka
35
36 1958). The data are not quantitative and inactivation rates cannot be determined.
37
38

39
40 *Effect of low pH.* Based on experience of freezing *Coxiella* in acidic media it is
41
42 believed that *Coxiella* may retain better viability in cheese at neutral pH than at pH
43
44 5.0 (Robert Heinzen, National Institute of Health, USA, pers. comm.). This is
45
46 supported by data from the 1950s that milk collected and maintained in aseptic
47
48 conditions remained infective for at least 45 days, but if allowed to become sour
49
50 (lower pH) it ceased to be infective within 24 hours (Combiesco *et al.* 1953).
51
52

53 54 **Summary of identified data gaps**

55
56
57
58
59
60

1
2
3 There are significant data gaps in the level of knowledge of *C. burnetii* with little or
4
5 no information on:-
6

- 7 1. Current farm prevalence and within herd/flock prevalence of *C. burnetii*
8 (ELISA and PCR data are available but will overestimate the prevalence);
9
- 10 2. Levels and viability of *C. burnetii* in sheep and goats' milk;
11
- 12 3. Survival of *C. burnetii* in unpasteurised milk and milk products;
13
- 14 4. Survival and removal of *C. burnetii* during the cheese-making processes and
15 manufacture of other milk products; and
16
- 17 5. Dose-response data for humans through the oral route.
18
19
20

21
22 The data gaps in part reflect the difficulties in routine culture of *C. burnetii* (Oyston
23 and Davies 2011) and also the lack of data on the viability of the organisms when
24 DNA is detected by PCR methods. It is concluded that there are insufficient data to
25 develop a quantitative risk assessment for *C. burnetii* in sheep and goats' milks, or in
26 milk products including cheese. There are, however, sufficient data to predict
27 exposures of *C. burnetii* (albeit in terms of GP_IP_ID₅₀) through consumption of
28 unpasteurised cows' milk and this is now described.
29
30
31
32
33

34 **A QUANTITATIVE EXPOSURE ASSESSMENT FOR CONSUMPTION OF** 35 36 37 38 39 40 41 **UNPASTEURISED COWS' MILK**

42 The specific question that the exposure assessment will address is, "*What is the*
43 *exposure to C. burnetii of a consumer through the cumulative consumption of*
44 *unpasteurised cows' milk over the period of one day?*". This may be broken down
45 into two outputs, namely:-
46
47
48
49

- 50 1. The probability of exposure through the cumulative daily consumption of
51 unpasteurised milk; and
52
53
54
55
56
57
58
59
60

1
2
3 2. The level of exposure, given exposure has occurred, to a person through
4 consumption of unpasteurised milk over the period of a day.
5
6

7 The exposure pathway is shown in Figure 1. The model parameters, based on the data
8 described previously are given in Table 1. The between-herd and within-herd
9 prevalences used are those for Northern Ireland (McCaughey *et al.* 2010) and are
10 broken down according to herd size. It is assumed that the probability that an infected
11 cow is shedding (p_{shedding}) is given by 22/72 (0.3055) according to the summed data
12 of Guatteo *et al.* (2012) over days 14, 21 and 28 post abortion due to *C. burnetii*. As a
13 worst case, it is assumed that a cow which does shed *C. burnetii* does so for the whole
14 year. A Normal distribution gave a good fit ($\chi^2 = 0.667, 1 \text{ df}; P = 0.88$) to the \log_{10} -
15 transformed titres for GP_IP_ID₅₀/ml milk from shedding cows and was used in the
16 risk assessment.
17
18
19
20
21
22
23
24
25
26
27
28

29 The quantitative model was implemented in Microsoft Excel, using the @RISK
30 software package to incorporate variation associated with herds and individual
31 animals in relation to infection, lactation and the levels of *C. burnetii* in milk. There
32 are no quantitative data to allow estimation of a decay rate for *C. burnetii* in milk and
33 it is assumed that no decay occurs between milking and consumption of fresh milk.
34
35
36
37
38
39

40 **The predicted mean level of *C. burnetii* in unpasteurised cows' bulk tank milk** 41 **(per herd) in the UK** 42 43 44

45 The model simulated each cow in a herd on a given day and each iteration of the
46 model represents the milk produced from a single herd on that day. In total 500,000
47 iterations were run representing 500,000 herd-days. For each iteration, the number of
48 cows (H) in the herd was randomly selected from the empirical distribution of herd
49 sizes for the 81 herds in England and Wales known to be producing unpasteurised
50 milk (data provided by UK Food Standards Agency). Taking into account the
51
52
53
54
55
56
57
58
59
60

1
2
3 between-herd prevalence (p_{Herd}), the within herd prevalence ($p_{\text{Within_herd}}$), the
4
5 probability of lactating ($p_{\text{Lactating}}$) and the probability of shedding given infection
6
7 (p_{Shedding}) (Table 1), binomial distributions were used to simulate whether or not each
8
9 cow in the herd was producing infected milk on that day. For each shedding cow the
10
11 number of *C. burnetii* GP_IP_ID₅₀S (C_{day}) contributed to the BTM on that day was
12
13 calculated as the product of the volume of milk produced by that cow on that day (V)
14
15 and the concentration (C_{ml}) of infectivity in the milk as drawn from a log-Normal
16
17 distribution fitted to the data of Enright *et al.* (1957). The total volume of milk in the
18
19 BTM was calculated as the sum of the volumes of milk (V) produced by all lactating
20
21 cows in the herd on that day. From the total *C. burnetii* shed from all cows ($C_{\text{BTM,Day}}$)
22
23 and the total volume of milk produced by the herd, the mean level of *C. burnetii* in the
24
25 bulk tank milk ($C_{\text{BTM,Litre}}$) from the given herd on a given day was calculated.
26
27

28
29 Although there will be variation between the individual cows within the herd in the
30
31 amount of *C. burnetii* shed each day, the mean is appropriate here because the milk in
32
33 the bulk tank is stirred, and furthermore is not mixed with milk from other cattle
34
35 herds. This is because of restrictions in England on the sale of unpasteurised cows'
36
37 milk (Anon 2006). The simulated mean level ($C_{\text{BTM,Litre}}$) of *C. burnetii* is 4,189
38
39 GP_IP_ID₅₀ per litre of unpasteurised milk from the bulk tank with 2.5th and 97.5th
40
41 percentiles of 0 and 26,848 GP_IP_ID₅₀ per litre, respectively. This represents the
42
43 mean for the 81 unpasteurised milk-producing herds in England and Wales.
44
45

46
47 *Validation of predicted levels of infectivity in unpasteurised milk against published*
48
49 *PCR data.* The seemingly high values predicted for infectivity levels in BTM reflect
50
51 the values of up to 1,000 GP_IP_ID₅₀S per 2 ml of unpasteurised milk (Enright *et al.*
52
53 1957) to which the log-Normal distribution, used in the simulation here, was fitted.
54
55 The distribution for the number of *C. burnetii* GP_IP_ID₅₀ per ml of BTM milk as
56
57
58
59
60

1
2
3 simulated is presented in Figure 2. The GP_IP_ID₅₀s per ml are converted to
4
5 logarithms to enable direct comparison with the distribution presented in Valergakis
6
7 *et al.* (2012) of the qPCR units per ml of milk. The two distributions are similar *in*
8
9 *shape* with each having two peaks. The “negative samples” peak reflects negative
10
11 herds and also positive herds with non-shedding cows on that day. However, although
12
13 the shapes of the distributions have some similarity, the simulated *C. burnetii*
14
15 GP_IP_ID₅₀ values are shifted by some three logs lower compared to the qPCR data
16
17 of Valergakis *et al.* (2012). The arithmetic mean number of qPCR units in the BTM of
18
19 Valergakis *et al.* (2012) is estimated to be 7.36×10^6 per litre and 1,800-fold higher
20
21 than the simulated mean level of 4,189 GP_IP_ID₅₀ per litre. Thus the model would
22
23 appear to underestimate the levels of *C. burnetii* in BTM by some three orders of
24
25 magnitude compared to PCR data obtained from BTM in the south-west of England.
26
27 However, there are three considerations which could account for this discrepancy:-
28
29

- 30
31
32 1. The PCR primers used by Valergakis *et al.* (2012) target a sequence of DNA
33
34 that is present in multiple copies in each *C. burnetii* organism. Thus Klee *et al.*
35
36 (2006) report 23 IS1111 elements in the genome of the Nine Mile strain,
37
38 although the number varied between seven and 110 in other isolates;
- 39
40 2. Some of the DNA detected by the PCR may represent non-viable (dead) *C.*
41
42 *burnetii* organisms; and
- 43
44 3. A GP_IP_ID₅₀ from milk may comprise more than one bacterium such that
45
46 multiple *C. burnetii* genomes are present in a GP_IP_ID₅₀. As discussed
47
48 above, it is estimated here that there are between 2 and 112 *C. burnetii*
49
50 organisms per GP_IP_ID₅₀ in milk.
51
52

53
54 It is concluded, therefore, that the predicted GP_IP_ID₅₀ in BTM (Figure 2) are not
55
56 inconsistent with the published qPCR data for BTM (Valergakis *et al.* 2012). Thus if
57
58
59
60

1
2
3 each GP_IP_ID₅₀ comprised 50 bacteria each with 20 copies of the PCR target
4
5 sequence (Klee *et al.* 2006), then the number of PCR copies would be 1,000-fold the
6
7 number of GP_IP_ID₅₀. This could account for the differences in the predicted number
8
9 of GP_IP_ID₅₀ per ml of milk (Figure 2) and observed number of PCR copies/ml
10
11 (Valergakis *et al.* 2012).
12

13 14 **The probability and level of human exposure to *C. burnetii* due to the** 15 16 **consumption of unpasteurised cows' milk**

17
18 Exposures were drawn from a Poisson distribution with a mean of the product of the
19
20 simulated GP_IP_ID₅₀ per litre of unpasteurised milk ($C_{\text{BTM,Litre}}$) and the amount of
21
22 milk (0.127 kg) consumed per person per day ($M_{\text{Litre/Day}}$). The exposure assessment
23
24 predicted that the probability of exposure to viable *C. burnetii*, i.e. one or more
25
26 GP_IP_ID₅₀, through the consumption of unpasteurised milk in the UK is 0.4203 per
27
28 person per day and that the daily exposures, to those who are exposed, will be
29
30 relatively high with a mean 1,266 GP_IP_ID₅₀ per person day and 2.5th and 97.5th
31
32 percentiles of 2 and 7,524 GP_IP_ID₅₀ per person per day, respectively. The
33
34 magnitudes of these exposures may be over-estimated for three reasons which relate
35
36 to whether an infected animal is shedding on a given day:-
37
38
39

- 40 1. Duration of shedding. It is assumed that an infected cow which is shedding in
41
42 milk does so every day.
- 43
44 2. Use of serological data (ELISA) for between herd and within herd prevalences
45
46 may overestimate the proportion of animals infected at any given time.
- 47
48 3. Use of PCR data to estimate the probability of shedding by a cow that
49
50 experienced abortion due to *C. burnetii* (Guatteo *et al.* 2012) assumes that all
51
52 *C. burnetii* DNA in milk from an infected cow does indeed represent viable *C.*
53
54
55
56
57
58
59
60
burnetii.

1
2
3 In a sensitivity analysis the duration of shedding was reduced to one month (from one
4 year). This decreased the probability of exposure by four-fold to 0.1048 per person
5 per day and decreased the mean level of exposure in those who were exposed by
6 three-fold to 411.5 GP_IP_ID₅₀ per person per day (2.5th and 97.5th percentiles of 1
7 and 2,290 GP_IP_ID₅₀ per person per day, respectively)..

14 **ASSESSING THE RISK OF INFECTION THROUGH CONSUMPTION OF** 15 **UNPASTEURISED COWS' MILK**

16
17
18 The predictions here suggest that consumers of unpasteurised cows' milk are
19 frequently exposed to relatively high loadings of *C. burnetii*. Although it is not known
20 how to convert GP_IP_ID₅₀ units into human oral ID₅₀s (because of lack of human
21 oral dose-response data), it is likely that each one presents a low risk to humans
22 through the oral route. There are three lines of evidence that support this. These
23 reflect the route of infection, the mechanism of infection and the genotype. First, with
24 respect to the route of infection, Fonseca *et al.* (1949) demonstrated high infection
25 rates by *C. burnetii* in humans through intradermal challenge but low rates through
26 oral challenge (although it is not known if the challenge doses were the same).
27 Intraperitoneal challenge is similar to intradermal challenge and thus it may be argued
28 on the basis of the data of Fonseca *et al.* (1949) that a GP_IP_ID₅₀ presents a low risk
29 through the oral route (since 2 of 11 humans were infected by oral challenge
30 compared to 29 of 29 humans by intradermal in Fonseca *et al.* (1949)). Second, with
31 respect to the mechanism of infection, *C. burnetii* targets macrophages within the host
32 tissues in the infection process (Amara *et al.* 2012) and there are far fewer
33 macrophages in the gastrointestinal tract compared to the lung. Thus the lung tissue
34 with a high number of alveolar macrophages is a prime environment for initial
35 infection and the most common route of infection by *C. burnetii* (Mike Minnick,
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Montana University, personal communication). Thus, *C. burnetii* is less infectious
4 through the oral route compared to inhalation. Third, the genotype of *C. burnetii* may
5 be important in relation to human infection. The genotypes of *C. burnetii* found in a
6 study of commercially available cows' milk in Europe are similar with a dominant
7 genotype that is only incidentally found in humans thus suggesting that the risk of
8 obtaining Q fever via exposure to infected cattle may be much lower than via
9 exposure to infected small ruminants (Tilburg *et al.* 2012). Indeed, sequencing work
10 at AHVLA (Richard Ellis, AHVLA, personal communication) shows that sheep
11 isolates of Q fever are most closely related to those in humans. This is important as
12 sheep shed *C. burnetii* to a lesser extent in milk (Rodolakis *et al.* 2007) and there is
13 little sheep milk consumption in the UK.

24 25 26 27 **RISKS OF INFECTION THROUGH UNPASTEURISED MILK COMPARED** 28 29 **TO OTHER MILK PRODUCTS, NAMELY CHEESE**

30
31 The risks through unpasteurised cheeses may be lower than those for unpasteurised
32 milk. Eldin *et al.* (2013) conclude that although there is a high prevalence of infection
33 in farm animals in France, consumption of cheese does not seem to pose a public
34 health risk for transmission of *C. burnetii* because the pathogen is not viable. This
35 may reflect inactivation of the pathogen in some cheeses. Indeed, the viability of *C.*
36 *burnetii* appears to be lost in cheese with Hirai *et al.* (2012) reporting no viable *C.*
37 *burnetii* in 7 unpasteurised milk cheeses which were PCR-positive. However, *C.*
38 *burnetii* infectivity for guinea pigs was retained in a cottage-type cheese for a period
39 of observation of 42 days (Sipka 1958). Typically, the pH of cheese ranges from 5.1
40 to 5.9 with a few exceptions such as Camembert which has a pH of 7.44 (World's
41 Healthiest Foods 2013). The pH of cheddar cheese is 5.0 to 5.2 with >60 days'
42 maturation (often 6 to 24 months) (Banks 2006). Semi-soft cheeses such as Caerphilly
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 have pH values of 4.6 – 6.2 with 10-14 days' maturation (Banks 2006). It is possible
4
5 that the combination of time/process conditions (e.g. lower pH and longer maturation
6
7 times) in the manufacture of some hard cheeses is not conducive to survival of *C.*
8
9 *burnetii*. This is consistent with viable *C. burnetii* rarely being detected in
10
11 unpasteurised cheese (Hirai *et al.* 2012) compared to unpasteurised milk (Enright *et*
12
13 *al.* 1957; Loftis *et al.* 2010) and with stronger epidemiological evidence for human
14
15 cases through unpasteurised milk compared to unpasteurised cheese.
16
17

18 **RISKS OF INFECTION THROUGH MILK AND MILK PRODUCTS**

19 **COMPARED TO OTHER ROUTES**

20
21
22 Inhalation of aerosols from parturient fluids of infected animals is the primary mode
23
24 of transmission of *C. burnetii* to humans while ingestion (mainly through drinking
25
26 unpasteurised milk) is probably a minor factor in the transmission and is now even a
27
28 point of controversy (Maurin and Raoult 1999; Cerf and Condron 2006). This may
29
30 reflect a combination of lower exposures and lower infectivity through unpasteurised
31
32 milk, as is now discussed.
33
34

35 **Relative infectivity of *C. burnetii* from oral consumption of milk compared to**

36 **inhalation of aerosols from birth products.**

37
38
39 The *C. burnetii* bacterium may be less infectious through unpasteurised milk
40
41 compared to aerosolised bacteria from livestock births or abortions because, as
42
43 discussed above, *C. burnetii* is less infectious through the oral route compared to
44
45 inhalation reflecting the greater numbers of target macrophages in the lung. In
46
47 addition it is proposed here that, in terms of tissue origin, *C. burnetii* derived from
48
49 birth product tissue may be more infectious (on average per bacterium or genome
50
51 equivalent) through a given route (e.g. intraperitoneal challenge) than that derived
52
53 from milk. Thus it is estimated above that there may be between 2 and 112 *C. burnetii*
54
55
56
57
58
59
60

1
2
3 organisms per GP_IP_ID₅₀ in milk. In contrast, there is considerable evidence that a
4
5 GP_IP_ID₅₀ from the placenta comprises just one *C. burnetii* organism. Thus Kersh *et*
6
7 *al.* (2013) recorded 1.5 to 2.5 x 10⁸ genome equivalents per gram of placenta from
8
9 goats and Hansen *et al.* (2011) reported 10⁹ *icd* gene copies (single copy per
10
11 bacterium) per ml of eluate from cattle cotyledons in parturient cattle. These values
12
13 agree well with the average of 5.0 x 10⁸ GP_IP_ID₅₀ per gram of ovine placental
14
15 tissue (Welsh *et al.* 1951). Further experimental work is needed to confirm the
16
17 number of *C. burnetii* bacteria in a GP_IP_ID₅₀ from milk. The argument presented
18
19 here that there are between 2 and 112 bacteria per GP_IP_ID₅₀ from milk hinges on
20
21 the maximum of 1,000 GP_IP_ID₅₀ per 2 ml of milk as recorded by Enright *et al.*
22
23 (1957).
24
25

26
27 It would also be of interest to know whether the ratio of SCV to LCV is the same in
28
29 birth products as in milk. Significant differences would affect whether a *C. burnetii*
30
31 organism (as represented by a genome equivalent or single bacterium) in milk is as
32
33 infectious, on average, as one from birth products, for example. These are important
34
35 considerations for developing any risk assessment to compare risks through milk and
36
37 aerosolised birth products, particularly since relatively few PCR-based studies address
38
39 the viability in milk (Loftis *et al.* 2010).
40
41

42 43 **Relative exposures through birth products compared to unpasteurised milk**

44
45 The exposures to humans may be lower through consumption of unpasteurised milk
46
47 than through aerosols from birth products. Huge numbers of bacteria are produced
48
49 during abortion caused by *C. burnetii* and via livestock birth products (10⁹
50
51 GP_IP_ID₅₀s per gram of sheep placenta (Welsh *et al.* 1951)) compared to the mean
52
53 of 98.8 GP_IP_ID₅₀s per ml of unpasteurised milk from shedding cows. Roest *et al.*
54
55 (2012) give semi-quantitative PCR data on excretion of *C. burnetii* in goats' milk,
56
57
58
59
60

1
2
3 demonstrating very low levels compared to goat placental tissue. A recent air
4
5 sampling study (Kersh *et al.* 2013) on a goat farm in the USA has shown the mean
6
7 level of *C. burnetii* DNA (n = 30) to be 98 genome equivalents per 500 litres of air in
8
9 areas around the farm one year after an outbreak. The lung tidal volume for a person
10
11 is approximately 0.5 litre per breath, and a farm worker taking 15 breaths per minute
12
13 over an 8 h working day would inhale 3,600 litres which would equate to a mean of
14
15 706 *C. burnetii* genome equivalents per person per working day. Mean levels of *C.*
16
17 *burnetii* DNA were 4.6-fold higher on the farm during the outbreak compared to a
18
19 year later when the air sampling was undertaken (Kersh *et al.* 2013). Thus exposures
20
21 on the farm during the outbreak through inhalation may be at the level of >3,000 *C.*
22
23 *burnetii* genome equivalents per person per working day and considerably higher than
24
25 the 532 GP_IP_ID₅₀ per person day predicted above through consumption of
26
27 unpasteurised milk. Without knowing how many bacteria there are in a GP_IP_ID₅₀ in
28
29 milk it is not possible to compare directly exposures through inhalation with those
30
31 through consumption of unpasteurised milk. In relation to exposure to *C. burnetii* due
32
33 to contact with birth products, farmers, vets and abattoir workers are most at risk.
34
35 During lambing season, in particular, exposure to such products will increase and
36
37 therefore to mitigate this risk (and that of acquiring other zoonotic pathogens),
38
39 pregnant women are advised to avoid close contact with sheep (NHS, 2014). No
40
41 information is available on the numbers of consumers that drink raw milk.
42
43
44
45
46
47
48

49 **DISCUSSION**

50 A quantitative risk assessment for transmission of *C. burnetii* to humans through milk
51
52 and milk products is not feasible at present because much of the data required are
53
54 missing. For example, while there are data from the 1950s on the number of
55
56
57
58
59
60

1
2
3 GP_IP_ID₅₀ units per ml of unpasteurised milk, there are no dose-response data to
4
5 relate how infectious a GP_IP_ID₅₀ unit is to humans through the oral route and there
6
7 are no quantitative data on survival in milk or milk products over time. *C. burnetii* is
8
9 viable in naturally infected unpasteurised milk. Thus it is well documented that guinea
10
11 pigs and mice have been experimentally infected albeit through intraperitoneal
12
13 challenge (Bell *et al.* 1949; Enright *et al.* 1957; Loftis *et al.* 2010).
14
15

16 Using the data of Enright *et al.* (1957) on levels of viable *C. burnetii* in milk of
17
18 shedding cows, it has been possible here to model the daily exposures to consumers of
19
20 unpasteurised cows' milk. The simulations demonstrate that daily exposures to viable
21
22 *C. burnetii* (in terms of GP_IP_ID₅₀) per person through unpasteurised milk are high.
23
24 This is consistent with recently published data from qPCR studies on cow BTM
25
26 samples taken in south-west England. This raises the question of how infectious *C.*
27
28 *burnetii* in milk is to humans through the oral route. Several lines of evidence are
29
30 presented here that these predicted high daily exposures through consumption of
31
32 unpasteurised milk present a relatively low risk to public health. There is little
33
34 information on the amount of milk which is consumed unpasteurised in England and
35
36 Wales, although the proportion is likely to be small. Thus, on the basis that there are
37
38 7,011 cows in unpasteurised-milk producing herds in England and Wales (data
39
40 provided by UK Food Standards Agency) and 2,864,000 dairy cows in England and
41
42 Wales (Helen Gartner, AHVLA, personal communication), it may be estimated that
43
44 just 0.24% of the total cows' milk is consumed unpasteurised.
45
46
47
48

49 Although some authors have gone as far as challenging the designation of *C. burnetii*
50
51 as a foodborne pathogen, it is concluded here that the risks to humans from *C.*
52
53 *burnetii* through consumption of unpasteurised milk and milk products (including
54
55 cheese) are not negligible but they are lower in comparison to transmission via
56
57
58
59
60

1
2
3 inhalation of aerosols from parturient products and livestock contact. This reflects the
4
5 lower risk of infection of *C. burnetii* through the oral route compared to the inhalation
6
7 route and also the much higher loadings in birth products compared to milk,
8
9 potentially giving higher exposures across the population through aerosols. There is
10
11 also some tentative evidence presented here to suggest that the pathogen is less
12
13 infectious in milk than in placentas (per DNA copy), although this needs further
14
15 substantiation.
16

17
18 *C. burnetii* has spore-like environmental stability due to the resistant SCV
19
20 morphotype which probably exists in milk and accounts for the survival of infectivity
21
22 in milk and milk products over long periods. While there are no obvious barriers in
23
24 the manufacturing of milk products, the risks may be lower for certain cheeses than
25
26 milk, particularly those cheeses with long maturation times at low pH. This is
27
28 consistent with viable *C. burnetii* rarely being detected in unpasteurised cheese
29
30 compared to unpasteurised milk and with stronger epidemiological evidence for
31
32 human cases through unpasteurised milk compared to unpasteurised cheese. A major
33
34 source of uncertainty with regard to cheese is the degree of partition of the organism
35
36 into the curds and hence the proportion which is removed with the whey. Indeed if *C.*
37
38 *burnetii* is “water-soluble”, i.e. does not partition into fat, then some 95% could be
39
40 removed with the whey, reducing the level of exposure by 20-fold. Future studies
41
42 could involve using qPCR to estimate levels of *C. burnetii* DNA in the whey and
43
44 curds.
45
46
47
48

49 50 **ACKNOWLEDGEMENTS**

51
52 We thank the UK Food Standards Agency for funding this work. We thank Andrew
53
54 Hill of AHVLA, Richard Vipond and Tim Brooks of PHE and the Specialist
55
56 Cheesemakers Association for helpful discussion.
57
58
59
60

REFERENCES

- Amara, A., Bechah, Y. and Mege, J.-L. (2012) Immune response and *Coxiella burnetii* invasion. In *Coxiella burnetii*: Recent advances and new perspectives in research of the Q fever bacterium ed. Toman, R., Heinzen, R.A., Samuel, J.E. and Mege, J.-L. *Advances in Experimental Medicine and Biology* **984**, 287-298. Dordrecht: Springer Science.
- Anon (2006). The Food Hygiene (England) Regulations 2006. Statutory Instrument 2006 No. 14. Available at: http://www.legislation.gov.uk/ukxi/2006/14/pdfs/ukxi_20060014_en.pdf (accessed 11 August 2014).
- Anon (2013) Innovate with dairy - What exactly are curds and whey? Available at: <http://www.innovatewithdairy.com/Pages/Whatexactlyarecurdsandwhey.aspx> (accessed 3 November 2014).
- ARC (2013) General concepts regarding dairy cattle. Available at: <http://www.arc.agric.za/home.asp?PID=1&ToolID=2&ItemID=1927> (accessed 11 August 2014).
- Arricau Bouvery, N., Souriau, A., Lechopier, P. and Rodolakis, A. (2003) Experimental *Coxiella burnetii* infection in pregnant goats: excretion routes. *Vet Res* **34**, 423– 433.
- Babudieri, B. and Moscovici, C. (1950) Behaviour of *Coxiella burnetii* with physical and chemical treatment. *R.C. Ist super. Sanit.* **13**, 739-748.
- Banks, J.G. (2006) Risk assessment of *L. monocytogenes* in UK retailed cheese. Final report to Food Standards Agency, Project B12006. Available at: http://www.foodbase.org.uk//admintools/reportdocuments/29-1-251_B12006_final_report.pdf (accessed 3 November 2014).

- 1
2
3 Bell, E.J., Parker, R.R. and Stoenner, H.G. (1949) Q fever: Experimental Q fever in
4
5 cattle. *Am J Public Health* **39**, 478-484.
6
7 Brouqui, P., Badiaga, S. and Raoult, D. (2004) Q fever outbreak in homeless shelter.
8
9 *Emerging Infectious Diseases* **10**, 1297-1299.
10
11 Brown, G.L., Colwell, D.C. and Hooper, W.L. (1968). An outbreak of Q fever in
12
13 Staffordshire. *J Hyg* **66**, 649-655.
14
15 Capuano, F., Mancusi, A., Casalnuovo, F., Perugini, A., Proroga, Y., Guarino, A. and
16
17 Berri, M. (2012) Real-time PCR-based detection of *Coxiella burnetii* in cheeses. *Eur*
18
19 *Food Res Technol* **235**, 1,181-1,186.
20
21 Cerf, O. and Condron, R. (2006) *Coxiella burnetii* and milk pasteurisation: an early
22
23 application of the precautionary principle? *Epidemiology and Infection* **134**, 946-951.
24
25 Combesco, D., Dumitresco, N., Zarnea, G., Saragec, A., Essrig, M., and Ionescu, H.
26
27 (1953) Experimental and epidemiological studies of pulmonary typhus (Q fever). I.
28
29 Mechanisms of transmission of infection of pulmonary typhus. *Stud Cercet*
30
31 *Inframicrobiol Microbiol Parazitol* **4** 109.
32
33 Connolly, J.H., Coyle, P.V., Adgey, A.A.J., O'Neill, H.J. and Simpson, D.M. (1990)
34
35 Clinical Q fever in Northern Ireland 1962-1989. *The Ulster Medical Journal* **59**, 137-
36
37 144.
38
39 Cutler, S.J., Bouzid, M. and Cutler, R.R. (2007) Q fever. *Journal of Infection* **54**, 313-
40
41 318.
42
43 Department of Health (2011) Whole milk (3.8% fat) consumption for all consumers
44
45 of 19 to 64 years. Available at:
46
47 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/216485](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/216485/dh_128556.pdf)
48
49 [/dh_128556.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/216485/dh_128556.pdf) (accessed 7 November 2014).
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *burnetii* in the environments of goat farms associated with a Q fever outbreak. *Appl*
4
5 *Environ Microbiol* **79**, 1697-1703.

6
7 Kingshay (2013) Kingshay dairy costings – National – published 16 April 2013.

8
9 Available at: [http://www.dairyco.org.uk/market-information/farming-data/kingshay-](http://www.dairyco.org.uk/market-information/farming-data/kingshay-dairy-costings/kingshay-dairy-costings-national/)
10
11 [dairy-costings/kingshay-dairy-costings-national/](http://www.dairyco.org.uk/market-information/farming-data/kingshay-dairy-costings/kingshay-dairy-costings-national/) (accessed 20 March 2014).

12
13
14 Klee, S.R., Tyczka, J., Ellerbrok, H., Franz, T., Linke, S., Baljer, G. and Appel, B.
15
16 (2006) Highly sensitive real-time PCR for specific detection and quantification of
17
18 *Coxiella burnetii*. *BMC Microbiology* **6**, 2.

19
20
21 Krumbiegel, E.R. and Wisniewski, H.J. (1970) Q fever in Milwaukee. Consumption
22
23 of infected raw milk by human volunteers. *Arch Environ Health* **21**, 63-65.

24
25 Lambton, S.L., Smith, R., Gillard, K. and Pritchard, G.C. (*unpublished*). Survey of
26
27 sheep and goats sera for Q-fever.

28
29
30 Langley, J.M., Marrie, T.J., Covert, A., Waag, D.M. and Williams, J.C. (1988) Poker
31
32 players' pneumonia. An urban outbreak of Q fever following exposure to a parturient
33
34 cat. *New Eng J Med* **319**, 354-356.

35
36
37 Loftis, A.D., Priestley, R.A. and Massung, R.F. (2010). Detection of *Coxiella burnetii*
38
39 in commercially available raw milk from the United States. *Foodborne Pathog and*
40
41 *Dis* **7**, 1453–56.

42
43 Marmion, B.P. and Harvey, M.S. (1956). The varying epidemiology of 'Q' fever in
44
45 the south east region of Great Britain. I. In an urban area. *J Hyg* **54**, 533-546.

46
47 Maurin, M. and Raoult, D. (1999). Q fever. *Clin Microbiol Rev* **12**, 518-553.

48
49 McCaughey, C., Murray, L.J., McKenna, J.P., Menzies, F.D., McCullough, S.J.,

50
51 O'Neill, H.J.O., Wyatt, D.E., Cardwell, C.R. and Coyle, P.V. (2010) *Coxiella burnetii*
52
53 (Q fever) seroprevalence in cattle. *Epidemiology and Infection* **138**, 21-27.
54
55
56
57
58
59
60

1
2
3 Minnick, M.F. and Raghavan, R. (2011) Genetics of *Coxiella burnetii*: on the path of
4
5 speciation. *Future Microbiology* **6**, 1,297-1,314.
6

7 Minnick, M.F. and Reghavan, R. (2012) Development biology of *Coxiella burnetii*. In
8
9 *Coxiella burnetii*: Recent advances and new perspectives in research of the Q fever
10
11 bacterium ed. Toman, R., Heinzen, R.A., Samuel, J.E. and Mege, J.-L. *Advances in*
12
13 *Experimental Medicine and Biology* **984**, 231-248. Dordrecht: Springer Science.
14

15
16 Omsland, A. and Heinzen, R.A. (2011) Life on the outside: The rescue of *Coxiella*
17
18 *burnetii* from its host cell. *Annual Review of Microbiology* **65**, 111-128.
19

20
21 Oyston, P.C.F. and Davies, C. (2011) Q fever: the neglected biothreat agent. *J Med*
22
23 *Microbiol* **60**, 9-21.
24

25 NHS (2014) Why should pregnant women avoid sheep during the lambing season?

26
27 Available at:

28
29 <http://www.nhs.uk/chq/pages/934.aspx?CategoryID=54&SubCategoryID=131>

30
31 (accessed 11 August 2014).
32
33

34 Paape, M.J., Bannerman, D.D., Zhao, X. and Lee, J.W. (2003) The bovine neutrophil:
35
36 Structure and function in blood and milk. *Vet Res* **34**, 597-627.
37

38 Paiba, G.A., Green, L.E., Lloyd, G., Patel, D. and Morgan, K.L. (1999). Prevalence
39
40 of antibodies to *Coxiella burnetii* (Q fever) in bulk tank milk in England and Wales.
41
42 *Veterinary Record* **144**, 519-522.
43

44
45 Pebody, R.G., Wall, P.G., Ryan, M.J. and Fairley, C. (1996). Epidemiological
46
47 features of *Coxiella burnetii* infection in England and Wales: 1984 to 1994.
48

49
50 *Communicable Disease Report* **6**, R128-R132.
51

52 Raoult, D., Tissot-Dupont, H., Foucault, C., Gouvernet, J., Fournier, P.E., Bernit, E.,

53
54 Stein, A., Nesri, M. *et al.* (2000). Q fever 1985-1998: Clinical and epidemiological
55
56 features of 1,383 infections. *Medicine* **79**, 109-123.
57
58
59
60

- 1
2
3 Rodolakis, A., Berri, M., He'chard, C., Caudron, C., Souriau, A., Bodier,
4
5 C.C., Blanchard, B., Camuset, P. *et al.* (2007). Comparison of *Coxiella burnetii*
6
7 shedding in milk of dairy bovine, caprine, and ovine herds. *J Dairy Sci* **90**, 5352–
8
9 5360.
- 10
11 Roest, H.-J., van Gelderen, B., Dinkla, A., Frangoulidis, D., van Zijderveld, F., Rebel,
12
13 J. and van Keulen, L. (2012) Q Fever in Pregnant Goats: Pathogenesis and Excretion
14
15 of *Coxiella burnetii*. *PLoS ONE* **7**, e48949. doi:10.1371/journal.pone.0048949
- 16
17 Rousset, E., Berri, M., Durand, B., Dufour, P., Prigent, M., Delcroix, T., Touratier, A.
18
19 and Rodolakis, A. (2009) *Coxiella burnetii* shedding routes and antibody response
20
21 after outbreaks of Q fever-induced abortion in dairy goat herds. *Appl Environ*
22
23 *Microbiol* **75**, 428-433.
- 24
25
26
27 Sipka, M. (1958) Survival of *Rickettsia burneti* in cheese. *Veterinarski Glasnik* **12**, 9-
28
29 12.
- 30
31
32 Schimmer, B., Lutikholt, S., Hautvast, J.L.A., Graat, E.A.M., Vellema, P. and van
33
34 Duynhoven, Y.T.H.P. (2011) Seroprevalence and risk factors of Q fever in goats on
35
36 commercial dairy goat farms in the Netherlands, 2009-2010. *BMC Veterinary*
37
38 *Research* **7**, 81.
- 39
40
41 Signs, K.A., Stobierski, M.G. and Gandhi, T.N. (2012) Q fever cluster among raw
42
43 milk drinkers in Michigan, 2011. *Clinical Infectious Disease* **55**, 1387-9.
- 44
45 Thomas, D.R., Treweek, L., Salmon, R.L., Kench, S.M., Coleman, T.J., Meadows, D.,
46
47 Morgan-Capner, P. and Caul, E.O. (1995) The risk of acquiring Q fever on farms: a
48
49 seroepidemiological study. *Occup Environ Med* **52**, 644-647.
- 50
51
52 Tilburg, J. H. C., Roest, H. J, I. J., Nabuurs-Franssen, M. H., Horrevorts, A. M.,
53
54 Klaassen, C. H. W. (2012) Genotyping Reveals the Presence of a Predominant
55
56
57
58
59
60

1
2
3 Genotype of *Coxiella burnetii* in Consumer Milk Products. *Journal of Clinical*
4
5 *Microbiology* **50**, 2156-2158.

6
7 Tissot-Dupont, H., Amadei, M.A., Nezri, M. and Raoult, D. (2004) Wind in
8
9 November, Q fever in December. *Emerg Infect Dis* **10**, 1264–1269.

10
11 Valergakis, G.E., Russell, C., Grogono-Thomas, R., Bradley, A.J. and Eisler, M.C.
12
13 (2012). *Coxiella burnetii* in bulk tank milk of dairy cattle in south-west England.

14
15 *Veterinary Record* **171**, 156.

16
17 Van Woerden, H.C., Mason, B.W., Nehaul, L.K., Smith, R., Salmon, R.L., Healy, B.,
18
19

20
21 Valappil, M., Westmoreland, D. *et al.* (2004) Q fever outbreak in industrial setting.

22
23 *Emerging Infectious Diseases* **10**, 1282-1289.

24
25 Welsh, H.H., Lennette, E.H., Abinanti, F.R. and Winn, J.F. (1951) *Public Health Rep*
26
27 *Wash* **66**, 1473-1477.

28
29 Wilson, L.E., Couper, S., Prempeh, H., Young, D., Pllock, K.G.J., Stewart, W.C.,
30
31

32
33 Browning, L.M. and Donaghy, M. (2010) Investigation of a Q fever outbreak in a
34
35 Scottish co-located slaughterhouse and cutting plant. *Zoonoses and Public Health* **57**,
36
37 493-498.

38
39 World's Healthiest Foods (2013). Available at:

40
41 <http://www.whfoods.com/genpage.php?tname=george&dbid=100> (accessed 21 June
42
43 2013).

44
45 Zubkova, R.I. (1957) Survival of *Rickettsia burneti* in milk and milk products.

46
47 *Journal of Microbiology, Epidemiology and Immunobiology* **28**, 42-46.

Table 1: Summary of data used for estimating probability and levels of *C. burnetii* in BTM (per herd) for cattle in the baseline model.

Description	Parameter	Summary of data probability distribution	Reference
Number of dairy cows per herd	H	Used empirical data for 81 cattle herds in England and Wales supplying unpasteurised milk.	Provided by UK Food Standards Agency
Probability herd is positive	p_{Herd}	$\begin{cases} 0.318 \text{ if } H < 50 \\ 0.600 \text{ if } 50 \leq H \leq 100 \\ 0.781 \text{ if } H > 100 \end{cases}$	McCaughey <i>et al.</i> (2010)
Probability animal is positive given herd is positive	$p_{\text{Within_herd}}$	$\begin{cases} 0.034 \text{ if } H < 50 \\ 0.102 \text{ if } 50 \leq H \leq 100 \\ 0.125 \text{ if } H > 100 \end{cases}$	McCaughey <i>et al.</i> (2010)
Probability animal is lactating	$p_{\text{Lactating}}$	Pert (265, Uniform (300,305); 340)/365	ARC (2013).
Probability animal is shedding <i>C. burnetii</i> in milk given animal is lactating and infected	p_{Shedding}	22 of 72 infected cows (0.3055)	Guatteo <i>et al.</i> (2012)
Volume of milk (per animal per day)	V_i	Normal (25.6, 1.263) (litre)	Kingshay (2013)
<i>Coxiella burnetii</i> concentration in milk (Shedders)	C_{ml}	Guinea pig intraperitoneal ID ₅₀ per ml distributed as $0.5 \times 10^{\text{Normal}(1.333, 1.0847)}$	Enright <i>et al.</i> (1957)
Cumulative milk consumption per person per day	$M_{\text{Litre/Day}}$	0.127 (litre per person per day)	Department of Health (2011)




Figure 1: Schematic diagram for the probability of exposure and levels of *C. burnetii* per person per day through consumption of unpasteurised milk. The model outputs are in boxes.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2: Simulated GP_IP_ID₅₀s of *Coxiella burnetii* per ml of unpasteurised BTM milk plotted on a log scale for comparison with qPCR data of Valergakis *et al.* (2012).