

# The Connection of the Degasperis–Procesi Equation with the Vakhnenko Equation

Vyacheslav O. VAKHNENKO <sup>†</sup> and John PARKES <sup>‡</sup>

<sup>†</sup> *Institute for Geophysics, 32 Palladin Ave., 03680 Kyiv, Ukraine*

E-mail: *vakhnenko@bitp.kiev.ua*

<sup>‡</sup> *University of Strathclyde, Richmond St., Glasgow G1 1XH, UK*

E-mail: *ejp@maths.strath.ac.uk*

Travelling-wave solutions of the Degasperis–Procesi equation (DPE) are investigated. The solutions are characterized by two parameters. Hump-like, loop-like and coshoidal periodic-wave solutions are found; hump-like, loop-like and peakon solitary-wave solutions are obtained as well. Hone and Wang showed a connection between the DPE and the Vakhnenko equation (VE). Comparing the solutions of the DPE and the VE, we observe that, for both equations at interaction of waves, there are three kinds of phaseshift that depend on the ratio of wave amplitudes. In particular, there is a case when two interacted waves have phaseshifts in the positive direction.

## 1 Introduction

This report deals with the Degasperis–Procesi equation (DPE) [1–3]

$$u_t - u_{txx} + 4uu_x = 3u_xu_{xx} + uu_{xxx} \quad (1)$$

which is contained in the family of equations [1]

$$u_t - u_{txx} + (b + 1)uu_x = bu_xu_{xx} + uu_{xxx}, \quad (2)$$

(parameter  $b$  is constant). When  $b = 2$ , equation (2) reduces to the well-known Camassa–Holm equation (CH) [4, 5]

$$u_t - u_{txx} + 3uu_x = 2u_xu_{xx} + uu_{xxx}. \quad (3)$$


Mikhailov and Novikov developed a powerful extension of the symmetry classification method [6], and applying this to the equation (2) they found that only the cases  $b = 2, 3$  could possess infinitely many commuting symmetries, and so only these two cases are integrable. In a recent paper Degasperis, Holm and Hone [2] constructed a Lax pair of the equation (1), and hence proved the integrability of the Degasperis–Procesi equation. Unfortunately, there are no solutions of the direct problem and the inverse problem of the spectral equation from this Lax pair.

Hone and Wang [7] have shown the connection of the DPE (1) with the integrable Vakhnenko equation (VE) [8–10]

$$(u_t + uu_x)_x + u = 0. \quad (4)$$

Indeed, the transformation  $x \rightarrow \varepsilon x - t/3\varepsilon$ ,  $t \rightarrow \varepsilon t$ ,  $u \rightarrow u - 1/3\varepsilon^2$  reduces the DPE to the VE when  $\varepsilon \rightarrow 0$

$$[(u_t + uu_x)_x + u]_x = \varepsilon^2(u_t + 4uu_x). \quad (5)$$



**Figure 1.** The polynomial  $f(z)$ . The interval of integration is between the roots  $z_2$  and  $z_3$ .

Recently, we have investigated equation (4) by direct integration [8,9], by Hirota's method [11] as well as by the inverse scattering transform method [10]. In the context of this paper, it is of interest that the VE has two families of travelling-wave solutions (see Figs. 1, 2 in [8]); in particular, there is a soliton solution in loop-like form.

Comparing the properties of the solutions of the DPE (1) and the VE (4), we observe that for both equations under interaction of waves, there are three kinds of phaseshift that depend on the ratio of wave amplitudes [2, 12, 13]. In particular, there is a case when two interacted waves have phaseshifts in the positive direction.

However, it engages our attention that soliton solutions have not been observed for the DPE recently. If the VE has a loop-like solution, it follows that the DPE should admit a loop-like solution. In this paper we investigate the travelling-wave solutions of the DPE. The solutions are characterized by two parameters. Hump-like, loop-like and coshoidal periodic-wave solutions are found; hump-like, loop-like and peakon solitary-wave solutions are obtained as well.

## 2 Travelling-wave solutions

We consider the DPE (1). Restricting our attention to travelling waves, we introduce new variables

$$z = (u - v)/|v|, \quad \eta = x - vt - x_0, \quad \tau = |v|t, \quad (6)$$

where  $v$  and  $x_0$  are arbitrary constants with  $v \neq 0$ . In this case the initial equation is reduced to the ODE

$$(zz_\eta)_{\eta\eta} = (4z + 3c)z_\eta \quad \text{with} \quad c := v/|v| = \pm 1. \quad (7)$$

After two integrations we have

$$(zz_\eta)^2 = f(z) \quad (8)$$

with

$$f(z) = z^4 + 2cz^3 + Az^2 + B = (z - z_1)(z - z_2)(z - z_3)(z - z_4),$$

where  $f(z)$  is a polynomial of fourth order and is shown in Fig. 1.




Figure 2. Four possible cases for the polynomial  $f(z)$ .

Analyzing equation (8), we observe that the roots of the polynomial  $f(z)$  are real and the interval of integration is between the roots  $z_2$  and  $z_3$ . We can write two forms of solution. The first form of solution is

$$z = \frac{z_2 - z_1 \operatorname{sn}^2(w|m)}{1 - n \operatorname{sn}^2(w|m)} \quad \text{with} \quad n = \frac{z_3 - z_2}{z_3 - z_1}, \tag{9}$$

where

$$w = p\zeta, \quad p = \frac{1}{2}\sqrt{(z_4 - z_2)(z_3 - z_1)}, \quad m = \frac{(z_3 - z_2)(z_4 - z_1)}{(z_4 - z_2)(z_3 - z_1)}$$

and

$$\eta = [wz_1 + (z_2 - z_1)\Pi(n; w|m)]/p. \tag{10}$$

Here  $\operatorname{sn}(w|m)$  is a Jacobian elliptic function and  $\Pi(n; w|m)$  is the elliptic integral of the third kind. The second form of solution is

$$z = \frac{z_3 - z_4 \operatorname{sn}^2(w|m)}{1 - n \operatorname{sn}^2(w|m)} \quad \text{with} \quad n = \frac{z_3 - z_2}{z_4 - z_2} \tag{11}$$

and

$$\eta = [wz_4 + (z_4 - z_3)\Pi(n; w|m)]/p. \tag{12}$$

Hence the solution is given in parametric form with  $w$  as a parameter. The solutions are characterized by the two parameters  $A$  and  $B$ , or equivalently  $A$  and  $m$ .


### 3 Classification of solutions

The equation (7) is invariant under the transformation  $z \rightarrow -z, c \rightarrow -c$ ; this corresponds to the transformation  $u \rightarrow -u, v \rightarrow -v$ . Hence we need to consider only the case  $c = 1$  (i.e.  $v > 0$ ).

Four cases are possible for the polynomial  $f(z)$  corresponding to different ranges of values of  $A$  (see Fig. 2). Examples of the corresponding solutions are illustrated in Fig. 3. In the first column are periodic solutions with  $m \neq 1$ . In the second column are the corresponding limiting cases with  $m = 1$ . These are solitary wave solutions, except for the figure in the second line which is a periodic peakon solution.

### 4 Conclusion

Travelling-wave solutions of the Degasperis–Procesi equation are investigated. The solutions are characterized by two parameters. Hump-like, loop-like and coshoidal periodic-wave solutions are found; hump-like, loop-like and peakon solitary-wave solutions are obtained as well.



**Figure 3.** Travelling-wave solutions.

## Acknowledgements

This research was supported in part by STCU, Project N 1747.

- [1] Degasperis A. and Procesi M., Asymptotic integrability, in *Symmetry and Perturbation Theory*, Editors A. Degasperis and G. Gaeta, World Scientific, 1999, 23–37.
- [2] Degasperis A., Holm D.D. and Hone A.N.W., A new integrable equation with peakon solutions, *Theor. Math. Phys.*, 2002, V.133, 1461–1472.
- [3] Degasperis A., Holm D.D. and Hone A.N.W., Integrable and non-integrable equations with peakons, in *Non-linear Physics: Theory and Experiment II (2002, Gallipoli)*, Editors M.J. Ablowitz, M. Boiti, F. Pempinelli and B. Prinari, Singapore, World Scientific, 2003, 37–43; nlin.SI/0209008.
- [4] Camassa R. and Holm D.D., An integrable shallow water equation with peaked solitons, *Phys. Rev. Lett.*, 1993, V.71, 1661–1664.
- [5] Fuchssteiner B. and Fokas A.S., Symplectic structures, their Bäcklund transformations and hereditary symmetries, *Physica D*, 1981, V.4, 47–66.

- 
- [6] Mikhailov A.V. and Novikov V.S., Perturbative symmetry approach, *J. Phys. A: Math. Gen.*, 2002, V.35, N 22, 4775–4790.
  - [7] Hone A.N.W. and Wang J.P., Prolongation algebras and Hamiltonian operators for peakon equations, *Inverse Problems*, 2003, V.19, 129–145.
  - [8] Vakhnenko V.A., Solitons in a nonlinear model medium, *J. Phys. A: Math. Gen.*, 1992, V.25, 4181–4187.
  - [9] Parkes E.J., The stability of solutions of Vakhnenko’s equation, *J. Phys. A: Math. Gen.*, 1993, V.26, 6469–6475.
  - [10] Vakhnenko V.O. and Parkes E.J., The calculation of multi-soliton solutions of the Vakhnenko equation by the inverse scattering method, *Chaos Solitons and Fractals*, 2002, V.13, 1819–1826.
  - [11] Morrison A.J., Parkes E.J. and Vakhnenko V.O., The  $N$  loop soliton solution of the Vakhnenko equation, *Nonlinearity*, 1999, V.12, 1427–1437.
  - [12] Vakhnenko V.O. and Parkes E.J., A novel nonlinear evolution equation integrable by the inverse scattering method, in Proceedings of Fourth International Conference “Symmetry in Nonlinear Mathematical Physics” (9–15 July, 2001, Kyiv), Editors A.G. Nikitin, V.M. Boyko and R.O. Popovych, Kyiv, Institute of Mathematics, 2002, V.43, Part 1, 384–391.
  - [13] Vakhnenko V.O. and Parkes E.J., The two loop soliton of the Vakhnenko equation, *Nonlinearity*, 1998, V.11, 1457–1464.