

# A technique to record the sedentary to walk movement during free living mobility: A comparison of healthy and stroke populations

Kerr<sup>1</sup>, Rafferty<sup>2</sup>, Hollands<sup>3</sup>, Barber<sup>4</sup>, Granat<sup>3</sup>

1 Biomedical Engineering, University of Strathclyde, UK.

2 Institute for Applied Health Research, Glasgow Caledonian University, UK.

3 School of Health Sciences, University of Salford, UK.

4 Stroke MCN, NHS Lanarkshire, UK.

## Corresponding author

Dr Andy Kerr

Department of Biomedical Engineering

University of Strathclyde

Wolfson Centre

106 Rottenrow East

Glasgow

G4 0NW

Email: [a.kerr@strath.ac.uk](mailto:a.kerr@strath.ac.uk)

T: 0141 548 2855

## Introduction

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Everyday life involves frequent transitions between postures and movements, such as sitting to standing and standing to walking. Variability within and between individuals performing these transitions is a hallmark of normal movement and a consequence of the abundance of motor solutions available to healthy individuals(1). In general, and irrespective of environmental and task specific factors, this motor flexibility allows healthy individuals to perform manifold daily activities without hesitation. However, limited motor flexibility, resulting from impairments observed in conditions like stroke, Parkinson’s disease and, more generally, age related frailty, can result in stereotypical, slow and hesitant transitions(2-4). To date movement transitions, e.g. sit-to-stand and sit-to-walk have been studied under controlled laboratory conditions (3-5) employing detailed biomechanical and muscle activation measurement techniques, while this provides important understanding of movement at the body functions and structure level (6) it provides only limited understanding of everyday movement activity. Studying these transitions during everyday life could help resolve problems such as the recovery of community mobility after stroke (7). Activity monitors can classify postures (8) and measure the time taken to change postures. Taken together these parameters allow the reporting of transitions in movement such as sit/lying (sedentary) to walk, during free-living.

The aim of this study was to test a new method for quantifying a sedentary to walk transition using the time period between a sedentary posture (sitting/Lying) and a bout of walking with populations of differing levels of mobility.

## Methods

### Participants

Data were extracted from two physical activity studies, providing two contrasting populations:

1) Stroke patients (n=34), including 31 infarcts and 3 haemorrhagic, recently (<14days) discharged from being an in-patient (median length of stay 44 days (IQR 18 to 62)) but still receiving rehabilitation input as part of their early supported discharge (ESD). Eleven individuals were living alone .They were aged  $68.9\pm 11.8$  years, height  $1.67\pm 0.2$ m and weight  $73.1\pm 18.6$ kg, 18 were male and there were variable levels of mobility (Modified Rivermead Mobility Index, median 34, IQR 30-37). The original study (UKCRN15472) was approved by the West of Scotland Ethics committee (13/WS/0150).

2) An age matched control group (n=30) was recruited consecutively from the local community. They were aged  $66.8\pm 10.5$  years and included 18 males. Ethical approval was granted by the Glasgow Caledonian University, School of Health and Life Sciences, Ethics Committee.

Data were collected from all participants in the same manner, an activity monitor (dimensions 45mm, 25mm, 5mm, and <15g in weight) consisting of a triaxial accelerometer (activPAL3, PAL Technologies Ltd, Glasgow, UK) was attached, using Tegaderm™ (3M, Neuss, Germany), to the anterior aspect of the participant's thigh (unaffected side for stroke patients and right side for controls). All participants were asked to continue their everyday activities as normal. After a minimum of 48 hours of continuous recording during the working week (Monday – Friday) the activity monitor was removed and the stored data

1 downloaded for processing. The average monitoring period was 92.26 hours (SD 40.61) for  
2 the stroke group and 164.80 hours (SD 12.80) for the control group.  
3

#### 4 5 Data processing 6

7  
8  
9 The activPAL3 samples data at 20Hz, these data are then classified into events using  
10 proprietary algorithms. The use of a single sensor limits the system's ability to differentiate  
11 between a lying and seated position, therefore events were identified as sedentary (either  
12 sitting or lying), standing and walking. Consecutive stride events were combined to give  
13 walking events. Each event has a start time and duration associated with it. The output from  
14 the device has been validated for classification of sedentary, standing, and walking activities  
15 in a range of populations (9).  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

26  
27 The sedentary to walk (STW) transition time was then calculated as follows:  
28

29  
30 Start time of the walking event – end time of the previous sedentary event.  
31  
32

33  
34 Based on this calculation four different categories of STW transition were determined using  
35 values gathered from laboratory studies (3, 10, 11).  
36  
37

38  
39  
40 1) Fluent STW: walking starts within 2s of a change in posture from sedentary. This time  
41 frame was based on healthy older adults being able to complete an entire (initiation to end)  
42 sit-to-walk transition within 1.8-2.3s (2, 3, 11). Two seconds was therefore considered a  
43 reasonable, maximum, time delay to consider it a single fluent movement.  
44  
45  
46  
47  
48  
49

50  
51 2) Hesitant STW: the walking event starts between 2s and 10s after the end of a sedentary  
52 event. Adults at risk of falling and stroke survivors can perform the whole STW transition, on  
53 average, within 10s (95% CI), including pauses in the movement (10, 11). Ten seconds was  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 therefore considered a reasonable maximum time delay, to consider it a single, if hesitant,  
2 movement.  
3

4  
5  
6 3) Separated STW: Walking occurring after a sedentary event with a substantial delay (>10s).

7  
8 This value was selected to be reflective of a disconnected sedentary to walk movement  
9 based on a hesitant STW being within a maximum of 10s.  
10

11  
12  
13 With a further classification of:

14  
15  
16 4) Sedentary to stand to sedentary (STSTS): There was a change from sedentary recorded  
17 without a subsequent bout of walking before a return to sedentary.  
18  
19

20  
21  
22 See figure 1 for illustration of these transitions using raw accelerometer data.  
23

24  
25  
26 Insert figure 1  
27

28  
29  
30 Figure 1: Illustration of transitions using raw accelerometer data.  
31

32  
33  
34 To explore the validity of these definitions the whole dataset (stroke and healthy age  
35 matched controls) was separated into discrete time bins (0-2, 2-4, 4-6, 6-8, 8-10, 10-15, 15-  
36  
37 20, ... >40) and plotted against the percentage of transitions for that group.  
38  
39

40  
41  
42 Statistical analysis  
43

44  
45  
46 Statistical differences for the percentages of these transitions between the groups were  
47 tested using the Mann-Whitney Test, and an alpha level of 0.05 was set for significance.  
48  
49

50  
51 **Dependence between the Modified Rivermead Mobility Index and the physical activity**

52 **monitor was explored with Spearman's rank correlation, all statistical tests were carried out**  
53  
54  
55 **with Minitab (Penn, USA).**  
56  
57

58  
59  
60 Results  
61  
62  
63  
64  
65

1 Walking followed a transition from sedentary on 91.8% of occasions in the control group  
2 compared to 68.0% (SD 11.9) in the stroke group. Only a median of 9.14% (IQR, 4.50-17.46)  
3 of the transitions performed by the stroke group per day were fluent (<2s delay between  
4 standing up and walking) compared to a mean of 43.96% for the controls, see table 1 and  
5 figure 1. In contrast 33.9±19.5% of transitions in the stroke group were categorised as STSTS  
6 compared to just 8.20±5.42% in the control group. These differences were statistically  
7 significant for both the fluent (p<0.001) and the STSTS (p<0.001). There were no significant  
8 differences between the groups for hesitant and separated transitions. Hesitant transitions  
9 accounted for 22.87±6.54% and 23.94±13.56% for controls and stroke respectively (p=0.69)  
10 while separated transitions accounted for 25.97±6.18% and 30.12±13.00% for controls and  
11 stroke respectively (p=0.11). There was a good positive correlation (Spearman rho 0.55,  
12 p=0.01) between the Modified Rivermead Mobility Index and percentage of daily fluent STW  
13 transitions, indicating stroke survivors with better mobility performed a greater percentage  
14 of fluent STW transitions.

15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37 Insert Figure 2

38  
39  
40  
41 Figure 2: Sedentary to walk transition categories expressed as a percentage of total and  
42 separated into time bins.  
43  
44

45  
46 Insert table 1

47  
48  
49 Table 1: Average (variance) number of daily transitions according to group with percentages  
50 of total  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Discussion

We present a new method for measuring and categorising transitions between sedentary postures and walking during everyday living. Using transition time we categorised; 1) a fluent sedentary to walk transition (<2s), 2) a hesitant transition (2-10s), 3) a separated transition (>10s, but walking does occur) and finally 4) a sedentary to stand to sedentary (STSTS) transition (i.e. no walking occurs). When applied to two populations, with (stroke) and without (healthy control) mobility impairment, this categorisation technique revealed significant differences, illustrating its potential value to mobility screening and rehabilitation research.

Using this technique, for example, it is evident that the primary reason for standing up in everyday life is to walk; 92% of the sedentary-to-stand transitions were followed by walking in healthy individuals. This finding supports the use of mobility tests that combine sit to stand and walking (12), as a better reflection of real world mobility. The advantage of the presented technique is that it can measure an individual's actual mobility at home over long periods of time, improving the measurement validity. Using the transition definitions a more detailed profile of an individual's mobility can be gained; fluent transitions, for example were much more common (43%) in the healthy older adults compared to the stroke population (9%), and better scores on the Modified Rivermead Mobility Index for the stroke patients were reasonably well correlated ( $r=0.55$ ) with the percentage of daily fluent STW transitions. These findings may be useful in detecting subtle changes in mobility.

## Limitations

Limited clinical information on the stroke sample prevented a more robust analysis of factors such as stroke severity, the use of assistance and psychological factors such as fear

1 of falling. The data were all derived from single site acceleration signals and the accuracy of  
2 the classification algorithms may be at risk with very slow moving individuals such as stroke  
3 survivors. Finally we recognise that in the absence of definitive free-living cut-off values the  
4 presented values of less than 2s, between 2 and 10s, and greater than 10s, whilst based on  
5 literature, may need to be adjusted in future as more data becomes available. To facilitate  
6 development of this technique we have presented the percentage data for 2 second bins  
7 (figure 2) to allow future researchers to explore different cut –off points.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

## 19 Conclusion

20  
21  
22 A novel technique for classifying movement transitions in everyday life found statistically  
23 significant differences in the type of transition (fluent, hesitant and separated) performed  
24 by groups with differing levels of mobility, creating opportunities to further understand  
25 community mobility.  
26  
27  
28  
29  
30  
31

## 32 Acknowledgements

33  
34  
35 The study was supported by the Scottish Stroke Research Network who assisted in  
36 participant screening and recruitment and Chest Heart and Stroke Scotland who provided  
37 funds for research equipment and associated expenses  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



## References

1. Stergiou N, Decker LM. Human movement variability, nonlinear dynamics, and pathology: Is there a connection? *Human Movement Science*. 2011;30(5):869-88.
2. Malouin F, McFadyen B, Dion L, Richards CL. A fluidity scale for evaluating the motor strategy of the rise-to-walk task after stroke. *Clinical Rehabilitation*. 2003 Sep;17(6):674-84.
3. Frykberg GE, Aberg AC, Halvorsen K, Borg J, Hirschfeld H. Temporal coordination of the sit-to-walk task in subjects with stroke and in controls. *Arch Phys Med Rehabil*. 2009 Jun;90(6):1009-17.
4. Buckley TA, Pitsikoulis C, Hass CJ. Dynamic postural stability during sit-to-walk transitions in Parkinson disease patients. *Mov Disord*. 2008 Jul 15;23(9):1274-80.
5. Kang GE, Gross MM. Emotional influences on sit-to-walk in healthy young adults. *Human Movement Science*. 2015;40:341-51.
6. World Health Organisation. International Classification of Functioning, Disability and Health (ICF). 2016 [cited 2016 14/05/2016]; Website].
7. Hyndman D, Ashburn A, Stack E. Fall events among people with stroke living in the community: Circumstances of falls and characteristics of fallers. *Arch Phys Med Rehabil*. 2002;83(2):165-70.
8. Kim Y, Barry VW, Kang M. Validation of the ActiGraph GT3X and activPAL Accelerometers for the Assessment of Sedentary Behavior. *Measurement in Physical Education and Exercise Science*. 2015 2015/07/03;19(3):125-37.
9. Grant PM, Dall PM, Mitchell SL, Granat MH. Activity monitor accuracy in measuring step number and cadence in community-dwelling older adults. *Journal of Aging and Physical Activity*. 2008;16:3.
10. Dion L, Malouin F, McFadyen B, Richards CL. Assessing mobility and locomotor coordination after stroke with the rise-to-walk task. *NEUROREHABIL NEURAL REPAIR*. 2003 Jun;17(2):83-92.
11. Kerr A, Rafferty D, Kerr KM, Durward B. Timing phases of the sit-to-walk movement: validity of a clinical test. *Gait Posture*. 2007 Jun;26(1):11-6.
12. Vernon S, Paterson K, Bower K, McGinley J, Miller K, Pua Y-H, et al. Quantifying Individual Components of the Timed Up and Go Using the Kinect in People Living With Stroke. *Neurorehabilitation and Neural Repair*. 2015 January 1, 2015;29(1):48-53.
13. SIGN. Management of patients with stroke: rehabilitation, prevention and management of complications, and discharge planning. In: Scotland NQI, editor. Edinburgh: SIGN; 2010.

Table 1: Average (variance) number of daily transitions according to group with percentages of total

	All (N=64) Mean (SD)	Stroke (n=34) Mean (SD)	Controls (n=30) Mean (SD)	Comparison p-value
Fluent	14 (11)	4 (2.14-11.56)*	22 (9)	<0.001
STW	26.54% (18.92)	9.14% (4.50-17.46)*	42.96% (12.58)	<0.001
Hesitant	9 (14)	14 (18)	5 (6)	0.004
STW	23.44% (10.78)	23.94% (13.56)	22.87% (6.54)	0.595
Separated	18 (16)	8 (8)	28 (15)	<0.001
STW	28.17% (10.51)	30.12% (13.00)	25.97% (6.18)	0.74
Sedentary to stand to Sedentary	8 (16) 21.85% (19.50)	14 (20) 33.88% (19.54)	2 (6) 8.20% (5.42)	<0.001 <0.001

\*median and IQR range reported as data were not normally distributed

7. Figure1

[Click here to download high resolution image](#)





Figure 2  
[Click here to download high resolution image](#)



### Research highlights

- A novel method for measuring movement transitions during everyday life is presented.
- Healthy participants walk after standing up on almost every occasion (92%).
- Typically this is a single action sedentary to walk movement.
- Stroke survivors walk less frequently after standing up (66%).
- Typically walking from a sedentary position is performed in a hesitant and/or separated pattern in stroke survivors.