

Associations of Vigorous-intensity Physical Activity with Biomarkers in Youth

Short title: Vigorous Physical Activity, Biomarkers, Youth

Justin B. Moore, PhD, MS,¹ Michael W. Beets, PhD, MPH, MEd,² Keith Brazendale, MS,² Steven N. Blair, PED,^{2,3} Russell R. Pate, PhD,² Lars B. Andersen, PhD,⁴ Sigmund A. Anderssen, PhD,⁵ Anders Grøntved, PhD, MPH, MSc,⁴ Pedro C. Hallal, PhD,⁶ Katarzyna Kordas, PhD,⁷ Susi Kriemler, PhD,⁸ John J. Reilly, PhD,⁹ Luis B. Sardinha, PhD,¹⁰

Affiliations: ¹Wake Forest School of Medicine, Department of Family and Community Medicine, Winston-Salem, North Carolina, US; ² University of South Carolina, Arnold School of Public Health, Department of Exercise Science, Columbia, South Carolina, US; ³University of South Carolina, Arnold School of Public Health, Department of Epidemiology and Biostatistics, Columbia, South Carolina, US; ⁴University of Southern Denmark, Department of Sport Science and Clinical Biomechanics, Odense, Denmark; ⁵Norwegian School of Sport Science, Department of Sport Medicine, Oslo, Norway; ⁶Federal University of Pelotas, Pelotas, Brazil; ⁷University of Bristol, School of Social and Community Medicine, Bristol, UK; ⁸University of Zürich, Epidemiology, Biostatistics and Public Health Institute, Zürich, Switzerland; ⁹University of Strathclyde, Physical Activity for Health Group, School of Psychological Sciences and Health, Glasgow, UK; ¹⁰Technical University of Lisbon, Faculty of Human Movement, Lisbon, Portugal; On behalf of the International Children's Accelerometry Database Collaborators

Address correspondence to: Justin B. Moore, PhD, MS, FACSM, Associate Professor
Department of Family & Community Medicine, Wake Forest School of Medicine, Wake Forest Baptist Medical Center. Winston-Salem, NC 27157, jusmoore@wakehealth.edu, 336.716.3702.

Abbreviations: CPM – Counts per minute; HDL – High Density Lipoprotein; ICAD – International Children's Accelerometry Database; LDL – Low Density Lipoprotein; MPA – Moderate Physical Activity; MVPA – Moderate-to-vigorous Physical Activity; VPA – Vigorous Physical Activity; WC – Waist Circumference.

ABSTRACT

Introduction: Physical activity (PA) conveys known cardiometabolic benefits to youth, but the contribution of vigorous-intensity PA (VPA) to these benefits is unknown. Therefore, we sought to determine, a) the associations between VPA and cardiometabolic biomarkers independent of moderate-intensity PA (MPA) and time sedentary, and b) the accelerometer cutpoint that best represents the threshold for health-promoting VPA in youth.

Methods: Data from the International Children's Accelerometry Database (ICAD) were analyzed in 2015. The relationship between cardiometabolic biomarkers and 4 categories of VPA estimated via 3 sets of cutpoints were examined using isotemporal substitution quantile regression modeling at the 10th, 25th, 50th, 75th, and 90th percentile of the distribution of each biomarker, separately. Age, sex, accelerometer wear time, sedentary time, and MPA were controlled for while allowing substitution for light-intensity PA. Data from 11,588 youth (4-18yrs) from 11 ICAD studies (collected 1998-2009) were analyzed.

Results: Only 32 of 360 significant associations were observed. Significant, negative relationships were observed for VPA with waist circumference and insulin. Replacing light intensity PA with VPA (corresponding to at the 25th to 90th percentiles of VPA) was associated with a .67 (-1.33, -0.01; P = .048) to 7.30cm (-11.01, -3.58; P < .001) lower waist circumference using Evenson and ICAD cutpoints (i.e., higher CPM). VPA levels were associated with 12.60 (-21.28, -3.92; P = .004) to 27.03 pmol/l (-45.03, -9.03; P = .003) lower insulin levels at the 75th to 90th percentiles using Evenson and ICAD cutpoints when substituted for light PA.

Conclusions: Substituting light PA with VPA was inversely associated with waist circumference and insulin. However, VPA was inconsistently related to the remaining biomarkers after controlling for time sedentary and MPA.

Keywords: Movement; cardiometabolic; adiposity; insulin.

1 INTRODUCTION

2 Emerging research utilizing international samples (7, 17) has indicated that many children
3 globally are spending an insufficient amount of time engaging in physical activity (PA) and an
4 excessive amount of time engaging in sedentary behaviors. Engaging in international guideline
5 recommendations (38) of 60 minutes per day (min/day) of moderate-to-vigorous physical
6 activity (MVPA) is inversely associated with biomarkers of cardiometabolic health (13, 25)
7 including lower rates of obesity (17) independent of time spent sedentary. While the benefits of
8 MVPA are well established cross-sectionally (7) and longitudinally (6), few studies of PA in
9 youth have examined the contribution of specific intensities to the association, despite a growing
10 body of literature that suggests that vigorous-intensity physical activity (VPA) may be more
11 important for the prevention and amelioration of cardiometabolic risk factors (13, 39). A small
12 number of studies have employed an objective measure of PA to examine associations with
13 cardiometabolic biomarkers. These studies suggest that VPA is independently associated with
14 cardiorespiratory fitness (positive) (23), BMI (negative) (17), adiposity (negative) (32), HDL
15 cholesterol (positive) (22), fasting glucose (negative) (31), and fasting insulin (negative) (1).
16 However, an extensive examination of the literature suggests that the relationship between VPA
17 and cardiometabolic biomarkers is inconsistent, potentially due to small samples, definition of
18 VPA, and other methodological limitations (11).

19

20 Complicating examinations of the relations between VPA and cardiometabolic biomarkers is an
21 issue of measurement of VPA, or more specifically, the threshold for which VPA occurs. While
22 imperfect, accelerometers are still considered one of the best objective measures available for
23 epidemiological studies of PA (5, 28), but the processing of data generated by accelerometers

24 (e.g., “counts”) lacks uniformity or consistency across studies (18), which can lead to
25 misclassification of exercise intensity (12) and/or lack of comparability across studies (3, 4).
26 Since the choice of cutpoint is a de facto selection of an intensity threshold with all other sources
27 of variability held constant (e.g., monitor brand, epoch), and no standard exists for the VPA
28 cutpoint, it is imperative to consider a range of accelerometer cutpoints for VPA if the
29 relationship between VPA and cardiometabolic biomarkers is to be studied.

30

31 The benefits of MVPA in youth are well established, but little research has been conducted to
32 examine the contribution of PA intensity in cardiometabolic health in youth. Therefore, the
33 objective of the present investigation was to determine, a) the associations between VPA and
34 cardiometabolic biomarkers independent of moderate physical activity (MPA) and sedentary
35 time, and b) the accelerometer cutpoint that best represents the threshold for health-promoting
36 VPA in a diverse sample of youth.

37

38 **MATERIALS/SUBJECTS AND METHODS**

39 **Study Design**

40 Data were utilized from the International Children’s Accelerometry Database (ICAD,
41 <http://www.mrc-epid.cam.ac.uk/Research/Studies/>), which was established to pool data on PA
42 from studies in youth worldwide. A comprehensive description of the ICAD can be found
43 elsewhere (34). Briefly, in 2008 19 studies were identified from a PubMed search that used an
44 Actigraph (Actigraph, LLC, Pensacola, FL, USA) accelerometer and included a minimum of 400
45 participants aged 3 to 18 years. Six additional studies were identified through professional
46 colleagues, with 21 studies ultimately contributing data to the final database (7, 34). For the

47 current study, 11 studies were included (7, 34), which are presented in brief with the variables
48 each contributed in Table 1 [details of the Avon Longitudinal Study of Parents and Children
49 (ALSPAC) are available at www.bris.ac.uk/alspac and including the data that are available via a
50 fully searchable data dictionary ([http://www.bris.ac.uk/alspac/researchers/data-access/data-](http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary)
51 [dictionary](http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary))]. Ethical approval for the present study was attained from participating institutions,
52 and data-sharing agreements were established prior to contribution of data.

53

54 **Participants**

55 Data from 11,588 youth (4-18yrs), representing 11 studies from Brazil, Europe, and the United
56 States from the ICAD were analyzed. Data from studies conducted between 1998 and 2009 were
57 included in the present analyses if the dataset contained PA, age, sex, and at least one biomarker
58 of a cardiometabolic risk [defined as “A characteristic that is objectively measured and evaluated
59 as an indicator of normal biological processes, pathogenic processes, or pharmacologic responses
60 to a therapeutic intervention” (2)].

61

62 **Measurements**

63 *Physical activity.* A comprehensive description of the measurement of PA has been published
64 previously (34). ICAD data were reanalyzed to allow for comparability across studies by
65 aggregating data to a 60-second epoch. The criterion of 60 minutes of consecutive zeros was
66 utilized to designate non-wear time, with a tolerance for 2 minutes of nonzero epochs (35).
67 Participants with three or more days with 600 minutes of valid wear time were included in
68 analyses. VPA was defined by cutpoints from Pate (26), Evenson (8), and the ICAD workgroup
69 (7, 34). These cutpoints were selected because they represent the most generous, lowest

70 threshold defining VPA [Pate $\geq 3,365$ counts/min (CPM)], a medium threshold (Evenson \geq
71 4,012 CPM), to the most stringent, highest threshold for VPA (ICAD $\geq 6,000$ CPM).

72

73 *Cardiometabolic biomarkers.* Eight cardiometabolic biomarkers reflecting a diverse array of
74 health indices were collected, including; waist circumference [as a proxy for adiposity (30)];
75 systolic and diastolic blood pressure (hemodynamics); high-density lipoprotein cholesterol, low-
76 density lipoprotein cholesterol, fasting triglycerides (lipid metabolism); fasting glucose and
77 fasting insulin (glucose metabolism). Details of data collection procedures can be found
78 elsewhere (7, 34). Waist circumference (WC) was assessed midway between the lower rib
79 margin and the iliac crest using a metal tape (10), except in the NHANES (National Health and
80 Nutrition Examination Survey) where WC was measured just above the iliac crest at the mid-
81 axillary line using similar equipment (36). Resting blood pressure was measured using standard
82 procedures, reported previously (7). Markers of lipid and glucose metabolism were assessed
83 using standard clinical procedures described in detail elsewhere (36).

84

85 **Statistical Analysis**

86 Descriptive analyses of accelerometer-derived estimates of $\text{min}\cdot\text{day}^{-1}$ spent in sedentary, MPA,
87 and VPA were computed across all studies using three sets of cutpoints to define PA intensities.
88 To evaluate the cross-sectional association of cardiometabolic biomarkers and time spent in
89 VPA, a series of isothermal substitution quantile regression models were estimated for each set
90 of cutpoints separately (20, 21, 40). Quantile regression models were employed since biomarkers
91 are often non-normal in their distribution, and quantile regression models are not influenced by
92 normality and are free from distributional assumptions (19). Individual models for each

93 biomarker as the dependent variable were estimated. Time spent in VPA, defined by one of the 3
94 sets of cutpoints, separately, served as the primary independent variable. Because of its non-
95 normal distribution, $\text{min}\cdot\text{day}^{-1}$ spent in VPA was placed into 4 categories – none (0mins/d –
96 reference category), low (lower 33%), middle (middle 33%), and high (upper 33%) – based on
97 the distribution of VPA for each of the 3 sets of cutpoints. The relationship between cardio-
98 metabolic biomarkers and 4 categories of VPA min/d [none (0 min/d – reference category), low
99 (7.2_{Pate} , 4.0_{Evenson} , 1.5_{ICAD} min/d), medium (18.6_{Pate} , 11.0_{Evenson} , 3.5_{ICAD} min/d), and high (42.7_{Pate} ,
100 28.9_{Evenson} , 11.9_{ICAD} min/d)] estimated via 3 sets of cutpoints [Pate: sedentary = 0 - 152
101 counts/min (CPM), MPA = 1677 – 3364, VPA = $\geq 3,365$ CPM; Evenson: sedentary = 0 - 100
102 CPM, MPA = 2296 – 4011, VPA = $\geq 4,012$ CPM; and ICAD: sedentary = 0 - 100 CPM, MPA =
103 3000 – 6000, VPA = $\geq 6,001$ CPM] —were examined using isotemporal substitution quantile
104 regression modeled at the 10th, 25th, 50th, 75th, and 90th percentiles of the distribution of each
105 biomarker. Included in each model were age (years), sex, average total daily wear time, and
106 $\text{min}\cdot\text{day}^{-1}$ in sedentary and MPA distilled using the corresponding cutpoint for VPA. Since light
107 PA (LPA) was the only intensity excluded from the models, all estimates are interpreted as
108 substituting “x” amount of LPA with VPA. Separate models were estimated for each study and
109 for each set of cutpoints used to define VPA within each study. An example of the modeling
110 approach is: insulin serving as the dependent variable, with 3 separate models using VPA levels
111 (i.e., low, middle and high, with no VPA as the referent group) reduced with each of the sets of
112 cutpoints for each study, run separately. Statistical significance was set at $P = .05$.

113

114 Meta-analytical techniques were used to combine the quantile regression model coefficients and
115 standard errors for each biomarker across the 11 studies for each of the 3 sets of cutpoints,

116 separately. Random effects inverse variance weighting was used to pool effects across studies
117 and within study for each set of cutpoints. The study served as the unit of analysis for each
118 quantile and category of VPA. For instance, the VPA estimates representing the lowest 33rd of
119 the distribution of VPA regressed on the 10th quantile of insulin were combined across all studies
120 for a given biomarker. All quantile regression analyses were conducted in 2015 using Stata
121 (v.13.0, College Station, TX) and all meta-analytic analyses were conducted using
122 Comprehensive Meta-Analysis (v2.2, Englewood, NJ).

123

124 **RESULTS**

125 Descriptive information for each study is presented in Table 2. The average amount of VPA
126 $\text{min}\cdot\text{day}^{-1}$ for each set of cutpoints (highest to lowest) by tertile ranged from 1.5 to 7.2 min/day
127 for the lowest tertile, the medium tertile 3.5 to 18.6 min/day, and the highest tertile 11.9 to 42.7
128 min/day. The results of the pooled meta-analytic effects for each quantile and level of VPA
129 across each cardiometabolic biomarker are presented in Table 3.

130

131 **Relationship of volume of VPA with cardiometabolic biomarkers**

132 Substituting LPA with VPA was inconsistently related to systolic/diastolic blood pressure,
133 fasting triglycerides, HDL, or LDL after controlling for time sedentary and MPA at all tertiles of
134 VPA volume, with only 32 of a possible 360 associations statistically significant ($P < .05$).
135 Independent of $\text{min}\cdot\text{day}^{-1}$ spent sedentary and in MPA, substituting LPA with VPA was
136 associated with a .67 to 7.30 cm smaller waist circumference at the 50th to 90th percentiles.
137 Relationships were observed for all three tertiles of VPA, but relationships at the lowest tertile of
138 VPA volume were significant at only the highest cutpoint value (i.e., ICAD). Substituting LPA

139 with VPA was associated with 12.6 to 27.0 pmol/l lower insulin values at the 75th to 90th
140 percentiles. Relationships were observed for all three tertiles of VPA, but relationships at the
141 lowest tertile of VPA were significant at only the highest tertiles of VPA volume for the highest
142 cutpoint value (i.e., ICAD).

143

144 **Influence of cutpoint**

145 Independent of min·day⁻¹ spent sedentary and in MPA, substituting LPA with the high volume of
146 VPA defined via Pate cutpoints was associated with a smaller waist circumference only at the
147 90th percentile. For VPA determined via Evenson cutpoints, substituting LPA for medium and
148 high VPA levels were associated with a smaller waist circumference at the 25th to 90th centiles.
149 Substituting LPA with the lowest, medium, and highest volumes of VPA reduced via ICAD
150 cutpoints was associated with a smaller waist circumference at the 50th to 90th, the 75th and 90th,
151 and the 25th to 90th, respectively. Across all other biomarkers (i.e., SBP, DBP, HDL, LDL,
152 glucose, and triglycerides), no consistent associations or patterns were observed, with only 9
153 significant associations observed from a possible 270 tested (<5%; see Figure).

154

155 **DISCUSSION**

156 The present study is the first of this scope (e.g., sample size, diversity of national origin) to
157 examine the relationship between VPA and cardiometabolic biomarkers in youth. The results are
158 consistent with previous studies using more homogeneous samples, such as Carson et al. (6)
159 where no association was found between diastolic blood pressure and VPA, but a significant
160 negative association was reported between waist circumference and VPA in children of the 2nd
161 and 3rd quartiles (relative to the 1st). The more nuanced analyses presented here, taken with those

162 of Carson et al. (6), provide additional insight into the complex relationship between VPA and
163 cardiometabolic biomarkers (11, 25). The results suggest that substituting modest amounts of
164 LPA for VPA may have cardiometabolic benefits above and beyond those conveyed by MPA
165 and avoidance of sedentary behavior (24). Of potentially greater importance, the current results
166 suggest that these health supportive associations are most pronounced in those who have
167 undesirable levels of these biomarkers, specifically those with relatively large waist
168 circumference or fasting insulin levels. If these relationships are found to be robust in
169 longitudinal and experimental studies, then a specific frequency and duration of VPA could be
170 incorporated as a distinct component of a PA “prescription” for youth (24). However, it must be
171 noted that VPA was independently associated with only two of the markers examined. Therefore,
172 while VPA may relay meaningful health benefits, the number of markers exhibiting those
173 benefits may be few relative to less intense movement.

174

175 These results, taken with a growing body of literature demonstrating the independent health
176 benefits of VPA for youth (6, 11, 14, 16, 17, 23, 24, 37), support the assertion that this intensity
177 should be considered when setting policy recommendations for PA of youth. For example, it has
178 been shown previously that as little as 9 (15) to 14 minutes (17) of VPA per day is associated
179 with less adiposity in Canadian (15) and multinational samples of youth (17). These previous
180 findings, derived from independent samples, are consistent with the present findings showing an
181 association of substituting 11.9 to 42.7 min/day of LPA for VPA. While this is a considerable
182 range, with the top end (42.7 min/day) potentially impractical, consistent benefits were seen for
183 VPA defined by the ICAD cutpoints, which even in the high volume category represented 11.9
184 min/day of VPA, is potentially achievable for most youth (39). Therefore, the present findings

185 suggest a modest duration (e.g., approximately 10 min) of high intensity PA may be related to
186 health benefits in youth who exhibit undesirable levels of insulin or waist circumference.

187

188 While the present study has a number of strengths, including an objective measure of PA, a large
189 sample size, a diverse and international sample, and an advanced analytical approach, the present
190 results should be considered in light of a number of limitations. First, all data were cross-
191 sectional in nature, therefore causality cannot be assumed. For example, it is possible that
192 children with smaller waist circumference are more vigorously active because it is less
193 cumbersome for them to do so. However, the nature of our analyses, which examined the
194 relationship of VPA and waist circumference at different quantiles of waist circumference, is less
195 supportive of this possibility. Second, while these cross-sectional results are supportive of VPA
196 specific PA recommendations for youth, it is unknown if changes in youth VPA levels will result
197 in meaningful changes in diastolic blood pressure, HDL, cholesterol, insulin or adiposity. While
198 a recent study is supportive of the latter three (29), the literature is mixed on the relationship
199 between increased VPA and blood pressure (9, 11, 27, 33), and very few studies have examined
200 the responsiveness of insulin or other markers of glucose metabolism (11, 13). Third, the
201 database we utilized lacks standardized dietary data or genetic data that might confound the
202 observed relationships. For example, children with higher levels of VPA may consume fewer
203 calories, or possess a genetic make-up supportive of a positive biomarker profile. This possibility
204 cannot be ruled out using the currently available data. Despite these limitations, this study
205 represents one of the largest to date that examined VPA in relation to cardiometabolic
206 biomarkers in youth.

207

208 In summary, the present results suggest few significant or clinically meaningful associations
209 between VPA and most cardiometabolic biomarkers studied in youth, but health promoting
210 associations were observed between VPA and select cardiometabolic biomarkers (i.e., insulin,
211 waist circumference), with the associations observed at higher levels of the biomarkers and
212 higher volumes of VPA. As such, VPA may have unique metabolic health benefits beyond those
213 conveyed by MPA or minimizing time spent sedentary. The present results also suggest that
214 higher VPA cutpoints represent an intensity that is associated with healthier insulin levels and
215 waist circumference. Future longitudinal and intervention studies are needed to determine the
216 temporal relationship between these variables, the modifiability of VPA, and the effect of
217 increased VPA on biomarkers in youth. If these results are indeed robust, then a less time
218 consuming, more intense dose of PA may be a viable option for youth seeking to achieve or
219 maintain cardiovascular health.

ACKNOWLEDGEMENTS

The results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation, and the results of the present study do not constitute endorsement by ACSM. We would like to thank all participants and funders of the original studies that contributed data to ICAD. We also gratefully acknowledge the contribution of Professor Chris Riddoch, Professor Ken Judge, and Dr. Pippa Griew to the development of ICAD.

ICAD COLLABORATORS

The ICAD Collaborators include: Prof LB Andersen, University of Southern Denmark, Odense, Denmark (Copenhagen School Child Intervention Study (CoSCIS)); Prof S Anderssen, Norwegian School for Sport Science, Oslo, Norway (European Youth Heart Study (EYHS), Norway); Prof G Cardon, Department of Movement and Sports Sciences, Ghent University, Belgium (Belgium Pre-School Study); Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Hyattsville, MD USA (National Health and Nutrition Examination Survey (NHANES)); Prof A Cooper, Centre for Exercise, Nutrition and Health Sciences, University of Bristol, UK (Personal and Environmental Associations with Children's Health (PEACH)); Dr. R Davey, Centre for Research and Action in Public Health, University of Canberra, Australia (Children's Health and Activity Monitoring for Schools (CHAMPS)); Prof U Ekelund, Norwegian School of Sport Sciences, Oslo, Norway & MRC Epidemiology Unit, University of Cambridge, UK; Dr. DW Esliger, School of Exercise and Health Sciences, Loughborough University, UK; Dr. K Froberg, University of Southern Denmark, Odense, Denmark (European Youth Heart Study (EYHS), Denmark); Dr. P Hallal, Postgraduate Program in Epidemiology, Federal University of Pelotas, Brazil (1993 Pelotas Birth Cohort); Prof KF Janz, Department of Health and Human Physiology, Department of Epidemiology, University of Iowa, Iowa City, US (Iowa Bone Development Study); Dr. K Kordas, School of Social and Community Medicine, University of Bristol, UK (Avon Longitudinal Study of Parents and Children (ALSPAC)); Dr. S Kriemler, Institute of Social and Preventive Medicine, University of Zürich, Switzerland (Kinder-Sportstudie (KISS)); Dr. A Page, Centre for Exercise, Nutrition and Health Sciences, University of Bristol, UK; Prof R Pate, Department of Exercise Science, University of South Carolina, Columbia, US (Physical Activity in Pre-school Children (CHAMPS-US) and Project Trial of Activity for Adolescent Girls (Project TAAG)); Dr. JJ Puder, Service of Endocrinology, Diabetes and Metabolism, Centre Hospitalier Universitaire Vaudois, University of Lausanne, Switzerland (Ballabeina Study); Prof J Reilly, Physical Activity for Health Group, School of Psychological Sciences and Health, University of Strathclyde, Glasgow, UK (Movement and Activity Glasgow Intervention in Children (MAGIC)); Prof J Salmon, School of Exercise and Nutrition Sciences, Deakin University, Melbourne, Australia (Children Living in Active Neighbourhoods (CLAN)); Prof LB Sardinha, Exercise and Health Laboratory, Faculty of Human Movement, Technical University of Lisbon, Portugal (European Youth Heart Study (EYHS), Portugal); Dr. LB Sherar, School of Sports, Exercise and Health Sciences, Loughborough University, UK; Dr. A Timperio, Centre for Physical Activity and Nutrition Research, Deakin University Melbourne, Australia (Healthy Eating and Play Study (HEAPS)); Dr. EMF van Sluijs, MRC Epidemiology Unit, University of Cambridge, UK (Sport, Physical activity and Eating behaviour: Environmental Determinants in Young people (SPEEDY)).

Funding source: The pooling of the data was funded through a grant from the National Prevention Research Initiative (Grant Number: G0701877) (<http://www.mrc.ac.uk/research/initiatives/national-prevention-research-initiative-npri/>). The funding partners relevant to this award are: British Heart Foundation; Cancer Research UK; Department of Health; Diabetes UK; Economic and Social Research Council; Medical Research Council; Research and Development Office for the Northern Ireland Health and Social Services; Chief Scientist Office; Scottish Executive Health Department; The Stroke Association; Welsh Assembly Government and World Cancer Research Fund. This work was additionally supported by the Medical Research Council [MC_UU_12015/3; MC_UU_12015/7], Bristol University, Loughborough University, and Norwegian School of Sport Sciences.

Financial disclosure: The authors have no financial relationships relevant to this article to disclose.

Conflict of interest: The authors have no conflicts of interest to disclose.

REFERENCES

1. Alhassan S, Robinson TN. Objectively measured physical activity and cardiovascular disease risk factors in African American girls. *Ethn Dis*. 2008;18(4):421-6.
2. Biomarkers Definitions Working G. Biomarkers and surrogate endpoints: preferred definitions and conceptual framework. *Clin Pharmacol Ther*. 2001;69(3):89-95.
3. Bornstein DB, Beets MW, Byun W, Welk G, Bottai M, Dowda M, et al. Equating accelerometer estimates of moderate-to-vigorous physical activity: In search of the Rosetta Stone. *J Sci Med Sport*. 2011;14(5):404-10.
4. Brazendale K, Beets MW, Bornstein DB, Moore JB, Pate RR, Weaver RG, et al. Equating accelerometer estimates among youth: The Rosetta Stone 2. *J Sci Med Sport*. 2016;19(3):242-9.
5. Butte NF, Ekelund U, Westerterp KR. Assessing Physical Activity Using Wearable Monitors: Measures of Physical Activity. *Med Sci Sports Exerc*. 2012;44(1S):S5-S12.
6. Carson V, Rinaldi RL, Torrance B, Maximova K, Ball GD, Majumdar SR, et al. Vigorous physical activity and longitudinal associations with cardiometabolic risk factors in youth. *Int J Obes (Lond)*. 2014;38(1):16-21.
7. Ekelund U, Luan Ja, Sherar LB, Esliger DW, Griew P, Cooper A. Moderate to Vigorous Physical Activity and Sedentary Time and Cardiometabolic Risk Factors in Children and Adolescents. *JAMA*. 2012;307(7):704-12.
8. Evenson KR, Catellier DJ, Gill K, Ondrak KS, McMurray RG. Calibration of two objective measures of physical activity for children. *J Sports Sci*. 2008;26(14):1557-65.
9. Farpour-Lambert NJ, Aggoun Y, Marchand LM, Martin XE, Herrmann FR, Beghetti M. Physical activity reduces systemic blood pressure and improves early markers of atherosclerosis in pre-pubertal obese children. *J Am Coll Cardiol*. 2009;54(25):2396-406.
10. Golding J, Pembrey M, Jones R, Team AS. ALSPAC--the Avon Longitudinal Study of Parents and Children. I. Study methodology. *Paediatr Perinat Epidemiol*. 2001;15(1):74-87.
11. Gralla MH, McDonald SM, Breneman C, Beets MW, Moore JB. Associations of Objectively Measured Vigorous Physical Activity With Body Composition, Cardiorespiratory Fitness, and Cardiometabolic Health in Youth: A Review. *Am J Lifestyle Med*. 2016.
12. Guinhouya CB, Lemdani M, Vilhelm C, Durocher A, Hubert H. Actigraph-defined moderate-to-vigorous physical activity cut-off points among children: statistical and biobehavioural relevance. *Acta Paediatrica*. 2009;98(4):708-14.

13. Gutin B, Owens S. The influence of physical activity on cardiometabolic biomarkers in youths: a review. *Pediatr Exerc Sci*. 2011;23(2):169-85.
14. Gutin B, Yin Z, Humphries MC, Barbeau P. Relations of moderate and vigorous physical activity to fitness and fatness in adolescents. *Am J Clin Nutr*. 2005;81(4):746-50.
15. Hay J, Maximova K, Durksen A, Carson V, Rinaldi RL, Torrance B, et al. Physical activity intensity and cardiometabolic risk in youth. *Arch Pediatr Adolesc Med*. 2012;166(11):1022-9.
16. Irving HM, Adlaf EM, Allison KR, Paglia A, Dwyer JJM, Goodman J. Trends in vigorous physical activity participation among Ontario adolescents, 1997-2001. *Can J Public Health*. 2003;94(4):272-4.
17. Katzmarzyk PT, Barreira TV, Broyles ST, Champagne CM, Chaput JP, Fogelholm M, et al. Physical Activity, Sedentary Time, and Obesity in an International Sample of Children. *Med Sci Sports Exerc*. 2015;47(10):2062-9.
18. Kim Y, Beets MW, Welk GJ. Everything you wanted to know about selecting the "right" Actigraph accelerometer cut-points for youth, but...: a systematic review. *J Sci Med Sport*. 2012;15(4):311-21.
19. Koenker R, Bassett G. Regression Quantiles. *Econometrica*. 1978;46(1):33-50.
20. Koenker R, Hallock KF. Quantile Regression. *J Econ Perspect*. 2001;15(4):143-56.
21. Koenker R, Xiao ZJ. Inference on the quantile regression process. *Econometrica*. 2002;70(4):1583-612.
22. Martinez-Gomez D, Eisenmann JC, Moya JM, Gomez-Martinez S, Marcos A, Veiga OL. The role of physical activity and fitness on the metabolic syndrome in adolescents: effect of different scores. The AFINOS Study. *J Physiol Biochem*. 2009;65(3):277-89.
23. Moore JB, Beets MW, Barr-Anderson DJ, Evenson KR. Sedentary time and vigorous physical activity are independently associated with cardiorespiratory fitness in middle school youth. *J Sports Sci*. 2013;31(14):1520-5.
24. Owens S, Galloway R, Gutin B. The Case for Vigorous Physical Activity in Youth. *Am J Lifestyle Med*. 2015.
25. Owens S, Gutin B. Physical Activity and Cardiometabolic Biomarkers in Youths: A 2013 Update. *Curr Cardiovasc Risk Rep*. 2014;8(2):1-8.
26. Pate RR, Almeida MJ, McIver KL, Pfeiffer KA, Dowda M. Validation and calibration of an accelerometer in preschool children. *Obesity (Silver Spring)*. 2006;14(11):2000-6.

27. Platat C, Wagner A, Klumpp T, Schweitzer B, Simon C. Relationships of physical activity with metabolic syndrome features and low-grade inflammation in adolescents. *Diabetologia*. 2006;49(9):2078-85.
28. Rachele JN, McPhail SM, Washington TL, Cuddihy TF. Practical physical activity measurement in youth: a review of contemporary approaches. *World J Pediatr*. 2012;8(3):207-16.
29. Racil G, Ben Ounis O, Hammouda O, Kallel A, Zouhal H, Chamari K, et al. Effects of high vs. moderate exercise intensity during interval training on lipids and adiponectin levels in obese young females. *Eur J Appl Physiol*. 2013;113(10):2531-40.
30. Reilly JJ, Kelly J, Wilson DC. Accuracy of simple clinical and epidemiological definitions of childhood obesity: systematic review and evidence appraisal. *Obes Rev*. 2010;11(9):645-55.
31. Rizzo NS, Ruiz JR, Oja L, Veidebaum T, Sjostrom M. Associations between physical activity, body fat, and insulin resistance (homeostasis model assessment) in adolescents: the European Youth Heart Study. *Am J Clin Nutr*. 2008;87(3):586-92.
32. Ruiz JR, Rizzo NS, Hurtig-Wennlof A, Ortega FB, Warnberg J, Sjostrom M. Relations of total physical activity and intensity to fitness and fatness in children: the European Youth Heart Study. *Am J Clin Nutr*. 2006;84(2):299-303.
33. Schmitz KH, Jacobs DR, Jr., Hong CP, Steinberger J, Moran A, Sinaiko AR. Association of physical activity with insulin sensitivity in children. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity*. 2002;26(10):1310-6.
34. Sherar LB, Griew P, Esliger DW, Cooper AR, Ekelund U, Judge K, et al. International Children's Accelerometry Database (ICAD): design and methods. *BMC Public Health*. 2011;11:485.
35. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc*. 2008;40(1):181-8.
36. US Department of Health and Human Services. Anthropometry and physical activity monitor procedures manual. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics; 2005. p. 132.
37. Vankim NA, Nelson TF. Vigorous physical activity, mental health, perceived stress, and socializing among college students. *Am J Health Promot*. 2013;28(1):7-15.
38. World Health Organization. Global Recommendations on Physical Activity for Health. Geneva, Switzerland: WHO Press, 2010.

39. Yin Z, Moore JB, Johnson MH, Vernon MM, Gutin B. The impact of a 3-year after-school obesity prevention program in elementary school children. *Childhood Obesity*. 2012;8(1):60-70.
40. Yu KM, Lu ZD, Stander J. Quantile regression: applications and current research areas. *J Roy Stat Soc D-Stat*. 2003;52:331-50.